

31
 $\frac{8}{4}$ $\frac{1}{13}$



Henry Field.

Dv 1010⁶ 1789

UNIVERSITÄTSBIBLIOTHEK
— Mathematische Nat. —
GUL. STR. 105
V 1904

COLLEGE OF PHYSICIANS.

Let the Book entitled THE PHARMACOPOEIA OF
THE ROYAL COLLEGE OF PHYSICIANS OF LONDON,
be printed.

Censors Board, October the 5th, 1787.

GEORGE BAKER, *President.*

HENRY REVELL REYNOLDS,
JAMES HERVEY,
JAMES ROBERTSON,
GEORGE FORDYCE, } *Censors.*

A
NEW TRANSLATION
OF THE
PHARMACOPOEIA
OF THE
ROYAL COLLEGE OF PHYSICIANS

OF
L O N D O N,
OF THE YEAR 1787;

WITH NOTES,
CRITICAL AND EXPLANATORY;
DOSES OF THE SEVERAL PREPARATIONS;

LIKEWISE

A TABLE OF THE QUANTITIES OF OPIUM AND QUICKSILVER
IN THE COMPOUND MEDICINES WHICH CONTAIN THEM,

AND A

LIST OF THE NEW NAMES;

TOGETHER WITH


LATIN AND ENGLISH INDEXES.

BY AN APOTHECARY.

L O N D O N:

PRINTED FOR J. JOHNSON, N^o 72, ST. PAUL'S
CHURCH-YARD.

M.DCC.LXXXIX.



TO
HIS MOST SACRED MAJESTY
GEORGE THE THIRD,
KING OF GREAT BRITAIN,
FRANCE, AND IRELAND,
DEFENDER OF THE FAITH,
DUKE OF BRUNSWIC AND LUNENBURGH,
ARCH-TREASURER AND ELECTOR
OF THE HOLY ROMAN EMPIRE, &c.

T H I S W O R K,
REVISED AND CORRECTED,

I S

D E D I C A T E D,
WITH THE GREATEST RESPECT AND REVERENCE,

B Y T H E

ROYAL COLLEGE OF PHYSICIANS OF LONDON,
KNOWING THAT IT WILL CLAIM HIS PATRONAGE
AND ACCEPTANCE, THE MORE IT IS ADAPTED
TO THE HEALTH OF THE PUBLIC.

UNIVERSITY OF CHALMERS
GEORGE THE THIRD
OF GREAT BRITAIN
AND IRELAND
MEMBER OF THE PARLIAM
OF GREAT BRITAIN AND IRELAND
BY J. H. B. W. O. R. E.
AND J. H. B. W. O. R. E.

PRINTED BY J. H. B. W. O. R. E.
IN THE CITY OF DUBLIN
BY J. H. B. W. O. R. E.
IN THE CITY OF DUBLIN
BY J. H. B. W. O. R. E.

P R E F A C E.

HALF a century is now nearly elapsed, since our predeceffors made a reform in our Dispensatory, which task we have now again undertaken ; this was executed in a manner which renders their industry and judgment very conspicuous and praise-worthy ; but although medicine has not perhaps kept pace in its progress with other useful arts, yet it has received assistances and improvements from various sources, which are neither few, nor of little value or estimation ; and more particularly from chemistry, which being lately studied with greater zeal and accuracy, and deeper judgment and attention, is well known to be much more, and better understood. Having therefore conceived, that our duty impelled us to bring both the substances which medicine employs, and their forms, anew' under our consideration, and weigh their importance, we did not think that we should fulfil this duty, or perform this work with propriety, without imbibing from the streams of modern chemistry whatever we saw useful in our undertaking, that so, from the addition of their

collected light, this part of medicine may shine with stronger, clearer, and brighter lustre. We had it principally in view to bring forward all those substances which chemistry furnishes to medicine, not only freed from impurity and error in the mode of preparation, but more perfect, and in a neater form, and in a more scientific order, than what the college had formerly used. Neither was all our labour and attention so employed and exhausted, in this very difficult part of our task, as to allow other parts to be neglected, or but just touched upon; on the contrary, the composition of each particular medicine was brought into view, and examined by itself with the most scrupulous attention, that if any thing was wanted it might be supplied, or if it contained any thing improper or unnecessary it might be cut off. Nor have we hesitated, in performing this, to throw out altogether such formulas as we judged useless, or to adopt and insert others which have more happy effects; so, however, as to avoid, on the one hand, any catching at novelty, and on the other, rejecting rashly what had been rendered habitual
by

by use. Particular care has been taken that there shall not remain any vestiges of medicines whose virtues are grounded on no better foundation than old womens tales. If, however, there may be found here and there remedies or prescriptions which may appear superfluous, or of no use, it is to be ascribed to our rather wishing to leave them to be corrected, or totally thrown out by our successors, than to obstinately set our faces against opinions, which, although not true, are nevertheless harmless. We have been particularly attentive to simplicity in prescription, and careful that medicines should not be mixed together, which would not readily coalesce, and which did not co-operate to the same purpose. Hence those pompous antidotes, composed of an enormous quantity of simples, huddled together without judgment, and for no reason, as being formed of substances collected together from all parts and all sources, of virtues totally repugnant, heaped together in an incongruous mass, are at length totally rejected. Even by this it appears, that we are no longer governed by a blind attachment to authority derived from inveterate habit, nor unbounded reverence for the antients. The

X P R E F A C E.

The terror and dread of poisons were impressed on the minds of the antients to an inconceivable degree, so as to produce a most painful anxiety, although it is now known that they were acquainted with but a very few. In our times their reception has been very dissimilar; and even medicine does not altogether look upon them as the most inveterate enemies of the human species, but, on the contrary, endeavours to bring them over to its party, and even hopes for aid from their alliance. Of these we have enumerated two or three, which we ourselves have made experiments with, being willing to adopt even others, if it should be found, by decisive experiment, that they also shall be proved capable of curing diseases; for it would be a degree of temerity extremely improper, for us to embrace those whose virtues are not sufficiently explored, as of experienced use, and advantage.

That deluge of unknown and new-coined names, which will appear as given to substances, already designed by others in general use, may induce some, at first sight, to consider us, in this respect, extremely culpable; since

since hardly any one does not wish rather to employ names for things with which he is already acquainted, than, as it were, learn a new language. But we beg that it may be considered, that these changes have the following objects: first, that those vain and absurd terms, taken from the works of old chemists, or otherwise applied, may, as much as can be, fall into disuse and oblivion; moreover, that the composition of a medicine may rather be known by its title, than an attempt be made to point out its virtues; and rather to show of what simple substances it consists, than in what diseases it may be employed, or to what part of the body its virtues are adapted; and lastly, that no medicine may be hidden by its name. As to the names we have given, according to our own fancy, to the three alkalis, one of which was in common use among the antients, and the other two but little altered from modern terms, they certainly carry upon their faces so much facility, and shorten so much the manner of expressing their compounds, that their introduction may at least be forgiven by physicians.

Never-

Nevertheless it is not to be denied, that so many novelties must bear hard upon those employed in pharmacy, while they begin to use them, and are not as yet versed in them; but this difficulty, however great it may be, unless we be mistaken, will be easily got the better of, and gradually go off of itself, giving way to an habit more convenient, more pleasant, and more useful.

We are by no means unconscious how great and difficult a work it would be, to produce a pure and perfect Pharmacopœia in all points — one which would please all men would be altogether an unhopeful attempt: nothing of this kind have we either undertaken, or professed to have performed; being ready to congratulate ourselves with the purest satisfaction if the work which we have set forth in these pages, with a wish to benefit the health of mankind, should in any degree produce this effect, alleviate the evils of disease, render the method of cure more conspicuous, and the restoration of health more expeditious.

ADVER-

ADVERTISEMENT.

BEFORE a criticism on any work of science is attempted, he who criticises ought to enquire into the means which the authors of the work had of being acquainted with the subject, in order that he may know whether he be a more competent judge of the symmetry of the different parts, and manner of constructing it, than the authors; and whether they meant to apply it to any purpose particularly, or whether there might be some supplemental effects, which the authors had not at all in view. Let us consider, therefore, with what view, and what lights, the College of Physicians of London have published their new Dispensatory. The best intelligence I have been able to acquire is this:—The College having resolved to publish a new edition of their Dispensatory, formed a committee of all the fellows, who met twice a week for two hours, to take into consideration what should be omitted, what should be added, and what should be amended of their former edition. They began by reading over their former edition, and receiving observations and propositions for the above purpose, which were argued and determined upon by a majority of the fellows present, when the matter appeared doubtful. This committee was attended by many of the first practitioners; and those who did not attend sent their proposals for alteration and amendment. The last edition being read through in the committee, and the several alterations and amendments approved of, being made to it, it was
printed

printed as a specimen. This specimen was sent to all the members of the college, to several apothecaries, and made as public as possible; with a request to all medical men to make observations, and propose amendments to the college in writing. A great many communications were made to the college in consequence, which were all read to the committee; and such of them as appeared eligible were admitted. A second specimen was formed, and made public, in the same manner, with the same request. This also produced many more communications, which were also read through; and some doubts arising about the propriety of certain chemical operations, a committee was formed to meet a committee of the Company of Apothecaries; which Company furnished this committee of the College of Physicians with their laboratory, materials, the assistance of their own committee, and their operator, Mr. Griffiths, who is generally considered as an excellent pharmaceutical chymist. With these assistances they went through their second specimen; and having made many corrections and amendments, it was left to the President and three of the Fellows to superintend the publication. After so much attention, and the assistance of men of such reputation for knowledge, not only in the practice of medicine, but also in the various auxiliary sciences, it behoves any one, who ventures to make remarks, to proceed with great diffidence, lest he should only convict himself of his own ignorance.

The remarks which accompany this translation are made partly to elucidate, as well as to consider the propriety of the several parts of this edition of the Dispensatory.

ORDER

ORDER OF THE CHAPTERS.

PONDERA, MENSURÆ, &c.	Page
MATERIA MEDICA	5
PRÆPARATIONES SIMPLICIORES	37
CONSERVÆ	44
SUCCI	47
EXTRACTA ET RESINÆ	49
OLEA EXPRESSA	57
OLEA DISTILLATA	58
SALES	65
MAGNESIA	91
PRÆPARATA E SULPHURE	93
PRÆPARATA EX ANTIMONIO	96
PRÆPARATUM EX ARGENTO	102
PRÆPARATA E FERRO	103
PRÆPARATA EX HYDRARGYRO	107
PRÆPARATA E PLUMBO	117
PRÆPARATUM E STANNO	119
PRÆPARATA E ZINCO	120
AQUÆ DISTILLATÆ	122
SPIRITUS DISTILLATI	127
DECOCTA ET INFUSA	138
VINA MEDICATA	151
TINCTURÆ	156
MISTURÆ	178
SYRUPI	183
MELLA MEDICATA	192
PULVERES	196
TROCHISCI	207
PILULÆ	211
ELECTUARIA	216
CONFECTIONES	219
AQUÆ MEDICATÆ	221
EMPLASTRA	224
UNGUENTA ET LINIMENTA	232
CERATA	243
EPITHEMATA	247

ORDER

ORDER OF THE CHAPTERS.

	Page
WEIGHTS, MEASURES, &c.	1
MATERIA MEDICA	5
EASY PREPARATIONS	37
CONSERVES	44
JUICES	47
EXTRACTS AND RESINS	49
EXPRESSED OILS	57
ESSENTIAL OILS	58
SALTS	65
MAGNESIA	91
PREPARATIONS OF SULPHUR	93
PREPARATIONS OF ANTIMONY	96
PREPARATION OF SILVER	102
PREPARATIONS OF IRON	103
PREPARATIONS OF QUICKSILVER	107
PREPARATIONS OF LEAD	117
PREPARATION OF TIN	119
PREPARATIONS OF ZINC	120
DISTILLED WATERS	122
DISTILLED SPIRITS	127
DECOCTIONS AND INFUSIONS	138
MEDICATED WINES	151
TINCTURES	156
MIXTURES	178
SYRUPS	183
MEDICATED HONIES	192
POWDERS	196
LOZENGES	207
PILLS	211
ELECTUARIES	216
CONFECTIONS	219
MEDICATED WATERS	221
PLAISTERS	224
OINTMENTS AND LINIMENTS	232
CERATES	243
POULTICES	247

THE
LONDON DISPENSATORY.

WEIGHTS AND MEASURES.

TWO kinds of weights are used in this kingdom, one for gold and silver, the other for other merchandises; the first we call Troy weight, the other Avoirdupois weight. The pounds are differently divided. The pound troy into twelve ounces, avoirdupois into sixteen. These weights differ both in their pounds and ounces; the avoirdupois pound is greater, the ounce less than the troy*. We use the troy pound, but we divide it differently from goldsmiths, to wit:

B

A pound

* The troy ounce contains 480 grains, the avoirdupois $437\frac{1}{2}$ grains.

A pound	} contains	{ twelve ounces.
An ounce		{ eight drachms.
A drachm		{ three scruples.
A scruple		{ twenty grains.

The measures are also of two kinds in this kingdom, ale measure and wine measure. We use wine measure, of which

A pint	} contains	{ sixteen ounces,
An ounce		{ eight drachms,

according to our division; and a gallon contains eight pints.

To avoid ambiguity, wherever we use the words ounces or drachms, we put w and m to distinguish weight and measure, in as much as an ounce and a drachm measure bear no general proportion to an ounce and a drachm weight.

* We consider mortars made of copper and brass as unfit, in most cases, to be employed
in

* Bell-metal mortars are those which have been in common use, and certainly can do no hurt in powdering dry roots, herbs, &c. where there is no moisture employed,
and

in the preparation of medicines. We also order that measures, funnels, and evaporating vessels, made of copper, lead, or any mixture of which either of these is an ingredient, should be carefully avoided.

B 2 We

and they are kept perfectly clean. Iron mortars are hard, but it is difficult to keep them clean from rust. For moist substances earthen-ware mortars, such as are made by Wedgwood are the best. Substances are more easily ground to powder by mills, which are of two kinds; one made of iron, as those employed for grinding coffee, pepper, &c. to use these the substances to be powdered should first be broken down into small pieces, the mill should be set wide, and the matter run through, by which means it is procured in gross powder. Then the mill should be set close, and the substance be run through so as to procure it fine. The other kind of mill is a stone roller running round a floor on a centre fixed at one end of the roller; the roller should be from six inches to a foot thick, and of four or five feet diameter, and turn upon an iron axis. It should be of some hard stone which is not calcareous. The floor should be made of the same stone, circular, and surrounded by a ledge to prevent the materials from being thrown off. Mills of either kind are not proper for grinding stony or metallic substances, for the purposes of medicine. When any vegetable or animal substance is to be powdered, it should be exposed to a heat not exceeding 100° until it is quite dry and crisp.

4 THE LONDON DISPENSATORY.

We use Fahrenheit's thermometer.

By boiling heat we are to be understood to mean from 200° to 212° .*

A gentle heat is from 90° to 100° according to us.

As often as we use the word specific gravity, we suppose the body heated to 55° .

* The heat of boiling water with the ordinary pressure of the atmosphere is 212° , but in taking a vessel off the fire and pouring the water upon drugs to make infusions, it falls to about 200° .

MATERIA

MATERIA MEDICA.

A.

Abrotonum, *the leaf.* Artemisia *Abrotanum.*

Linnæi Species

Plantarum.

Abfinthium mariti-
mum.

Artemisia *maritima,*

L. S. P.

Sea Wormwood,

the top.

B 3

Abfinthium,

It is very difficult to form a catalogue of Materia Medica, so as to enumerate those articles which every apothecary is expected to have ready in his shop. Some medicines have not only very great efficacy, but also great constancy in their operations. Others are less efficacious, and very inconstant; of these sometimes one set come into fashion, and sometimes another. Enumerations of Materia Medica have therefore been very various: and it has often happened that persons, many of them not in the medical

Abfinthium, vulgare. Artemifia *Abfynthium*,
 Wormwood, *the* L. S. P.
herb.

Acetofa

medical line, have stumbled upon some old Pharmacopœia, and found some medicine recommended in a particular difeafe, and having exhibited it, they have found or fup-
 pofed they have found it of efficacy in curing the patient, have vaunted it into practice, and continued it in vogue for fome time, until the fame inefficacy, or inconfancy, or fome detrimental quality, which threw it out of practice on a former occafion, again throws it into oblivion. Of this we have had late examples in cicuta, digitalis rubra, and various others, which, although fome of them have been introduced in the prefent Materia Medica of the College, will probably of themfelves fall out of fome future edition, and others of fimilar kinds be introduced.

Befides thefe there will always be changes of medicines having fimilar virtues, fince it is impoffible that an apothecary can keep all the fubftances which ever have been employed in medicine, of which even there is good evidence of their being efficacious. It is to be expected therefore, that one country will felect one fet of thefe, and another another fet, and with equal propriety, if they choofe fuch as will produce all the effects that medicine has hitherto been able to attain.

There is another great difficulty in forming a lift of Materia Medica arifing from the defcription the antients gave of plants, animals, and minerals, being fo defective

?

that

Acetosa pratensis, the Rumex Acetosa,
leaf. L. S. P.

Acidum vitriolicum.

Vitriolic acid.

The specific gravity is
to that of distilled
water as 1,850 to
1000.

B 4

Aconitum,

that we can hardly determine now to what substances the names they used are to be applied. As in white hellebore for instance, if we were to give the doses of the root which we call by that name, that they were accustomed to give, we should destroy almost all persons to whom it was exhibited. Modern writers on Natural History have given much more accurate descriptions; and Linnè has improved upon his predecessors to a very great degree, although he has no way arrived at perfection. There is also a ground on which his names are improper to be employed in medicine, which is that he has used such as are altogether incompatible with those employed by the greatest number of, and best medical authors, insomuch that their works, in as far as goes to the application of medicines, must become entirely useless. The College seem to have endeavoured to avoid this difficulty by using the old names, and affixing the synonyms of Linnè and other late Natural Historians. There is one thing however to be considered, that the names, which have been

8 THE LONDON DISPENSATORY.

Aconitum, <i>the herb.</i>	Aconitum <i>Napellus,</i> L. S. P.
Adeps <i>fuilla.</i>	
Hog's lard.	
Allium.	Allium <i>sativum,</i> L. S. P.
Garlic, <i>the root.</i>	
Aloë <i>Barbadensis,</i> focotorina.	Aloe <i>perfoliata,</i> L. S. P.
Althæa.	Althæa <i>officinalis,</i> L. S. P.
Marshmallow, <i>the</i> <i>root, the leaf.</i>	
Alumen.	Argilla <i>vitriolata.</i>
Allum.	
Ammoniacum, <i>the</i> <i>gum-resin.</i>	

Amygdala

in use for many of the articles of the *Materia Medica* have been very various in various practical authors, so that, perhaps, it would be better if for once they could be fixed so as not to admit of farther alteration; but for this purpose it would be necessary to form a glossary, in which the synonyms as used by authors should be added.

It is hardly practicable to give a translation of all the articles of the *Materia Medica* into English names, they have been so extremely vague, I have therefore in many cases retained the Latin name.

Amygdala amara, Bitter almond.	Amygdalus <i>communis</i> , L. S. P.
——— dulcis, Sweet almond, <i>the nut.</i>	
Anethum. Dill, <i>the seed.</i>	Anethum <i>graveolens</i> , L. S. P.
Angelica, <i>the root,</i> <i>stem, leaf, and seed.</i>	Angelica <i>Archangelic</i> , L. S. P.
Anisum, <i>the seed.</i>	Pimpinella <i>Anisum</i> , L. S. P.
Antimonium. Antimony.	Antimonium Sulphu- raturum.
Arabicum Gummi. Gum Arabic.	Mimosa <i>nilotica</i> , L. S. P.
Argentum. Silver.	
Arnica, <i>the herb,</i> <i>flower, and root.</i>	Arnica <i>montana</i> , L. S. P.
Arum, <i>the fresh root.</i>	Arum <i>maculatum</i> , L. S. P.
Aſa ſœtida, <i>the gum</i> <i>refin.</i>	Ferula <i>Aſa ſœtida</i> , L. S. P.
	Afarum,

10 THE LONDON DISPENSATORY.

Afarum, <i>the leaf.</i>	Afarum europæum, L. S. P.
Avena, <i>the oat.</i>	Avena sativa, L. S. P.
Aurantium hispalense. Orange tree, <i>the leaf,</i> <i>the flower, the juice</i> <i>and rind of the</i> <i>fruit.</i>	Citrus Aurantium, L. S. P.

B.

Balsamum Canadense.	Pinus Balsamea, L. S. P.
Copaivum.	Copaifera officinalis, L. S. P.
Peruvia- num.	Myroxylon peruife- rum. Linnæi Sup- plementum Planta- rum.
Toluta- num.	Toluiфера Balsamum, L. S. P.
Bardana, <i>the root.</i>	Arctium Lappa, L. S. P.
Barilla.	Natron impurum. Beccabunga,

Beccabunga, *the herb.* Veronica *Beccabunga*,
L. S. P.

Benzoë, *the resin.* Styrax *Benzoë.* Acta
Philosophica Lon-
dinenfis.

Biftorta, *the root.* Polygonum *Biftorta*,
L. S. P.

Bolus Gallicus,
Borax, Natron boracitatum.

C.

Calamus aromaticus, *the root.* Acorus *Calamus*,
L. S. P.

Calx, Lapis calcareus purus
Quick lime. recens ustus.

Camphora, Laurus *Camphora*.
Camphor. L. S. P.

Canella alba, *the bark.*

Cantharis, Meloë *vesicatorius*.
The Spanish Fly. Linnæi Systema
Naturæ.

Cardamine, *the flower.* Cardamine *pratensis*,
L. S. P.

Cardamomum.

12 THE LONDON DISPENSATORY.

Cardamomum minus.	Amomum repens.
Lesser Cardamum, <i>the</i>	Sonnerati iter.
<i>seed.</i>	
Carduus benedictus,	Centaurea benedicta,
<i>the herb.</i>	L. S. P.
Carica.	Ficus Carica, L. S. P.
Figs.	
Caruon.	Carum Carui, L. S. P.
Carraway, <i>the seed.</i>	
Caryophyllum aroma-	Caryophyllum aroma-
ticum.	ticum, L. S. P.
Cloves and their essen-	
tial oil.	
Caryophyllum ru-	Dianthus Caryophyl-
brum, <i>the flower.</i>	lus, L. S. P.
Cascarilla, <i>the bark.</i>	
Cassia fistularis, <i>the</i>	Cassia Fistula, L. S. P.
<i>fruit.</i>	
Castoreum Russicum.	
Russian Castor.	
Catechu, vulgo Terra	Mimosa Catechu,
Japonica.	L. Suppl. P.
Catechu, commonly	
called Japan earth,	

Centaureum

Centaureum minus. *Gentiana Centaurium,*
 Lesser Centaury, *the* L. S. P.
top.

Cera flava.
 Yellow wax.
 alba.

White wax.

Chamœmelum. *Anthemis nobilis,*
 Camomile, *the single* L. S. P.
flower.

Chelæ cancrorum. *Cancer Pagurus,*
 Crabs claws. L. S. N.

Cicuta. *Conium maculatum,*
 Hemlock, *the herb,* L. S. P.
flower, and seed.

Cinara. *Cynara Scolymus,*
 Artichoke, *the leaf.* L. S. P.

Cineres clavellati. *Kali impurum.*
 Pearl ashes.

Cinnamomum. *Laurus Cinnamomum,*
 Cinnamon, *the bark,* L. S. P.
and its essential oil.

Coccinella.

Cochineal.

Cochlearai

14 THE LONDON DISPENSATORY.

Cochlearia hortenſis. Cochlearia *officinalis*,
Garden Scurvy-graſs, L. S. P.
the herb.

Colchicum, *the fresh* Colchicum *autumnale*,
root. L. S. P.

Colocynthis. Cucumis *Colocynthis*,
Colocynthida, *the pith* L. S. P.
of the fruit.

Colomba.

Columba, *the root.*

Contraſyerva, *the root.* Dorſtenia *Contraſy-*
erva, L. S. P.

Corallium rubrum. Iſis *nobilis*, L. S. N.
Red Coral.

Coriandrum. Coriandrum *ſativum*,
Coriander, *the ſeed.* L. S. P.

Cornu cervi.

Hartſhorn.

Creta.

Chalk.

Crocus. Crocus *ſativus*,
Saffron, *the ſtigma of* L. S. P.
the flower.

Cubeba.

Cubeba.	Piper <i>Cubeba</i> , L. Suppl. P.
Cucumis agreffis. Wild cucumber, <i>the</i> <i>fresh fruit.</i>	Momordica <i>Elate-</i> <i>rium</i> , L. S. P.
Cuminum. Cumin, <i>the seed.</i>	Cuminum <i>Cyminum</i> , L. S. P.
Cuprum. Copper.	
Ærugo, Verdigris.	
Vitriolum cœru- leum, Blue Vitriol.	Cuprum vitriolatum.
Curcuma, <i>the</i> <i>root.</i>	Curcuma <i>longa</i> , L. S. P.
Cydonium malum. Quince, <i>the apple,</i> <i>and its seed.</i>	Pyrus <i>Cydonia</i> , L. S. P.
Cynosbatus. Hip, <i>the fruit.</i>	Rosa <i>canina</i> , L. S. P.

Daucus

D.

Daucus sylvestris.	Daucus <i>Carota</i> ,
Wild carrot, <i>the seed.</i>	L. S. P.
Digitalis.	Digitalis <i>purpurea</i> ,
Fox-glove, <i>the herb.</i>	L. S. P.

E.

Elemi, <i>a resin.</i>	Amyris <i>Elemifera</i> ,
	L. S. P.
Enula campana.	Enula <i>Helenium</i> ,
Elecampane, <i>the root.</i>	L. S. P.
Eryngium, <i>the root.</i>	Eryngium <i>maritimum</i> ,
	L. S. P.

F.

Ferrum.	
Iron.	
<i>Vitriolum Viride.</i>	Ferrum <i>vitriolatum</i> .
<i>Green Vitriol.</i>	
Filix.	Polypodium <i>Filix</i>
Fern, <i>the root.</i>	mas, L. S. P.
	Feniculum

Feniculum dulce. Anethum *Feniculum*,
 Sweet fennel, *the seed.* L. S. P.
 Fenum Græcum, Trigonella *Fenum*
the seed. *Græcum*, L. S. P.

G.

Galbanum, *the gum-* Bubon *Galbanum*,
resin. L. S. P.

Galla.

Gall.

Gambogia.

Gamboge, *a gum-*
resin.

Genista.

Spartium *scoparium*,
 L. S. P.

Broom, *the top, the*
seed.

Gentiana.

Gentiana lutea, L. S. P.

Gentian, *the root.*

Ginseng, *the root**

Panax quinquefolium,
 L. S. P.

C

Glycyrrhiza

* I should rather have expected this article in a Chinese Materia Medica.

Glycyrrhiza, Glycyrrhiza *glabra*,
Liquorice, *the root.* L. S. P.

Granatum. Punica *Granatum*,
Pomegranate, *the* L. S. P.

flower called Balau-
stium, and the rind
of the fruit.

Gratiola, *the herb.* Gratiola *officinalis*,
L. S. P.

Guaiacum, *the wood,* Guaiacum *officinale*,
bark, and gum- L. S. P.
resin.

H.

Helleboraster, *the* Helleborus *fætidus*,
leaf. L. S. P.

Helleborus *albus.* Veratrum *album*,
White hellebore, *the* L. S. P.

root.

niger. Helleborus *niger*,

Black hellebore, *the* L. S. P.
root.

Hordeum.

Hordeum.	Hordeum <i>distichon</i> ,
Barley, <i>the seed.</i>	L. S. P.
<i>Pearl Barley.</i>	
Hydrargyrus.	
Quicksilver.	
Cinnabaris.	Hydrargyrus sulphu-
Cinnabar.	ratus.
Hypericum, <i>the</i>	Hypericum <i>perfora-</i>
<i>flower.</i>	<i>tum</i> , L. S. P.

I.

Jalapium.
 Jalap, *the root.**

C 2

Ichthyocolla.

• This and the two following articles have no synonyms, which I observe happens likewise in several of the other articles of the *Materia Medica*. As in the works of the Linnæus's as well as of other authors several of these are described, it led me to enquire from what cause these synonyms were omitted. The most obvious one was, that the College were not satisfied with the identity of the species, and that it was better to give no specification of the plants and animals from whence the articles were procured than a doubtful one. On enquiry, I find there are several species of convolvulus in the botanic gardens

Ichthyocolla.	
Ifinglafs.	
Ipecacuanha.	
Ipecacuan, <i>the root.</i>	
Iris, <i>the root.</i>	Iris <i>florentina</i> , L. S. P.
Juglans.	Juglans <i>regia</i> , L. S. P.
Wallnut, <i>the unripe fruit.</i>	
Juniperus.	Juniperus <i>communis</i> ,
Juniper, <i>the berry, and top.</i>	L. S. P.

K.

Kino, <i>a resin.</i>	Gummi Gambiense.
-----------------------	------------------

Ladanium.

gardens of this country under the name of jalap, all of which have been imported as the true jalap plant, and that the evidence of the identity of the species is extremely deficient in all the other cases. There is also another reason, which in some instances I suppose may have had an effect: it is, that the same substance is not uncommonly the product of several vegetables, as manna, sugar, &c.

L.

Ladanum, <i>a resin.</i>	Cistus creticus, L. S. P.
Lavendula.	Lavandula Spica, L. S. P.
Lavander, <i>the flower.</i>	L. S. P.
Laurus.	Laurus nobilis, L. S. P.
Laurel, <i>the leaf, and berry.</i>	
Lignum Campe- chense.	Hæmatoxylum Cam- pechianum, L. S. P.
Logwood.	
Limon.	Citrus Medica, L. S. P.
Lemon, <i>the juice, the external rind, and the oil called essen- tial.</i>	
Linum,	Linum usitatissimum, L. S. P.
Flax, <i>the seed.</i>	
Lujula, <i>the leaf.</i>	Oxalis Acetocella, L. S. P.

C 3

Majorana.

M.

Majorana.	<i>Origanum Majorana,</i>
Marjoram, <i>the herb.</i>	L. S. P.
Malva.	<i>Malva sylvestris,</i>
Mallow, <i>the leaf, and</i> <i>flower.</i>	L. S. P.
Manna.	
Marrubium album.	<i>Marrubium vulgare,</i>
White hoarhound, <i>the</i> <i>herb.</i>	L. S. P.
Marum syriacum, <i>the</i> <i>herb.</i>	<i>Teucrium Marum,</i> L. S. P.
Mastiche.	<i>Pistacia Lentiscus,</i>
Mastich, <i>the resin.</i>	L. S. P.
Mel.	
Honey.	
Melissa.	<i>Melissa officinalis,</i>
Balm, <i>the herb.</i>	L. S. P.
Mentha piperitis.	<i>Mentha piperita,</i>
Peppermint, <i>the herb.</i>	L. S. P.
<i>sativa.</i>	<i>Mentha spicata.</i>
	Mint,

Mint, <i>the herb.</i>	Hudsoni Flora An- glica.
Mezereum, <i>the bark of the root.</i>	Daphne Mezereum, L. S. P.
Millepeda.	Oniscus <i>Asellus,</i>
Milleped.	L. S. P.
Morum.	Morus <i>nigra,</i> L. S. P.
Mulberry, <i>the fruit.</i>	
Mofchus.	
Musk.	
Myrrha, <i>a gum-resin.</i>	

N.

Nasturtium aquati- cum.	Sisymbrium <i>Nastur- tium aquaticum,</i>
Water-cress, <i>the fresh herb.</i>	L. S. P.
Nicotiana.	Nicotiana <i>Tabacum,</i>
Tobacco, <i>the leaf.</i>	L. S. P.
Nitrum.	Kali nitratum.
Nux moschata.	Myristica <i>Moschata,</i>
Nutmeg, <i>the essential oil, the expressed oil</i>	Acta Holmiensia.

C 4

commonly

*commonly called oil
of mace, and mace.*

O.

Olibanum, a gum- resin.	Juniperus lycia, L. S. P.
Oliva.	Olea europæa, L. S. P.
Olive, the oil.	
Opium.*	
Opopanax, a gum- resin.	Pastinaca Opopanax, L. S. P.
Origanum, the herb.	Origanum vulgare, L. S. P.
Ovum.	Ovum gallinaceum.
Egg.	

Papaver

* There has been no true botanical description of the species of poppy from which opium is made in Arabia. The most judicious of those who have observed the cultivation of opium of my acquaintance differ very much in opinion from Linnè, who probably never saw the plant. It is singular however that the College should not have followed the common opinion of authors. It is a mark of their care not to mislead. There has no writer given any tolerable account of the preparation of opium from the poppy.

P.

Papaver album.	Papaver <i>somniferum</i> ,
White poppy, <i>the</i> <i>head.*</i>	L. S. P.
erraticum.	<i>Rhæas</i> , L. S. P.
Corn poppy, <i>the</i> <i>flower.</i>	
Pareira brava, <i>the</i> <i>root.</i>	<i>Cissampelos pareira</i> ,
	L. S. P.
Parietaria, <i>the herb.</i>	<i>Parietaria officinalis</i> ,
	L. S. P.
Pentaphyllum, <i>the</i> <i>root.</i>	<i>Potentilla reptans</i> ,
	L. S. P.
Peruvianus cortex.	<i>Cinchona officinalis</i> ,
Peruvian Bark.	L. S. P.
Petroleum.	<i>Bitumen Petroleum</i> ,
	L. S. N.
Petrofelinum, <i>the root</i> ,	<i>Apium Petrofelinum</i> ,
<i>the seed.</i>	L. S. P.
Pimento, <i>the berry.</i>	<i>Myrtus Pimenta</i> ,
	L. S. P.
	Piper

* *Quandoque dormitat bonus Homerus!* It should have been the capsule. Caput is classical.

Piper Indicum.	Capficum <i>annuum</i> ,
Cayenne pepper, <i>the</i>	L. S. P.
<i>fruit, or rather the</i>	
<i>capsule.</i>	
longum.	Piper <i>longum</i> , L. S. P.
Long pepper, <i>the</i>	
<i>fruit.</i>	
nigrum.	Piper <i>nigrum</i> , L. S. P.
Black pepper, <i>the</i>	
<i>berry.</i>	
Pix Burgundica.*	
Burgundy pitch.	
liquida.	
Tar.	
Plumbum.	
Lead.	
Ceruffa,	
Ceruffe.	
Lithargyrus,	
Litharge.	
Minium.	

Prunum

* This is probably a mixture of different materials.

Prunum Gallicum.	Prunus domestica,
Prunes,	L. S. P.
fylvestre.	Prunus spinosa,
Sloe.	L. S. P.
Pulegium,	Mentha Pulegium,
Pennyroyal, <i>the herb,</i>	L. S. P.
<i>the flower.</i>	

Q.

Quassia, <i>the wood, the</i>	Quassia amara,
<i>bark, the root,</i>	L. S. P.
Quercus,	Quercus Robur,
Oak, <i>the bark,</i>	L. S. P.

R.

Raphanus rusticanus.	Cochlearia Armoracia,
Horse-radish, <i>the root.</i>	L. S. P.
Rhabarbarum.	Rheum palmatum,
Rhubarb, <i>the root.*</i>	L. S. P.

Ribes

* The College ought to have left the synonym undetermined; for the Tartars, although they principally take the rheum palmatum because its roots are large, yet frequently gather other species. And it is most probable that

Ribes nigrum.	Ribes <i>nigrum</i> ,
Black currant, <i>the</i>	L. S. P.
<i>fruit.</i>	
rubrum.	<i>rubrum</i> , L. S. P.
Red currant, <i>the fruit.</i>	
Ricinus, <i>the seed.</i>	Ricinus <i>communis</i> ,
Rosa damascena.	Rosa <i>centifolia</i> ,
Damask rose, <i>the</i>	L. S. P.
<i>flower leaf.</i>	
rubra.	Rosa <i>Gallica</i> , L. S. P.
Red rose, <i>the flower</i>	
<i>leaf.</i>	
Rosmarinus.	Rosmarinus <i>officinalis</i> ,
Rosemary, <i>the top, the</i>	L. S. P.
<i>flower.</i>	
Rubia.	Rubia <i>tinctorum</i> ,
Madder, <i>the root.</i>	L. S. P.
Rubus idæus.	Rubus <i>idæus</i> , L. S. P.
Raspberry, <i>the fruit.</i>	
	Ruta.

that the rheum palmatum is not the best, as there is a quantity selected for the use of the Imperial family of Russia, which is heavier, more compact, and less yellow than the roots which have been produced from the seeds of the rheum palmatum brought over by Dr. Mouncey.

Ruta. *Ruta graveolens,*
 Rue, *the herb.* L. S. P.

S.

Sabina. *Juniperus Sabina,*
 Savin, *the leaf.* L. S. P.

Saccharum non puri-
 ficatum.

Muscavada fugar.
 purificatum. Saccharum bis coc-
 tum.
 Double-refined fugar.

Sagapenum, a gum-
 resin.

Sal amarus. *Magnesia vitriolata.*
 Bitter or Epsom salt.

Sal ammoniacus. *Ammonia muriata.*

Sal ammoniac. *Natron muriatum.*

Sal muriaticus.
 Common salt.

Salvia. *Salvia officinalis,*
 Sage, *the leaf.* L. S. P.

Sambucus.

Sambucus.	Sambucus nigra,
Elder, <i>the interior</i>	L. S. P.
<i>bark,* the flower,</i>	
<i>and the berry.</i>	
Sanguis draconis.	
Dragon's blood, <i>a</i>	
<i>resin.</i>	
Santalum rubrum.	Pterocarpus <i>Santolinus</i> ,
Red Sanders, <i>the wood.</i>	L. Suppl. P.
Santonium, <i>the seed.</i>	Artemisia <i>Santonium</i> ,
	L. S. P.
Sapo.	Sapo ex oleo olivæ &
Soap.	natro confectus.
Sarcocolla, <i>a gum-</i>	
<i>resin.</i>	
Sarsaparilla, <i>the root.</i>	Smilax <i>Sarsaparilla</i> ,
	L. S. P.
Sassafras, <i>the wood, the</i>	Laurus <i>Sassafras</i> ,
<i>root, and the bark</i>	L. S. P.
<i>of the root.</i>	

Scammonium.

* Cortex interior is not Latin. It ought to have been liber.

Scammonium.	Convolvulus <i>Scamma-</i>
Scammony, a gum <i>resin.</i>	<i>nia</i> , L. S. P.
Scilla.	<i>Scilla maritima</i> ,
Squill, <i>the root.</i>	L. S. P.
Scordium, <i>the herb.</i>	<i>Teucrium Scordium</i> ,
	L. S. P.
Senna, <i>the leaf.</i>	<i>Cassia Senna</i> , L. S. P.
Seneka, <i>the root.</i>	<i>Polygala Senega</i> ,
	L. S. P.
Serpentaria Virginiana, <i>the root.</i>	<i>Aristolochia Serpen-</i> <i>taria</i> , L. S. P.
Sevum ovillum.	
Mutton fuet.	
Simarouba, <i>the bark.</i>	<i>Quassia Simarouba</i> ,
	L. Suppl. P.
Sinapi.	<i>Sinapis nigra</i> , L. S. P.
Mustard, <i>the seed.</i>	
Sium, <i>the herb.</i>	<i>Sium nodiflorum</i> ,
	L. S. P.
Spermaceti.	
Spigelia, <i>the root.</i>	<i>Spigelia marilandica</i> ,
	L. S. N.

Spina cervina, *the* Rhamnus catharticus,
berry. L. S. P.

Spiritus vinosus recti-
 ficatus.

Rectified spirit of wine,
 contains 95 parts of
 alcohol, and 5 parts
 of distilled water in
 100. Its specific
 gravity is to that of
 distilled water as
 835 to 1,000.

Spiritus vinosus te-
 nuior.

Weaker spirit of wine,
 contains 55 parts of
 alcohol, and 45 of
 distilled water in
 100. Its specific
 gravity is to that
 of distilled water as
 930 to 1,000.

Spongia.
 Sponge.

Spongia officinalis,
 L. S. N.

Stannum.

Stannum.

Tin.

Staphisagria, *the seed.* Delphinium *Staphisagria*, L. S. P.Styrax, *a resin.* Styrax *officinalis*,
L. S. P.

Succinum.

Amber.

Sulphur.

Sulphuris flores.

Flowers of sulphur.*

T.

Tamarindus. Tamarindus *Indica*,Tamarind, *the fruit.* L. S. P.Tanacetum. Tanacetum *vulgare*,Tansey, *the flower,*
the herb. L. S. P.Taraxacum. Leontodon *Taraxa-*Dandelion, *the root,*
the herb. cum, L. S. P.

D

Terebinthina.

* According to the usual form observed in this Materia Medica, Sulphuris Flores should have been in Italics; the properties not being altered by the sublimation.

Terebinthina vulgaris.

Common turpentine.

Chia.

Chia turpentine.

Testæ ostreorum.

Ostrea edulis, L. S. N.

Oyster-shells.

Thus.

Frankincense, a resin.

Tormentilla.

Tormentilla erecta,

Tormentil, the root.

L. S. P.

Tragacantha.

Astragalus Tragacan-

Tragacanth, a gum.

tha, L. S. P.

Trifolium paludosum,
the herb.

Menyanthes trifoliata,

L. S. P.

Triticum.

Triticum hybernum,

Wheat, the flower.

L. S. P.

starch.

Tuffilago, the herb.

Tuffilago Farfara,

L. S. P.

V.

Valeriana sylvestris.

Valeriana officinalis,

Wild Valerian, the
root.

L. S. P.

I

Viola.

Viola.	Viola odorata, L. S. P.
Violet, <i>the fresh</i> <i>flower.</i>	
Vitis.	Vitis vinifera, L. S. P.
The Vine.	
Uva passa.	
Raisins.	
Vinum.	
Wine.	
Tartarum.	Crude tartar.
Tartar.	
Tartari cryfalli.	Purified tartar.
Cryfalls of tartar.	
Acetum.	
Vinegar.	
Ulmus.	Ulmus campestris,
Elm, <i>the interior</i> <i>bark.</i>	L. S. P.
Urtica.	Urtica dioica, L. S. P.
The nettle, <i>the herb.</i>	
Uva urfi, <i>the leaf.</i>	Arbutus Uva urfi,
	L. S. P.

D 2

Zedoaria,

Z.

Zedoaria, <i>the root.</i>	Kaempferia <i>rotunda,</i> L. S. P.
Zincum.	
Zinc.	
Lapis calaminaris. Calamine.	Lapis calaminaris uf- tus.
Tutty.	
Tutty.	
Vitriolum album. White Vitriol.	Zincum vitriolatum.
Zingiber.	
Ginger, <i>the root.</i>	Amomum <i>Zingiber.</i>

P R Æ-

PRÆPARATIONES
SIMPLICIORES.

EASY PREPARATIONS.

QUORUNDAM AQUA NON SOLU-
BILIUM PRÆPARATIO.

PREPARATIONS OF SOME SUBSTAN-
CES WHICH ARE NOT SOLUBLE IN
WATER.

THEY are to be powdered in a mortar;
afterwards moistened with water, and
levigated upon a hard stone which is not cal-
careous, till they are reduced to a very fine
powder.* Dry this powder upon a mass of
chalk, covered with filtering paper, in a
warm, or at least dry place.

D 3

In

* The best stone to levigate these powders upon is red granite; or, if it can be procured, granite which is perfectly white. They might be made still finer by a second grinding on agate or jasper, well polished.

In this way are prepared,
 Antimony,
 Crabs claws,
 Coral,
 Chalk,*
 Calamine,
 Amber,
 Oyster-shells, first cleared from any dirt
 which may adhere to them.
 Tutty.

The Crabs claws should be broken first into
 small pieces, and washed with boiling water
 before they are levigated. †

Prepare verdigris in the same manner.

* This may be procured as well from manufactories,
 where it is called whitening.

† I suppose the College mean before they are reduced
 to a fine powder in the mortar.

ADIPIS

ADIPIS SUILLÆ SEVIQUE OVILLI
PRÆPARATIO.THE PREPARATION OF HOGS LARD
AND MUTTON SUET.

Cut them into small pieces, melt them in a moderate heat, then strain them from the membranes.

AMMONIACI PURIFICATIO.

THE PURIFICATION OF AMMONIAC.

Boil gum ammoniac, if it looks impure, in water till it becomes soft, then squeeze it through a hempen bag, and set it by till the resinous part subsides. Evaporate the water, and towards the end of the evaporation add the resinous part, and mix it with the gum.

Asa foetida and other gum-resins are to be purified in the same manner.

Any gum that melts easily, such as galbanum, may be purified by putting it into an

D 4

ox's

ox's bladder, and keeping it in warm water till it becomes so soft that it may be squeezed by a press through an hempen bag.*

CORNU CERVI USTIO.

THE BURNING OF HARTSHORN.

Burn pieces of hartshorn till they are quite white to the heart, then reduce them to a very fine powder.

HERBARUM ET FLORUM EXSICCATIO.

THE DRYING OF HERBS AND FLOWERS.

Strew them lightly, and dry them in a gentle heat. †

MELLIS

* It is much better to employ in medicine only such parcels of these resins and gum-resins as are naturally pure without any preparation.

† Do it in a heat between 90° and 100° of Fahrenheit's thermometer. This degree of heat is by much the best, preserving

MELLIS DESPUMATIO.

THE CLARIFICATION OF HONEY.

Melt honey in a water bath, and take off the scum.

MILLEPEDÆ PRÆPARATIO.

PREPARATION OF MILLEPEDES.

Put millepedes into a thin hempen bag, place them over hot weak spirits of wine, in a close vessel, that they may be killed and rendered brittle by the vapour.

PUL-

preserving even the essential oils more perfectly than in a lesser.

Where many medicines or a large quantity are required to be dried, a coale should be set in the middle of a room, the smoke being conveyed to the chimney by an iron pipe. Iron stands should be placed round the sides of the room. The substances to be dried should be spread in tin pans placed over one another on the stands, at about eight inches distance. The proper heat should be produced by the fire in the coale. Herbs dried in this manner retain their colour, their flavour, their taste, and their medicinal qualities much more perfectly.

PULPARUM EXTRACTIO.

THE EXTRACTION OF PULPS.

If pulpy fruits be unripe, or ripe and dry, fet them by in a moist place till they become soft; then squeeze the pulp through a hair sieve, boil them over a moderate fire, stirring them frequently, till they become of a proper consistence.*

Take out the pulp of cassia, and boil it to a proper thickness.

Press out the pulps of ripe and fresh fruits through a sieve, without boiling them.

SCILLÆ EXSICCATIO.

THE DRYING OF SQUILLS.

Take off the outer peel of the squill, cut it transversely into thin slices, and dry it in a gentle heat.

SPONGIÆ

* I am informed that this method of softening pulpy fruits was preferred to that of boiling them in water, from an experiment that had been made at Apothecary's Hall, which determined the Company to use this mode in preference.

SPONGIÆ USTIO.

THE BURNING OF SPONGE.

Cut a sponge in pieces, beat it till it is separated from the small stones which often adhere to it, then burn it in a close iron vessel until it becomes black and brittle; and lastly powder it very finely.

STYRACIS PURIFICATIO.

PURIFICATION OF STYRAX.

Dissolve styrax in rectified spirit of wine, strain it, then evaporate it in a gentle heat until it becomes of a proper consistence.

CON-

C O N S E R V Æ.

C O N S E R V E S.

CONSERVE of Lujula,
SEA WORMWOOD,
The RED ROSE,
The exterior Rind of the SE-
VILLE ORANGE.

TAKE off the foot-stalks from the leaves; take the flower-leaves, before they are thoroughly expanded, out of the flower-cup, take off the yellow points; rasp the external rind of the orange with a grater, beat them in a marble mortar with a wooden pestle, first the matter by itself, and then with three times its weight of double refined sugar, until they are mixed.

CON-

CONSERVA ARI.

CONSERVE OF ARUM.

Take of the root of fresh arum bruised half a
pound,

Double refined sugar one pound and
an half.

Beat them together in a mortar.

CONSERVA CYNOSBATI.

CONSERVE OF THE HIP.

Take of the pulp of fresh hips one pound,

Double refined sugar powdered w.
twenty ounces.

Mix them so as to form a conserve.

CONSERVA PRUNI SYLVESTRIS.

CONSERVE OF THE SLOE.

Put the sloes in water, then set them over
the fire to soften, taking care not to allow
them

46 THE LONDON DISPENSATORY.

them to burst; when they are soft enough take them out, press out the pulp, and mix it with three times its weight of double refined sugar, so as to make a conserve.

CONSERVA SCILLÆ.

CONSERVE OF THE SQUILL.

Take fresh squills w. one ounce,

Double refined sugar w. five ounces.

Beat them in a mortar into a conserve.

Keep all conserves, but particularly conserve of arum and the squill, in close vessels.

SUCCI.

S U C C I.
J U I C E S.

SUCCUS COCHLEARIÆ COMPOSITUS.

COMPOUND JUICE OF SCURVY-GRASS.

TAKE of juice of scurvy-grass two pints,
beccabunga,
water-cress, of each one pint,
Seville oranges M. twenty ounces.

Mix them; let the dregs subside, and pour off the liquor clear, or strain it.

SUCCUS BACCÆ SAMBUCI SPISSATUS.

INSPISSATED JUICE OF THE ELDER-BERRY.

Take of the juice of elder-berries, pressed out and clarified, two pints.

Evaporate

Evaporate it in a bath of water saturated with common salt, till it is of a proper consistence.*

Inspissate the juice of black currants, lemons, † and hemlock gathered when the flower is just making its appearance, in the same manner.

* I did not know the reason of the College using this kind of bath, and therefore tried the effects of it, and a common water bath. I found that I could not make watery fluids boil in a common water bath; but that they boiled in this, and the evaporation went on of course much faster; and on trying with a thermometer, I found the heat to be increased to 230° in the bath, and the substances were not rendered at all empeurymatic. I found that greater heat may be obtained by dissolving other salts in the water; but how far the heat may be increased without danger of burning, practice only can determine.

† Lemon juice should be evaporated until not more than one sixth remains.

EXTRACTA

EXTRACTA ET RESINÆ.

EXTRACTS AND RESINS.

EXTRACT of CAMOMILE,
BROOM TOPS,
GENTIAN,
LIQUORICE,
BLACK HELLEBORE,
RUE,
SAVIN.

BOIL them in distilled water, press out the liquor, strain it and set it by till the dregs subside; then evaporate them in a bath of water saturated with common salt, until they be of a consistence proper for pills.

The same kind of baths should be used in the preparation of all extracts, that the evaporation may be rightly performed.

E EXTRACTUM

EXTRACTUM COLOCYNTHIDIS
COMPOSITUM.COMPOUND EXTRACT OF COLOCYN-
THIDA.

Take of the pith of colocynthida, cut in
small pieces, w. six drachms,
of focotorine aloes, powdered, w.
one ounce and an half,
of scammony powdered, w. half an
ounce,
of lesser cardamom seeds powdered,
without the capsules, w. one
drachm,
of weak spirit of wine one pint.

Digest the colocynthida in the spirit of wine
in a gentle heat for four days; add the aloes
and the scammony to the tincture squeezed out
of the pith; distill off the spirit until the mass
is of a proper consistence for pills, adding the
powder of the cardamom seeds towards the
end.*

ELA-

* The spirit is here ordered to be distilled off, instead
of being evaporated. I suppose the College can only
mean

ELATERIUM.

Cut ripe wild cucumbers, squeeze the juice lightly out of them, pass it through a very fine hair sieve into a glass vessel, set it by for some hours until the thicker parts subside. Pour away the thinner part which floats at top, and put the rest into a filter,* filter away the thinner parts of it. Expose the thicker part which remains in the filter to a gentle heat, covering it with a linen cloth until it becomes perfectly dry.

E 2

EXTRAC.

mean by this the saving the spirit; in which case it will be sufficient to carry on the distillation until the spirit be all come over. The remaining part of the evaporation may be carried on in a proper bath in open vessels.

* The filter to be used on this occasion should be paper, such as is commonly called blotting paper. This paper, as we have it commonly in the shops, is stained; the stain frequently spoils the colour, and sometimes injures the substances filtered through it. Care therefore should be taken to procure it unstained.

EXTRACTUM LIGNI CAMPECHEN-
SIS.

EXTRACT OF LOGWOOD.

Take of logwood rasped one pound.*

Boil

* The College here and in several other cases have prescribed a particular quantity, when any quantity may be employed indiscriminately. I suppose they considered it as the quantity convenient for a private apothecary to make at one time. They have also in compound medicines given the proportions by ordering particular quantities, instead of proportional parts, for the same reason, I suppose.

Private apothecaries, especially in the country, who wish to serve their patients conscientiously, cannot be too anxious to make all preparations and compositions of medicines themselves. They may be judges of the simple medicines, and may buy only such as are perfect in their kind; but there are druggists who keep medicines at all prices, and take care often to make something similar to the compound ones as cheap as they can. It is true indeed, that great care is taken at Apothecary's Hall not only to have the best simple medicines, but also that they shall be properly and skilfully prepared and compounded, as I know, having once been admitted to see their warehouses and laboratory.

Those

which will be red and pellucid when hot, but yellow and turbid when cold; repeat the operation with fresh water until the liquor remains clear. When cold mix all these liquors together, and evaporate them to a proper consistence.

This extract should be prepared in two ways; one soft, fit for pills, one hard and capable of being powdered.*

EXTRACTUM CORTICIS PERUVI-
ANI CUM RESINA.

RESINOUS EXTRACT OF BARK.

Take of Peruvian bark, grossly powdered,
one pound,

Rectified spirit of wine four pints:

Digest

* A larger quantity of extract may be procured by these repeated boilings, but that will be from some of the substance of the bark which has no medical virtue being dissolved. The dose of course must be encreased, to have the same efficacy. I should therefore presume, that boiling it twice is as much as should be allowed.

Digest them for four days, and pour off the tincture. Boil the bark which remains in ten pints of distilled water in an open vessel, till it be evaporated to two pints. Strain the tincture and decoction separately; evaporate the water from the decoction, and distill the spirit from the tincture, till they begin to be thick; mix them together, and evaporate till they are of a consistence to form pills.

In the same manner should be prepared

Extract of Cascarilla,

Jalap.

EXTRACTUM SENNÆ.

EXTRACT OF SENNA.

Take of senna one pound,

Distilled water one gallon.

Boil the senna in the distilled water, after they are boiled add a little * rectified spirit of wine; strain the liquor, and evaporate it to a proper consistence.

E 4

OPIUM

* A little is an indefinite term, and certainly ought not to be employed in the College's prescription of medicines. Four ounces M. may be a proper quantity.

OPIUM PURIFICATUM.

PURIFIED OPIUM.

Take of opium, cut in small pieces, one pound,

Weak spirit of wine twelve pints.

Digest them in a gentle heat, shaking them now and then, till the opium be dissolved; strain the solution through paper. Distill the tincture so prepared to a proper consistence.*

Let purified opium be kept in a double form; one soft for making pills, the other hard that it may be powdered.

N O T E.

In making extracts, the liquor should be frequently stirred during the evaporation, especially after it begins to grow thick.

A little spirit of wine should be mixed with all softer watery extracts.†

OLEA

* It would be sufficient here to distill off the spirit and evaporate the remainder in an open vessel.

† To prevent them from growing mouldy.

OLEA EXPRESSA.

EXPRESSED OILS.

OLEUM AMYGDALÆ.

OIL OF THE ALMOND.

BEAT fresh, either sweet or bitter, almonds in a mortar, then press out the oil in a cold press.

In the same manner oil is to be expressed from the seeds of Flax bruised,

of Ricinus, the husk being first taken off,

and Mustard-seed bruised.*

OLEA

* These oils are, I believe, never prepared by the apothecary or druggist, but are always bought as merchandise, and might have been inserted therefore in the *Materia Medica*.

OLEA DISTILLATA.
ESSENTIAL OILS.

OLEUM ESSENTIALE ANISI.

ESSENTIAL OIL OF ANISEED,
CARRAWAY,
LAVENDER,
PEPPER-MINT,
COMMON MINT,
ORIGANUM,
PENNY-ROYAL,
ROSEMARY,
The JUNIPER-BERRY,
SASSAFRAS-ROOT.

THESSE oils are separated by distillation
in a common still with a worm tub, of
a proper magnitude. But to prevent burning,
a sufficient quantity of water is to be added to
the

the feeds. They should be digested with the water before the distillation.*

The water which distills over with the oil may be preserved for use.

OLEUM

* The maceration should be continued for different lengths of time, as the plants, or parts of plants, are of a softer or firmer texture. When they are of a soft texture, they may be put in over night and distilled the next morning. When of a firmer texture, they may be macerated four and twenty or six and thirty hours. It is better too to dry them a little before they are put into the water, as the water being imbibed again will crack the cells containing the oil, and let it out more easily. Where the essential oil is of sufficient value to afford it, it is better also to add salt in the distillation, to increase the heat of the boiling fluid, as essential oils are not so volatile as water.

A great deal of the oil remains suspended in the water, and this mixture is called a distilled water; where this is not valuable itself, it is better to use it in the next distillation, as by that means so large a proportion of the oil will not be waisted in it.

In some plants no essential oil at all is procured in the first distillation. But none of these are enumerated by the College.

The fluid should be made to run from the end of the worm into a vessel which will let the oil fall to the bottom, if it be of greater specific gravity than the water,
and

OLEUM PETROLEI.

OIL OF PETROLEUM.

Distill petroleum in a sand-heat.

OLEUM TEREBINTHINÆ.

OIL OF TURPENTINE.

Take of common turpentine five pounds,
water four pints.

Distill the turpentine with the water in a
copper still.

RESINA

and rise to the top if it be of less, while the bulk of the water runs off. It should not be allowed to run in a stream down into this vessel, but to fall upon the side of a funnel made flat for the purpose, what they call an air funnel, which should be fitted to the top of the vessel. The vessel should be filled with distilled water of the same plant, if it can be had, otherwise with plain distilled water, before the liquor begins to run into it from the worm. It should hold about a quart. As soon as no more oil collects, either the distillation should be stopped or the receiving vessel changed.

RESINA FLAVA.

YELLOW RESIN.

Remains in the still after the oil of turpentine is distilled off.

OLEUM TEREBINTHINÆ RECTIFICATUM.

RECTIFIED OIL OF TURPENTINE.

Take of oil of turpentine one pound,

Water four pints.

Distill them.*

OLEUM ANIMALE.

ANIMAL OIL.

Take of oil of hartshorn one pound.

Distill it three times. †

OLEUM

* When the oil of turpentine is all come over, the distillation should be stopped, leaving water enough to prevent the still from burning.

† The best manner of distilling this oil is to take an iron pot, such as is used in a sand heat, and fit to it an earthen-

OLEUM SUCCINI RECTIFICATUM.

RECTIFIED OIL OF AMBER.

Take of oil of amber one pound.

Distill it three times.

OLEUM VINI.

OIL OF WINE.

Take alcohol,

vitriolic acid, of each one pint.

Mix them gradually, and distill them,
taking care the black scum, which is apt to
arise,

earthen-ware head like the upper part of a retort. The oil should be put in the iron pot, and mixed with quicklime, powdered by slacking it in water, and afterwards heated red hot, so as to form a stiff paste. The head should be luted to the pot, and a tin receiver, such as a common tea cannister, should be fitted to it, and kept moist with a wet cloth wrapped round it. The distillation should then be performed with a moderate heat. This distillation should be repeated three times; the oil which is produced at the third distillation should be put
into

arise, does not pass into the receiver. Separate the oily part of the distilled liquor from the volatile vitriolic acid. Add to the oily part a sufficient quantity of water of pure kali, to take off the sulphureous smell; then distill off the æther with a gentle heat. The oil of wine remains in the retort swimming upon the watery fluid, from which it is to be separated.*

into a glass retort with a quart or two of water, and distilled as long as the oil comes over.

* Oil of wine may be also obtained by continuing the distillation in the preparation of æther after all the æther is distilled over. A very small portion of oil of wine is thus obtained. By using a larger proportion of vitriolic acid, the alkohol is more perfectly decomposed, little æther is produced, and a much larger proportion of oil of wine.

At first sight it might appear more proper to have brought all the processes for the decomposition of alkohol into æther, and oil of wine, together; but on more mature observation, I see it would be totally against the plan of the College, who have regarded nothing but similarity of form. Æther, indeed, might have been regarded as an oil rather than as a spirit.

This process succeeds best in a very thin flint glass retort, containing three or four times the quantity of the liquor

64 THE LONDON DISPENSATORY.

liquor to be distilled, and supported, or suspended, over a charcoal fire, which may be removed or extinguished immediately if the scum should rise too high. But in this case the operation must be performed where there is no danger from fire.

S A L E S.

S A L E S.

S A L T S.

ACIDUM VITRIOLICUM DILUTUM.

DILUTE VITRIOLIC ACID.

TAKE of vitriolic acid, w. one ounce,
 distilled water, w. eight ounces.

Mix them gradually.*

ACIDUM NITROSUM.

NITROUS ACID.

Take of purified nitre, w. fixty ounces,
 vitriolic acid, w. twenty-nine
 ounces.

F Mix

* They are ordered to be mixed gradually, lest the heat produced should break the vessel, which must be glass, or earthen-ware. But if a thin glass vessel, such as a Florence flask, be used, they may be poured in and mixed at once.

Mix and distill them.

The specific gravity of this acid is to that of distilled water as 1,550 to 1,000.*

ACIDUM NITROSUM DILUTUM.

DILUTE NITROUS ACID.

Take nitrous acid,
distilled water, of each one pound.

Mix them together.

ACIDUM MURIATICUM.

MURIATIC ACID.

Take of sea-salt dried ten pounds,
vitriolic acid six pounds,
water five pounds.

First

* This distillation is best performed in a glass retort and tubulated receiver, the tube placed downwards, and entering into a large phial, the neck of which nearly fits it. The tube should reach within a quarter of an inch of the bottom of the phial, that so none of the vapour of the acid

First mix the acid with the water, pour them upon the salt, and distill them.*

The specific gravity of this acid is to that of distilled water as 1,170 to 1,000. .

F 2

ACETUM

acid may escape without passing through the liquor, after the distillation has gone on some little way. If the acid be required of a very high colour, the heat towards the end of the distillation should be very great.

Part of the acid by a second distillation may be obtained colourless. But that is of no consequence in pharmacy.

* The cheapest mode of obtaining this acid is by performing the distillation in the following manner.

An iron pot is to be set in brick-work, and fitted with a flat lid, which may be screwed down to it. The lid should be lined on the inside with common luting of clay and sand, which should be dried, excepting where it rests upon the edge of the pot; it should then be screwed down tight. In the lid there should be two holes, one an inch and an half, the other three quarters of an inch in diameter. Into the largest hole a tube should be inserted, and luted fast with a lute of dry clay formed into a thick paste with linseed oil; this tube should rise up about six inches, and then be bent with an angle inclining downwards, and enter into the side of a vessel similar to that commonly employed for separating oils from watery fluids, and called
a separating

ACETUM DISTILLATUM.

DISTILLED VINEGAR.

Take of vinegar five pints.

Distill

a separating funnel; this should be fixed so that there may be placed under it a vessel containing water, the tube of the funnel passing six or eight inches down into the water. Into the other hole in the lid a tube should be fitted, and luted; this tube should rise eight inches, be bent down six inches, and then be bent up a foot.

Before the lid is screwed down upon the pot, a quantity of salt is to be put into the iron pot previously dried and decrepidated. The lid is then to be screwed down, the apparatus fixed, and the stopper put into the funnel. Vitriolic acid is then poured into the crooked tube; as soon as the bent part is full, a portion will fall into the pot upon the salt, a quantity of vapour of the acid will come over, and combine with the water. When vapour ceases to come over, more vitriolic acid is to be poured in: and this is to be continued as long as fresh vitriolic acid occasions any bubbling through the water. Then the fire is to be lighted; more vapour will come over. When vapour ceases to come over in a moderate heat, a little more vitriolic acid is to be poured in; if this produces more vapour, more and more is to be added as long as fresh bubbles arise. When no more vapour

Distill it with a moderate heat, in glass

F 3

vessels,

vapour is produced by the addition of vitriolic acid, the fire is to be raised to a red heat, and continued as long as there is any appearance of vapour coming over.

If there should be the appearance of any fluid in the upright part of the larger tube which conveys the vapour, the vessel with the water impregnated with acid is immediately to be removed, and the distillation stopped.

If during the distillation the water impregnating with the acid should become very hot, the vessel is to be removed, and another similar one put in its place. When that is become hot, it is to be removed, and the first one replaced: and this should be repeated until the liquor has acquired the proper specific gravity.

It sometimes happens in this operation that the vapour suddenly ceases to arise. If this should happen, the liquor will rise up into the first receiver or separating funnel: to prevent it from rising into the pot, the stopper of the first receiver is to be taken out until fresh vapour begins to be extricated, when the stopper should be replaced.

To procure the glauber salt, a quantity of lime sufficient to saturate any superfluous acid should be thrown upon the mass remaining in the pot, and a quantity of water sufficient to dissolve the salt should be applied. The whole should be made to boil, then be taken out into a vessel where the impurities may subside. The clear liquor should be taken off, evaporated, and crystallized in flat vessels.

vessels, as long as it comes over free from a burnt taste.*

ACIDUM ACETOSUM.

ACETOUS ACID.

Take of verdigris grossly powdered two pounds.

Dry it very thoroughly in a salt-water bath; then distill it in a sand heat; and rectify the acid by a second distillation.

The specific gravity of this acid is to that of distilled water as 1,050 to 1,000:†

SAL

* The strongest vinegar should be employed for this purpose, as in that case what comes over at the first is fit for use. If weak vinegar be taken, not only the trouble and expence of the first part of the distillation is lost, but the product acquires a burnt taste as soon as that which comes over is worth preserving.

A copper body with a glass head is the best apparatus for this distillation.

† I found upon trying this process and that which was given in the specimen, that the produce by the process in the specimen contained both fixed and volatile vitriolic acids,

SAL ET OLEUM SUCCINI.

SALT AND OIL OF AMBER.

Take of amber two pounds.

Distill it in a sand heat, gradually increasing the fire; there will arise up an acid liquor, an oil, and a salt rendered impure by a mixture of the oil.

SAL SUCCINI PURIFICATUS.

PURIFIED SALT OF AMBER.

Take of salt of amber half a pound,
distilled water one pint.

F 4

Boil

acids, and was obtained at about double the expence. The acid produced by the process in the Dispensatory does not shew the least mark of copper, not acquiring the smallest purple blue colour from the application of any quantity of volatile alkali.

The way to judge of the verdigris being perfectly dry, is by covering the vessel in which it was put in the bath with any cool cover of metal or glass for about a minute. If any vapour condenses on the metal or glass, the verdigris is not sufficiently dry.

The distillation should be performed in a common glass retort and receiver.

Boil the salt in the distilled water,* and set by the liquor that the salt may crystallize.

FLORES BENZOËS.

FLOWERS OF BENZOËS.

Take of powdered benzoës one pound.

Put them into an earthen pot in a sand heat, and sublime the flowers with a moderate heat into a paper cone fitted to the pot.

If the flowers should be yellow, mix them with white clay, and sublime them again.†

KALI

* The salt is by this means rendered purer than by another process, which is done by imbibing the oil from the salt by means of filtering paper, and subliming it from white clay; in which case a small portion of the oil adheres to the salt, which some have thought rendered it a better medicine, adding the antispasmodic quality of the oil, which in this preparation is not disagreeable to the taste or stomach, to the sedative quality of the salt.

† It is better to sublime them out of a retort, the neck of which is cut off near to the body.

KALI PRÆPARATUM.

PREPARED KALI.

Take of pearl-ashes two pounds,
distilled water boiling hot three
pints.

Dissolve the kali in the distilled water, and strain the solution through paper; then evaporate the liquor till a film appears upon the surface. Set it by for a night, that the neutral salts may crystallize. Afterwards pour off the liquor and evaporate the whole * water, frequently stirring it towards the end, lest the salt should adhere to the pot.

In the same manner impure kali may be purified, when it is found in the ashes of any vegetable.

The

* Not only the water of solution, but also the water of crystallization; for the crystals of kali contain a large portion of water. The salt therefore may be had in dry powder, and may still contain this water. If therefore the whole water be not evaporated, the dose will be different in the same weight.

The same salt may be prepared from tartar burnt until it becomes of an ash colour.*

AQUA KALI.

WATER OF KALI.

Take of kali one pound.

Set it in a moist place till it becomes fluid, then strain it.†

AQUA

* Kali is the common Arabic name for fixed vegetable and fixed fossil alkali, which the Arabians could not distinguish from one another, *al* being only the article *the*. The plants from whence large quantities of alkali are produced got all the common name of *kali*, from the salt; as we should express it in English the *kali plant*. Just as we say the *camphor tree*, or as in older English, *glass-wort*.

Giving single names to the alkalis gives great facility to the new names of compounds.

I suppose the French chemists have read Dean Swift's etymology, in which he proves, that all languages were derived from the English, and therefore have called fixed vegetable alkali *pot-ashes*.

† This solution differs from a saturated solution of kali in water, in that if any part of it should be uncombined with gas, it will attract it from the air, so that the whole shall be saturated with gas.

AQUA KALI PURI.

WATER OF PURE KALI.

Take of kali four pounds,
lime six pounds,
distilled water four gallons.

Pour four pounds of the water upon the lime, let them stand for an hour, then put in the kali and the rest of the water. Boil them for a quarter of an hour, let them stand to cool, and strain the liquor. A pint of this liquor ought to weigh sixteen ounces. If the liquor should effervesce with any acid, add more lime.

KALI PURUM.

PURE KALI.

Take water of pure * kali one gallon.

Evaporate

* Alkalis, it is well known, are often combined with a vapour which has been called by some mephitic air, by others

I

Evaporate it to dryness, melt it and pour it out.*

CALX CUM KALI PURO.

LIME MIXED WITH PURE KALI.

Take of lime, w. five pounds four ounces,
water of pure kali sixteen pints.

Boil the water of kali till three fourths of it are evaporated; moisten the lime with water, and let it fall into a powder; then sift it

others fixed air, others aerial acid, and others gas, and these names are not any of them generally adopted. The College seem to have hesitated in adopting either of them, and therefore I suppose have chosen to avoid giving a compound name to the compound of kali and this vapour, and have called it *Kali præparatum*. Which however cannot possibly remain, if the new system of compound names continues. The name of pure added to what used to be called caustic alkalis is certainly necessary, as it is now universally allowed that in this state they are uncombined with any other substance.

* This is generally employed by the surgeons in London as a caustic, in preference to the next preparation.

it into the liquor. Keep the paste thus formed in a stopped vessel.

NATRON PRÆPARATUM.

PREPARED NATRON.

Take of barilla powdered two pounds,
distilled water one gallon.

Boil the barilla in four pints of the water for half an hour, and strain it. Boil the part which remains in the strainer with the rest of the water, and strain it. Evaporate together the liquor from both boilings to two pints, and set them by for eight days.* Strain this liquor again, and after a proper evaporation †, set

* This part of the process, which I thought superfluous, I find on trial to be necessary, in order to allow a very fine black powder to subside, and which would otherwise contaminate the salt.

† The College have not here specified what a proper evaporation is: nor is it possible, because different masses of barilla contain different proportions of natron. The quantity of evaporation is to be ascertained by trial; to wit, by evaporating a little, letting it stand to cool, and when no crystals are formed evaporating a little more, and so on.

set the liquor by to crystallize. Dissolve the crystals in distilled water, strain the liquor, evaporate it, and set it by that fresh crystals may be formed.*

AMMONIA PRÆPARATA.†

PREPARED AMMONIA.

Take of sal ammoniac powdered one pound,
prepared

* This second crystallization is not only useful to render the crystals finer, but also to get rid of phlogisticated alkali, which is often found in barilla.

Barilla contains a great variety of different substances. Sometimes a little neutral salt, which is easily got rid of by the crystals of the alkali forming themselves fair, and large, and separate. The quantity of the neutral salts in barilla is hardly ever considerable. But the operator should take care not to get Alicant kelp instead of barilla, as it contains a large quantity of sea-salt.

After the first crystallization, a fresh evaporation gives a fresh quantity of natron; but if we go on evaporating, instead of crystals a gelatinous mass is obtained, which can only be used for purposes where any impure alkali may be employed.

† This name is quite a new one for volatile alkali. According to the College's plan of using compound names
for

prepared chalk two pounds.

Mix them together and sublime them.*

AQUA AMMONIÆ PURÆ.

WATER OF PURE AMMONIA.

Take of sal ammoniac one pound,

lime

for chemical compounds, it was necessary to give simple names to the alkalis. Names can only come from authority or usage; the College have therefore taken the Arabic name for what used to be called fixed vegetable alkali, the Greek name for what used to be called fixed fossil alkali, but there was no single name whatever used by any nation for volatile alkali; it was therefore necessary to invent one. I have heard some people object to this name as not being a pretty one. But different persons have different tastes.

I shall here take an example of the difference in the compound names, if single names had not been given to the alkalis. What formerly was called vitriolated tartar, is called by the College kali vitriolatum; it would have been, to have made it perfect in the other way, alkali vegetabili fixum causticum vitriolatum.

* This sublimation is best performed in an iron pot, with an earthen-ware head entering with a large tube into a receiver made of iron tinned, which we commonly call tin.

lime two pounds,
water one gallon.

Add two pints of the water to the lime, let them stand together an hour; then add the sal ammoniac and six pints of the water boiling hot, cover the vessel immediately; let the liquor stand to cool, pour it off, and distill a pint with a moderate heat.*

AQUA

* The College have given directions in this, and several other cases, only for the preparation of the medicine, without any directions what was to be done with, or what advantage was to be made of, the substances remaining after its formation. It is not to be supposed, however, that these are to be thrown away, if any use can be made of them. In the present case a great deal of volatile alkali would be lost.

If large quantities of aqua ammoniæ puræ should be wanted, the best way is to pour the clear liquor into a common still, wash the earth that remains with a fresh quantity of water, and throw it likewise into the still; repeat the washing two or three times. The calcareous earth is then to be thrown away. The liquor in the still is to be distilled through a common worm into a receiver, tying a bladder round the neck of the receiver and round the end of the worm, and pricking a hole in it with a pin. Two quarts are then to be distilled for every pound of sal ammoniac; the liquor distilled over is to be put into a
common

AQUA AMMONIÆ.
WATER OF AMMONIA.

Take of sal ammoniac one pound,
pearl-ash one pound and an half,
water four pints.

Distill two pints with a gentle heat.*

G LIQUOR

common retort, and distilled with a moderate heat till the quantity ordered by the College comes over; then the receiver is to be changed, and the distillation continued until what comes over does not smell of volatile alkali. The volatile alkali procured by this second distillation may be either used where a weaker solution is required, or it may be concentrated by another distillation.

Some portion of the volatile alkali will be lost if a common receiver be used; which may be avoided by employing one which has a tube coming out from its side, and after running a little way horizontally, is bent down into water contained in a phial. But the saving will hardly pay the expence of the apparatus, and the breakage of the glasses.

There will remain in the retort a compound of muriatic acid and calcareous earth, of which no use is made at present.

* In this preparation the ammonia is not perfectly saturated with gas, there being always some pure kali in pearl-
ash

LIQUOR VOLATILIS, SAL, ET OLEUM CORNU CERVI.

THE VOLATILE LIQUOR, SALT, AND OIL OF HARTSHORN.

Take of hartshorn ten pounds.

Distill it with a fire gradually encreased. A volatile liquor, a salt, and an oil will come over. Separate the salt and the oil, and distill the liquor three times.

Mix the salt with an equal weight of prepared chalk, sublimate it, and repeat the operation three times, or until the salt becomes white.

The

ash which will therefore detach part of the ammonia pura, it suggested itself to me that it would be better and cheaper to saturate water with prepared ammonia. On trying the experiment, there was very little difference in the expense; the smell of that produced by the College process was more pungent, and could not be entirely taken off by calx muriata. I do not therefore understand why this preparation was preferred, especially as the residuum is of no use, kali muriatum, formerly called digestive salt of filvius, not being now used in medicine.

The same volatile liquor, salt, and oil, may be procured from any part, excepting the fat, of any animal.*

G 2

KALI

* This distillation is best performed in an iron cylindrical vessel laid horizontally in a reverberatory furnace, the two ends of the cylinder coming out at the two sides of the furnace. One end should be open, with an iron lid to fit it pretty exactly; the other should be shut, excepting a hole near the top capable of receiving a tube two inches in diameter in the inside. This tube should pass through a vessel filled with water into a large tin receiver, which ought likewise to be immersed in water. From this receiver a tin tube should rise, consisting of joints fitted into one another closely, and should be altogether from forty to an hundred feet in length; the end of the tube may be brought into the ash-hole of the furnace. The cylinder being filled with bones, the lid is to be luted on, the fire raised, and the distillation continued as long as any vapour comes out of the end of the tube. The lid is then to be taken off, the bones raked out, if their ashes are wanted white, into a common grate, otherwise into water. The cylinder is immediately to be filled with fresh bones, the lid put on, and the distillation carried on in the same manner. This is to be repeated until it be judged that the receiver is sufficiently full; then the operation is to be stopped, and when all is cool, the salt, liquor, and oil, is to be taken out of the receiver, the tube

KALI VITRIOLATUM.

VITRIOLATED KALI.

Take of the salt which remains after the
distillation of nitrous acid two
pounds,
distilled water two gallons.

Evaporate the superfluous acid from the salt
over a strong open fire; then boil it for a
little while in the water; strain it, and set it
by to crystallize.

NATRON VITRIOLATUM.

VITRIOLATED NATRON.

Take of the salt which remains after the
distillation of muriatic acid two
pounds,
distilled

is to be taken to pieces, and the salt adhering to the sides
of it is to be beaten out.

The purification may be performed in glass vessels.

This salt contains a small quantity of empyreumatic oil,
which has been conceived to add to its antispasmodic vir-
tues in hysterical complaints.

distilled water two pints and an half.

Evaporate the superfluous acid from the salt with a strong open fire; then boil it in the water for a short time, strain it, and set it by to crystallize.*

NITRUM PURIFICATUM.

PURIFIED NITRE.

Take of nitre two pounds,
water four pints.

G 3 Boil

* Glauber salt is not only procured in this way, but also in making sal ammoniac, the process for which follows.

The ammonia procured from the distillation of bones is saturated with the vitriolic acid which is left after the crystallization in the manufactories of copperas. The vitriolated ammonia is boiled with sea salt in water, and set by for the glauber salt to crystallize. After the glauber salt is crystallized the water is evaporated, and the sal ammoniac sublimed. If the glauber salt should be tinged with iron, it is to be boiled with a little lime, and crystallized over again.

Boil the nitre in the water until it is dissolved, strain the liquor, and set it by to crystallize.

KALI ACETATUM.

ACETATED KALI.

Take of kali one pound.

Boil it with a moderate fire in four or five times its weight of distilled vinegar. When the effervescence is over, add more distilled vinegar at different times, until the greater part of the first vinegar being evaporated, the fresh makes no new effervescence; which happens generally when about twenty pints of vinegar have been used: then dry it slowly. There is left an impure salt, which is to be melted with a moderate fire, then dissolve it in water, and strain it through paper. If the melting be properly performed, the strained liquor will be colourless, otherwise of a brown colour. Lastly, evaporate this liquor out of a shallow glass vessel, stirring the salt as it concretes, that it may dry the sooner. It should

should be kept in a stopped vessel. The salt ought to be very white, and perfectly soluble both in water and alcohol without leaving any dregs. If the salt, although it be white, should let fall any sediment when dissolved in alcohol, its solution in alcohol is to be strained and dried again.*

AQUA AMMONIÆ ACETATÆ.

WATER OF ACETATED AMMONIA.

Take of ammonia two ounces,
distilled vinegar four pints, or as
much as is necessary to saturate
the ammonia, †

Mix them together.

KALI TARTARISATUM.

TARTARISED KALI.

Take of kali one pound,
G 4 crystals

* This salt some time ago was esteemed a powerful diuretic. It has by no means maintained its reputation.

† This salt being of nearly equal volatility with water is with difficulty procured in its solid form.

crystals of tartar three pounds,
 distilled water boiling hot one gal-
 lon.

Dissolve the kali in the water, and throw in gradually the crystals of tartar powdered. Let the liquor cool, and strain it through paper; evaporate part of the water, and set it by to crystallize.

NATRON TARTARISATUM.

TARTARISED NATRON.

Take of natron, w. twenty ounces,
 crystals of tartar powdered two
 pounds,
 distilled water boiling hot ten pints.

Dissolve the natron in the water, and add gradually the crystals of tartar; strain the liquor through paper, evaporate and set it by to crystallize.*

ALU-

* Both these salts, especially tartarised natron, are much fairer, if the saturation is made in the cold either in glass or earthen-ware vessels. But it requires a long digestion,

ALUMINIS PURIFICATIO.

PURIFIED ALUM.

Take of alum one pound,
chalk, w. one drachm,
distilled water one pint.

Boil them a little, strain them, and set them by to crySTALLIZE.*

ALUMEN USTUM.

BURNT ALUM.

Take of alum half a pound.

Burn it in a close vessel as long as it continues to bubble. †

If

digestion, and that they should be stirred every now and then during the digestion.

* There is a small quantity of superfluous vitriolic acid adheres to alum, which renders it unfit for lotions in several cases; this acid is removed by the chalk.

† Many chemical authors use the words *ustulatio*, and *ustro*, to express any decomposition made by heat in open vessels,

If the crystals of salts have any impurities adhering to them, they are first to be washed with the liquor left after the crystallization, and afterwards with a little distilled water, or rectified spirit of wine.

When any salt is crystallized out of any liquor, pour the liquor off, and strain it if necessary. Evaporate part of the liquor, and set it by until fresh crystals are formed; and repeat this as long as pure crystals can be procured.*

vessels, when part of the substance is thrown off in vapour. This sense of the word however by no means accords with our common idea of burning in English. In this operation the water of crystallization is evaporated from the alum.

* To obtain fine crystals, a large quantity of the materials should be set to crystallize at once, the evaporation should be only carried on so far as to allow a small quantity to crystallize at once, and the liquor should be suffered to cool very slowly.

MAGNESIA.

M A G N E S I A.

MAGNESIA ALBA.

WHITE MAGNESIA.

TAKE bitter salt,
kali, of each two pounds,
distilled water boiling hot twenty
pints.

Dissolve the bitter salt in ten pints of water, and the kali in the other ten pints; strain them through paper separately, and mix them together. Boil the liquor for a short time, and strain it through linen while it is yet hot; the magnesia will remain on the linen strainer; free it from the vitriolated kali by frequently washing it with distilled water.

MAGNESIA

MAGNESIA USTA.
BURNT MAGNESIA.

Take of white magnesia, w. four ounces.

Expose it to a strong fire for two hours.
After it cools, keep it in a stopped glass
vessel.*

* By this operation the gas is thrown off from the mag-
nesia, with a view of avoiding flatulency when it is ex-
hibited. But this is rather a refinement in speculation
than in practice, the gastric juices taking up the gas im-
mediately.

If pure kali be employed in the precipitation of mag-
nesia, the same intention will be fulfilled.

PRÆ-

PRÆPARATA E SULPHURE.

PREPARATIONS OF SULPHUR.

FLORES SULPHURES LOTI.

WASHED FLOWERS OF SULPHUR.

TAKE of flowers of sulphur one pound,
distilled water four pints.

Boil the flowers of sulphur for a little in the water, pour off this water, and wash away the acid with cold water; then dry the flowers.

KALI SULPHURATUM.

SULPHURATED KALI.

Take of flowers of sulphur, w. one ounce,
kali, w. five ounces.

Melt

Melt the sulphur with a moderate fire, mix in the salt, and continue to stir them briskly until they unite.*

OLEUM SULPHURATUM,
ET
PETROLEUM SULPHURATUM.
SULPHURATED OIL,
AND
SULPHURATED PETROLEUM.

Take of flowers of sulphur, w. four ounces,
olive oil, w. sixteen ounces.

Boil them together in a pot slightly covered until they unite.

In the same manner make sulphurated petroleum.

SULPHUR PRÆCIPITATUM.
PRECIPITATED SULPHUR.

Take of sulphurated kali six ounces,
distilled

* When the kali has begun to unite with the sulphur, the heat must be gradually encreased so as to keep the whole fluid.

PREPARATIONS OF SULPHUR. 95

distilled water one pound and an
half,

dilute vitriolic acid a sufficient
quantity.

Boil the sulphurated kali in the distilled
water until it is dissolved. Strain the liquor
through paper, and pour in the vitriolic acid.
Wash the precipitated powder with water re-
peatedly, until it becomes insipid.

PRÆPARATA

PRÆPARATA EX ANTI-
MONIO.

PREPARATIONS OF ANTI-
MONY.

ANTIMONIUM CALCINATUM.
CALCINED ANTIMONY.

TAKE of antimony powdered, w. eight
ounces,
nitre powdered two pounds.

Mix them, and throw them by a little at
a time into a red hot crucible; burn the white
substance left after the deflagration about half
an hour, when it is cold powder it, then wash
it with distilled water.*

CROCUS

* This calx of antimony rarely produces any effect,
and has been employed principally by practitioners who
were afraid of using an active preparation of this metal,
and yet chose to give antimony because it was the fashion.

CROCUS ANTIMONII.

CROCUS OF ANTIMONY.

Take antimony powdered,
nitre powdered, of each one pound,
common salt, w. one ounce.

Mix them, and throw them by a little at a time into a red hot crucible, then encrease the heat until the whole is melted; pour them out, and after they are cold separate the scoria.*

ANTIMONIUM MURIATUM.

MURIATED ANTIMONY.

Take crocus of antimony in powder,
vitriolic acid, of each one pound,
common salt dried two pounds.

H

Put

* They should be poured out into an iron inverted cone.

The effect of this process is to destroy part of the sulphur of the antimony, and leave the regulus combined with a smaller proportion of sulphur.

Put the vitriolic acid into a retort, mix the salt and the crocus together, put them into the retort by a little at a time; distill them in a sand heat. Let the distilled matter stand in an open vessel for some days, and pour off the liquid part from the dregs.

PULVIS ANTIMONIALIS.

ANTIMONIAL POWDER.

Take antimony grossly powdered,
hartshorn rasped, of each two
pounds.

Mix them, and throw them into a broad iron pot red hot; stir them constantly until they become of an ash colour. Powder the mass when it is cold, and put it into a coated crucible, lute into it another crucible inverted; there must be a small hole in the bottom of the inverted crucible. Put it into the fire, and heat it gradually until it becomes red hot; keep it in this degree of heat for two hours.

When

PREPARATIONS OF ANTIMONY. 99

When it has stood to be cold, take the mass out of the crucible, and powder it finely.*

SULPHUR ANTIMONII PRÆCIPITATUM.

PRECIPITATED SULPHUR OF ANTIMONY.

Take of antimony in powder two pounds,
water of pure kali four pints,
distilled water three pints.

Mix them together, and boil them with a moderate fire for three hours, stirring them constantly, and adding the distilled water as it is found necessary. Strain the liquor while hot through a double linen cloth, and pour into it by a little at a time, while it is yet warm, a sufficient quantity of dilute vitriolic

H 2 acid

* This is supposed to be the preparation known under the name of Dr. James's Powder.

acid to precipitate the sulphur. Wash off the vitriolated kali with warm water.*

ANTIMONIUM TARTARISATUM.

TARTARISED ANTIMONY.

Take of crocus of antimony in powder one pound and an half,
crystals of tartar two pounds,
distilled water two gallons.

Boil them in a glass vessel about a quarter of an hour, strain the liquor through paper, and set it by to crystallize. †

ANTI-

* In this operation the regulus of antimony is precipitated in fine powder mixed with the sulphur, but not combined with it, so as to leave it exposed to be acted upon by the salts of the stomach.

Care is to be taken to stir the whole very perfectly together when the precipitation is finished, there being different proportions of the regulus of antimony and sulphur in the first and latter precipitations.

† Lately a calx of antimony formed by the decomposition of muriated antimony has been employed to form
this

ANTIMONIUM VITRIFICATUM.

VITRIFIED ANTIMONY.

Take of powdered antimony, w. four ounces.

Burn it in an earthen vessel, encreasing the heat slowly, and stirring it with an iron rod, until it throws off no more sulphureous vapours. Put this powder in a crucible capable of containing a third part more. Adapt a cover to it, and put it into a fire at first moderate, but afterwards encreased until the matter melts, then pour it out.*

H 3

PRÆPA-

this combination; some likewise have used glass of antimony. Not seeing any reason for preferring the crôcus, I have made some enquiry into this subject, and have been informed that these three modes of preparation were made by the Committee at Apothecary's Hall, and tried at one of the Royal hospitals, and that this was found to be the preferable one: but the particulars I have not been able to learn.

This is the preparation of antimony which can be most depended upon, its only fault is the smallness of the quantity required for a dose when exhibited as an alterative.

* It should be poured out upon a slab.

PRÆPARATUM EX ARGENTO.

PREPARATION OF SILVER.

ARGENTUM NITRATUM.

NITRATED SILVER.

TAKE of silver, w. one ounce,
dilute nitrous acid, M. four
ounces.

Dissolve the silver in the acid in a glass vessel upon warm sand; then increase the heat gently, and evaporate it to dryness. Melt the mass in a crucible, taking care that the heat be not increased to a greater degree than is necessary, and pour it out into proper moulds.*

PRÆPA-

* Pure silver should be employed for this operation. If the silver the operator has be not pure, the best way is to precipitate it from the solution by means of copper, to take it out as soon as it is precipitated, wash it first with water, then with a little water of ammonia, and it will then be fit for the operation.

PRÆPARATA E FERRO.
PREPARATIONS OF IRON.

FERRUM AMMONIACALE.

AMMONIACAL IRON.

TAKE of iron filings one pound,
sal ammoniac two pounds.

Mix them, and sublime them. Rub the sublimate and the part which has not risen together, until they be thoroughly mixed, and sublime them a second time.*

H 4

FERRI

* In this operation part of the sal ammoniac is decomposed, the ammonia coming over pure, which may be condensed and preserved if the sublimation is performed in a retort to which is adapted a receiver containing water. Part of the iron of course is united with the muriatic acid. This compound, however, does not rise, the sublimate consists of iron combined with a part of the sal ammoniac which is not decomposed.

The

FERRI RUBIGO.

RUST OF IRON.

Take of iron filings one pound.

Expose them to the air, and keep them moist with water until they rust; then rub them in an iron mortar, pour on distilled water, and when the grosser parts have subsided, pour off the water with the fine powder suspended in it into another vessel. The remaining part, which cannot be rubbed to fine powder easily, is to be again exposed to the air and moisture to rust: it is then to be rubbed and washed in the same manner. When the fine powder has subsided, pour off the water and dry it.

FERRUM

The College have omitted the solution of this salt in spirits, which was called tincture of flowers of steel, and thought by many practitioners one of the best preparations of iron, as it is more capable of being diffused and suspended in a large proportion of water than any other of them, and may therefore be given with similar effects to these of chalybeate mineral waters.

FERRUM TARTARISATUM.

TARTARISED IRON.

Take of iron filings one pound,
powdered crystals of tartar two
pounds.

Mix them with distilled water into a thick
mass, expose them to the air in a broad
earthen vessel for eight days: then dry the
mass in a sand heat, and powder it very finely.

FERRUM VITRIOLATUM.

VITRIOLATED IRON.

Take iron filings,
vitriolic acid, of each, w. eight
ounces,
distilled water three pints.

Mix them in a glass vessel, and when they
have ceased to emit bubbles, let them stand
for some time upon warm sand; then pour
off the liquor, and strain it through paper,
make

make a proper evaporation, and set it by to crystallize.*

* The iron filings should be very pure and carefully separated from copper or brass, or any other metal, by the magnet; and the purity should be ascertained by digesting a few of them in a little water of ammonia. Otherwise this preparation will have no advantage over common green vitriol crystallized.

It is better not to saturate the vitriolic acid quite perfectly, as a little superfluous acid prevents the salt from decomposing so readily as it is otherwise apt to do.

PRÆPARATA

PRÆPARATA EX HY-
DRARGYRO.

PREPARATIONS OF QUICK-
SILVER.

HYDRARGYRUS PURIFICATUS.

PURIFIED QUICKSILVER.

Take quicksilver,

iron filings, of each four pounds.

Rub them together, and distill them out of
an iron vessel.*

HYDRAR-

* Quicksilver may commonly be had in the shops very pure, so that this operation is seldom necessary, only when after it has been squeezed through leather its surface soon loses its lustre.

To render quicksilver extremely pure, muriated mercury should be rubbed with an equal quantity of kali, and a little charcoal powdered, and then distilled in glass vessels.

HYDRARGYRUS ACETATUS.

ACETATED QUICKSILVER.

Take of purified quicksilver one pound,
dilute nitrous acid two pounds,
water of kali a sufficient quantity.

Dissolve the quicksilver in the acid in a glass vessel with a sand heat, then pour in the water of kali by a little at a time, until the calx of the quicksilver is precipitated. Wash the calx with a great deal of distilled water, and dry it with a gentle heat.

Take of the calx of quicksilver thus made
one pound,
acetous acid a sufficient quantity to
dissolve the calx.

Mix them in a glass vessel, and when the quicksilver is dissolved, strain the liquor through paper, then evaporate till a film covers the surface of it, and set it by to crystallize. Keep the crystals in a stopped vessel.

HYDRAR-

HYDRARGYRUS CALCINATUS.

CALCINED QUICKSILVER.

Take of purified quicksilver one pound.

Expose the quicksilver to a heat of about 600° in a glass bolt-head with a flattish bottom, until it concretes into a red powder.

HYDRARGYRUS CUM CRETA.

QUICKSILVER WITH CHALK.

Take of purified quicksilver, w. three
ounces,

powdered chalk, w. five ounces.

Rub them together until the globules disappear.*

HYDRAR-

* Quicksilver may be rubbed in this manner into very fine particles, but it is not at all altered by or combined with the chalk, as is evident from our being able to dissolve away the chalk by muriatic acid, and leave the quicksilver behind apparently in a very fine powder; but if this
apparent

HYDRARGYRUS MURIATUS.

MURIATED QUICKSILVER.

Take purified quicksilver,
vitriolic acid, of each two pounds,
common salt thoroughly dry three
pounds and an half.

Mix the acid with the quicksilver in a glass vessel, and boil them in a sand heat till the mass is thoroughly dry. Mix it when it is cold with the common salt in a glass vessel, then sublime it in a glass cucurbit with a fire gradually increased. Lastly, separate the sublimed matter from the scoriæ.*

CALOMELAS.

apparent powder be rubbed in a glass mortar, the small globules will again unite.

Quicksilver divided by rubbing it with a dry powder of any kind very rarely has any effect.

* This sublimation requires a deep pot for the sand heat, and to be so set that the fire may play round it to a considerable height, otherwise the sublimate is with great difficulty separated from the scoriæ.

A glass

PREPARATIONS OF QUICKSILVER. III

CALOMELAS.

CALOMEL.

Take of muriated quicksilver one pound,
purified quicksilver, w. nine ounces.

Rub them together till the globules disappear, and sublime them; rub the whole mass and sublime it again; in the same manner repeat the operation so that there shall be four sublimations in all. Afterwards powder the sublimate very finely, and wash it with boiling distilled water.

HYDRARGYRUS MURIATUS MITIS.

MILD MURIATED QUICKSILVER.

Take purified quicksilver,
dilute nitrous acid, of each half a
pound.

Mix

A glass head should be put on the top of the cucurbit to catch the vapour which rises higher than the sublimate, by this means a quantity of quicksilver will be saved.

The old operation appears a very inelegant one, but is still thought by some operators the cheapest.

Mix them in a glass vessel, and set them by till the quicksilver is dissolved. Heat them till the salt is dissolved. Pour the hot liquor into a glass vessel, into which there has first been put another hot liquor, consisting of

Common salt, w. four ounces,
Distilled water eight pints.

When a white powder has fallen to the bottom of the vessel, pour off the clear liquor, and wash the powder with hot water frequently until it is insipid. Lay the powder upon filtering paper, and dry it with a gentle heat.*

HYDRAR-

* The College observe in the Index, that this is a preparation which, although it was contained in former editions of the Dispensatory, was rejected in the last. Mr. Scheele supposed he had invented a new and easier method of forming and powdering calomel. This preparation and the former, however, are by no means the same, this being apter to act on the stomach and intestines. It would seem most probable that all the saline preparations of quicksilver are converted in the blood-vessels into muriated quicksilver, by the sea salt and sal ammoniac which is found in the blood. Vitriolated, nitrated, and acetated quicksilver, are converted partly into muriated
and

HYDRARGYRUS NITRATUS RUBER.
RED NITRATED QUICKSILVER.

Take purified quicksilver,
nitrous acid, of each one pound,
muriatic acid, w. one drachm.

Mix them in a glass vessel, and dissolve the quicksilver in a sand heat; then increase the heat until the matter is converted into red crystals.*

I

CALX

and partly into mild muriated quicksilver, as soon as they are touched by either of these salts; and calomel and mild muriated quicksilver are both of them totally convertible into muriated quicksilver by sal ammoniac. The difference therefore of the action of these preparations one might expect to be entirely in the stomach and intestines. However, this is but conjecture; and indeed no consequences should be drawn from chemical philosophy into the method of preparation, or method of action of medicines, unless confirmed by experience in diseases themselves; for the very slightest alteration in the manner of preparing a chemical substance will very frequently alter some property on which its medicinal effects may depend.

* The small quantity of muriatic acid renders the crystals larger and fairer.

Some

CALX HYDRARGYRI ALBA.
WHITE CALX OF QUICKSILVER.

Take muriated quicksilver,
fal ammoniac,
water of kali, of each half a
pound.

Diffolve first the fal ammoniac, and afterwards the muriated quicksilver in the same distilled water, then add the water of kali. Wash the precipitated powder until it be tasteless.

HYDRARGYRUS CUM SULPHURE.
QUICKSILVER WITH SULPHUR.

Take purified quicksilver,
flowers

Some portion of the nitrous acid is driven off by the heat, and a little yellowish matter sublimes into the upper part of the vessel, containing a considerable quantity of quicksilver.

Quicksilver may be obtained from the waste part in all these operations by distilling it with iron filings, or either of the fixed alkalis and a little charcoal powdered.

PREPARATIONS OF QUICKSILVER. 115

flowers of sulphur, of each one
pound.

Rub them together till the globules disappear.*

HYDRARGYRUS SULPHURATUS
RUBER.

RED SULPHURATED QUICKSILVER.

Take of purified quicksilver, w. forty
ounces,

fulphur, w. eight ounces.

Melt the sulphur and mix in the quick-
I 2 silver.

* This preparation was formerly called Æthiop's mineral. This name has also been given to a substance formed by melting sulphur and pouring quicksilver into it, stirring them till the quicksilver be combined with the sulphur. When they are only rubbed together, the quicksilver is in very fine globules, which are mixed with the sulphur in a fine powder, but each of them retains its properties. The sulphur acts as a mild laxative, the quicksilver has rarely any effect; it has however sometimes. When they are combined by melting together, the sulphur loses its laxative quality if the proportions be perfect, and the quicksilver is defended by it so as never to produce any effect.

silver. If the mixture should catch fire, extinguish it by covering the vessel. Powder the mass thus formed, and sublime it.*

HYDRARGYRUS VITRIOLATUS.
VITRIOLATED QUICKSILVER.

Take purified quicksilver,
vitriolic acid, of each one pound.

Mix them in a glass vessel, and heat them gradually till they unite in a white mass, which is to be entirely dried by a strong fire. This mass is to be washed with a large quantity of warm distilled water, which renders it of a yellow colour, and breaks it down into a fine powder. Rub this powder carefully with the water in a glass mortar. After the powder has subsided, pour off the water, and add fresh distilled water; do this repeatedly until the powder has no longer any taste.†

PRÆPARATA

* This compound is now generally considered as having no effect, excepting in fumigation.

† There is a considerable quantity of quicksilver in the washings which may be saved.

PRÆPARATA E PLUMBO.

PREPARATIONS OF LEAD.

CERUSSA ACETATA.

ACETATED CERUSSE.

TAKE of cerusse one pound,
 distilled vinegar one gallon and
 an half.

Boil the cerusse with the vinegar until the acid be saturated, then strain it through paper, and after a proper evaporation set it by to crytallize.

AQUA LITHARGYRI ACETATI.

WATER OF ACETATED LITHARGE.

Take of litharge, w. two pounds four ounces,
 distilled vinegar one gallon.

I 3

Mix

Mix them, and boil them down to six pints, constantly stirring them, and set them by. After the dregs have subsided strain the liquor.*

* This salt and the former would seem at first fight the same; but in making them I found them differ very much in their chemical properties. Whether they differ in their medicinal I am not informed.

The water of acetated litharge has been introduced into practice some time ago under the name of Goulard's wash for external applications. The uncertainty of its dose arising from the different strengths of the distilled vinegar should prevent it from being used internally; as, if lead is ever to be employed internally, we should be anxious to determine the dose accurately.

PRÆPARATUM

PRÆPARATUM E STANNO.

PREPARATION OF TIN.

STANNUM PULVERATUM.

POWDERED TIN.

TAKE of tin six pounds.

Melt it in an iron vessel, and stir it with an iron rod until a powder is formed and swims on the top. Take off this powder as it forms, let it stand to cool, and pass it through a sieve.

PRÆPARATA E ZINCO.
PREPARATIONS OF ZINC.

ZINCUM CALCINATUM.
CALCINED ZINC.

TAKE of zinc broken in pieces, w.
eight ounces.

Heat a large deep crucible placed leaning to one side white hot; throw in the zinc by a piece at a time, and place over the crucible another, so however that the air may be suffered to come to the hot zinc. Take the calx as it forms out of the apparatus, and pass the white lighter part through a sieve.*

ZINCUM

* This calx has been lately introduced for internal use being purer than cadmia or tutty, which are also procured by calcination by heat.

ZINCUM VITRIOLATUM.

VITRIOLATED ZINC.

Take of white vitriol one pound,
 vitriolic acid, w. one drachm,
 distilled water boiling hot three
 pints.

Mix them, and strain them through paper,
 and after a proper evaporation set them by in
 a cold place to crystallize.*

* White vitriol not uncommonly contains a little of
 some other metallic salts, as well as other impurities,
 which are separated by this process. I suppose the vitriolic
 acid is added, to prevent the salt from calcining on being
 dissolved in water, as it has been observed in the Philoso-
 phical Transactions by a member of the College, that
 metallic salts are apt to decompose if they be dissolved in
 a large proportion of water.

AQUÆ

AQUÆ DISTILLATÆ.
DISTILLED WATERS.*

AQUA DISTILLATA.
DISTILLED WATER.

TAKE of spring water ten gallons.

Distill over four pints, which are to be thrown away; then distill four gallons, which are

* Excepting the first, these distilled waters are mixtures of a small proportion of essential oil with water; the oil not being combined with, but simply suspended in water. Although they be procured in distilling the essential oils as before prescribed, the College seem to have ordered them to be prepared when the essential oil is not required by itself, because a sufficient quantity of the distilled water would not have been procured for use.

For this purpose a much larger quantity of water in proportion to the plant should be employed in the distillation, than where the essential oil is wanted separately.

are to be preserved in a glafs or earthen-ware bottle stopped with a glafs stopper.

AQUA ANETHI.

DILL WATER.

Take of dill seed bruised one pound,
water a sufficient quantity to prevent burning.

Distill over one gallon.

AQUA CINNAMOMI.

CINNAMON WATER.

Take of bark of cinnamon bruised one pound,
a sufficient quantity of water to prevent burning.

Steep the cinnamon in the water for four and twenty hours, distill them together, drawing over one gallon.

AQUA

AQUA FÆNICULI.

FENNEL WATER.

Take of the seeds of sweet fennel bruised
one pound,
a sufficient quantity of water to
prevent burning.

Distill over one gallon.

AQUA MENTHÆ PIPERITIDIS.

PEPPER-MINT WATER.

Take of pepper-mint thoroughly dried one
pound and an half,
a sufficient quantity of water to
prevent burning.

Distill over one gallon.

AQUA MENTHÆ SATIVÆ.

COMMON MINT WATER.

Take of common mint thoroughly dried
one pound and an half,
a sufficient

DISTILLED WATERS. 125

a sufficient quantity of water to
prevent burning.

Distill over one gallon.

AQUA PIMENTO.

JAMAICA PEPPER WATER.

Take of Jamaica pepper half a pound,
a sufficient quantity of water to
prevent burning.

Steep the Jamaica pepper in the water for
twenty-four hours, then distill them together,
drawing over one gallon.

AQUA PULEGII.

PENNYROYAL WATER.

Take of Pennyroyal thoroughly dried one
pound and an half,
a sufficient quantity of water to
prevent burning.

Distill over one gallon.

AQUA

AQUA ROSÆ.

ROSE WATER.

Take of the fresh flower leaves of the damascene roses, the yellow points being taken off, six pounds,
 a sufficient quantity of water to prevent burning.

Distill over one gallon.

We have ordered dry herbs for the distillation of their waters, because they cannot be had all the year fresh. When fresh ones are used, the proportional weight should be increased. Whether, however, they are used fresh or dry, we leave it in the option of the operator to vary the proportional weight, according to the climate and weather in which the herbs grew, or were gathered.

Plants and seeds which have been kept more than a year, are less proper for the distillation of their waters.

To each gallon of these waters add, m. five ounces of weaker spirit of wine.

SPIRITUS

SPIRITUS DISTILLATI.
DISTILLED SPIRITS.

ALKOHOL.

TAKE of rectified spirit of wine one gallon,
kali hot one pound and an half,
pure kali, w. one ounce.

Mix the spirit of wine with the pure kali, and add of the hot kali one pound; shake them together, and digest them for twenty-four hours. Pour off the spirit, add to it the rest of the kali, and distill it in a water-bath. Keep it in a vessel well stopped.

The specific gravity of alkohol is to that of distilled water as 815 to 1,000.*

SPIRITUS

* I suppose the pure kali is employed to take out the essential oil from the spirit. The London Rectifiers employ it to take off the flavour of malt spirits.

If

SPIRITUS ÆTHERIS VITRIOLICI.

SPIRIT OF VITRIOLIC ÆTHER.

Take rectified spirit of wine,
vitriolic acid, of each one pound.

Pour the acid by a little at a time into the spirit, and shake them till they are mixed; then distill the spirit of vitriolic æther with a moderate fire, from a retort into a tubulated receiver, the tube of which enters into another receiver, to which it is fitted, until sulphureous vapours begin to arise.

ÆTHER

If it be wished to have pure alkohol, the end of the worm should be fitted into a receiver, by tying a bladder which has been rendered soft by moistening it, but dried as well as possible with a cloth, round the end of the worm in the neck of the receiver. Small holes should be made with a pin through the bladder. Alkohol attracts water so strongly from the air, that if it runs in a stream through it into the receiver, it will be considerably diluted.

The kali should be heated nearly red hot, especially that which is put in the still; and in my opinion it would be better to distill the spirit after it is poured off from the first kali, and then digest it with the second portion of kali, afterwards to pour it off and distill it by itself.

ÆTHER VITRIOLICUS.

VITRIOLIC ÆTHER.

Take of spirit of vitriolic æther two pounds,
 water of pure kali, M. one ounce,
 Skake them together, and distill over with
 a gentle heat, M. fourteen ounces.*

K SPIRITUS

* The process for forming æther from alcohol and vitriolic acid has been variously given and reasoned upon by different authors. What seems to me the best idea is, that alcohol consists of phlogiston, acetous acid, and water. When vitriolic acid is applied to alcohol, it unites with part of the phlogiston, and sets loose part of the water, leaving the acetous acid combined with a less proportion of phlogiston and water forming æther. If the exact proportion of vitriolic acid is not employed, if there be too little, or if there be water combined with the alcohol, or vitriolic acid, the alcohol is not all decomposed; the proportion of æther is therefore less, and the remaining alcohol is combined with the æther, and renders part or the whole of it soluble in water. By the re-distillation as ordered in this process, the æther is separated from the part of the alcohol which comes over along with it, but this does not avoid the loss of the alcohol which remains un-decomposed.

SPIRITUS ÆTHERIS NITROSI.

SPIRIT OF NITROUS ÆTHER.

Take of rectified spirit of wine two pints,
nitrous acid half a pound.

Mix

un-decomposed. In order therefore to procure the æther in its full quantity, the pure alkohol, instead of rectified spirit, should be employed. Considering, however, that the College might have some reason for preferring a process in which there is an evident loss, I bought a small quantity at Apothecary's Hall, and compared it with some I made with the same proportions of pure alkohol and acid, and found them differ only in that mine smelt a little more of volatile vitriolic acid, but it agreed in all other properties.

If the distillation be continued, oil of wine is procured in very small quantity, not nearly sufficient for the present consumption, Hoffman's anodyne liquor having been more used lately, being recommended by the physician of the greatest business in this country.

If vitriolic acid be applied in a larger proportion, it decomposes part of the æther, uniting with the phlogiston, and also detaching a part of the water, leaving the acetous acid combined with less phlogiston and water, so forming oil of wine and very little æther; as in the process given by the College for oil of wine.

In

Mix them by pouring the acid into the spirit, and distill over with a gentle heat, w. one pound ten ounces.*

SPIRITUS AMMONIÆ.

SPIRIT OF AMMONIA.

Take weak spirit of wine three pints,
sal ammoniac, w. four ounces,
pearl-ash, w. six ounces.

K 2

Mix

In the process prescribed by the College for making æther, if what comes over first in the first distillation of vitriolic æther be taken away by changing the receiver, more æther will be procured in the second distillation.

The vitriolic acid which remains in the retort will decompose about half the first proportion of alkohol.

* This operation should be performed in a tubulated retort, and tubulated receiver, the tube going down into another receiver. The spirit should be put into the retort, the acid poured in at once, and the tube of the retort immediately be shut. If the acid be put into the retort first, an ebullition often takes place, and occasions the destruction of the whole apparatus.

Nitrous æther differs from vitriolic in the smell principally. It has however been considered as a very different medicine.

Mix them, and distill over with a moderate fire one pint and an half.

SPIRITUS AMMONIÆ FOETIDUS.

FOETID SPIRIT OF AMMONIA.

Take weak spirit of wine six pints,
fal ammoniac one pound,
asa foetida, w. four ounces,
pearl-ash one pound and an half.

Mix them, and distill over with a moderate heat five pints.*

SPIRITUS ANISI COMPOSITUS.

COMPOUND SPIRIT OF ANISE.

Take aniseed bruised,
angelica seed bruised, of each half
a pound,
weak spirit of wine one gallon,
a sufficient

* There being a little caustic alkali in the pearl-ash, these two preparations will be somewhat more pungent than if prepared kali had been used.

DISTILLED SPIRITS. 133

a sufficient quantity of water to
prevent burning.

Distill over one gallon.*

SPIRITUS CARUI.

SPIRIT OF CARAWAY.

Take caraway feeds bruised half a pound,
weak spirit of wine one gallon,
water a sufficient quantity to pre-
vent burning.

Distill over one gallon.

SPIRITUS CINNAMOMI.

SPIRIT OF CINNAMON.

- Take bark of cinnamon bruised one pound,
weak spirit of wine one gallon,

K 3 a sufficient

* In this and the following prescriptions, the essential oils of the plants are dissolved in the spirit. They were formerly called, improperly, compound or spirituous waters.

134 THE LONDON DISPENSATORY.

a sufficient quantity of water to
prevent burning.

Distill over one gallon.

SPIRITUS JUNIPERI COMPOSITUS.
COMPOUND SPIRIT OF JUNIPER.

Take of juniper berries bruised one pound,
caraway seeds bruised,
fennel seeds, of each, w. one ounce
and an half,
weak spirit of wine one gallon,
a sufficient quantity of water to
prevent burning.

Distill over one gallon.

SPIRITUS LAVENDULÆ.
SPIRIT OF LAVENDER.

Take of fresh lavender flowers one pound
and an half,

z

weak

DISTILLED SPIRITS. 135

weak spirit of wine one gallon,
Distill over in a water bath five pints.

SPIRITUS MENTHÆ PIPERITIDIS.

SPIRIT OF PEPPERMINT.

Take the dried herb of peppermint one
pound and an half,
weak spirit of wine one gallon,
a sufficient quantity of water to
prevent burning.
Distill over one gallon.

SPIRITUS MENTHÆ SATIVÆ.

SPIRIT OF GARDEN MINT.

Take the dried herb of garden mint one
pound and an half,
weak spirit of wine one gallon,
a sufficient quantity of water to
prevent burning.
Distill over one gallon.

K 4

SPIRITUS

SPIRITUS NUCIS MOSCHATÆ.

SPIRIT OF NUTMEG.

Take of nutmegs bruised, w. two ounces,
weak spirit of wine one gallon,
a sufficient quantity of water to
prevent burning.

Distill over one gallon.

SPIRITUS PIMENTO.

SPIRIT OF PIMENTO.

Take the bruised berries of pimento, w.
two ounces,
weak spirit of wine one gallon,
a sufficient quantity of water to
prevent burning.

Distill over one gallon.

SPIRITUS PULEGII.

SPIRIT OF PENNYROYAL.

Take of the dry herb of Pennyroyal one
pound and an half,
weak spirit of wine one gallon,
a sufficient

a sufficient quantity of water to
prevent burning.

Distill over one gallon.

SPIRITUS RAPHANI COMPOSITUS.
COMPOUND SPIRIT OF HORSE-RADISH.

Take the fresh root of horse-radish,
the exterior rind of Seville oranges
dried, of each two pounds,
of the fresh herb of garden scurvy-
grafs four pounds,
bruised nutmegs, w. one ounce,
weak spirit of wine two gallons,
a sufficient quantity of water to
prevent burning.

Distill over two gallons.

SPIRITUS RORISMARINI.

SPIRIT OF ROSEMARY.

Take of the fresh tops of rosemary one
pound and an half,
weak spirit of wine one gallon.

Distill over five pints in a water bath.

DECOCTA

DECOCTA ET INFUSA.
DECOCTIONS AND INFU-
SIONS.

DECOCTUM CORNU CERVI,
DECOCTION OF HARTSHORN.

TAKE of burnt hartshorn prepared, w,
two ounces,
gum arabic, w. six drachms,
distilled water three pints.

Boil them to two pints, constantly stirring
them, and strain the liquor.*

DECOCTUM

* In this preparation there is very little, if any, of the calcareous earth or phosphorated lime dissolved in the water, and therefore it should only be strained through a coarse strainer. There seems also no advantage obtained by boiling it so long.

DECOCTUM CORTICIS PERUVIANI.
DECOCTION OF PERUVIAN BARK.

Take of Peruvian bark powdered, w. one
ounce,
distilled water, M. one pint and
three ounces.

Boil them ten minutes in a covered vessel,
and strain the liquor while it is hot.*

DECOCTUM PRO ENEMATE.
DECOCTION FOR AN ENEMA.

Take of the dry leaves of mallow, w. one
ounce,
dry chamomile flowers, w. half
an ounce,
water, one pint.

Boil them, and strain the liquor.

DECOCTUM

* In this decoction of bark I suppose the College mean
to retain the flavour of the essential oil. It seems calcu-
lated to be employed only as a strengthening remedy.

DECOCTUM PRO FOMENTO.

DECOCTION FOR FOMENTATION.

Take dry leaves of southernwood,
dry tops of sea wormwood,
dry chamomile flowers, of each,
w. one ounce,
dry laurel leaves, w. half an ounce,
distilled water six pints.

Boil them a little, and strain the liquor.*

DECOCTUM HELLEBORI.

DECOCTION OF HELLEBORE.

Take of hellebore root powdered, w. one
ounce,
distilled water two pints,
rectified spirit of wine, w. two
ounces.

Boil

* There does not seem to be much reason for an admixture of these herbs in fomentation; the astringency in the laurel leaves, particularly, seems averse to the emollient effects of a fomentation.

Boil the water with the root to one pint, let the liquor cool, strain it, and add the spirit.

DECOCTUM HORDEI.

BARLEY WATER.

Take of pearl barley, w. two ounces,
distilled water four pints.

Wash the barley clean with cold water, then boil it a little in about a pint of water, pour off this water, add the distilled water boiling hot, boil it down to two pints, and strain the liquor.*

DECOCTUM HORDEI COMPOSITUM.

COMPOUND DECOCTION OF BARLEY.

Take of decoction of barley two pints,
raisins, the stones being taken out,
figs

* Boiling the barley in the first water coagulates it, and renders the solution afterwards less rosy and more agreeable.

figs sliced, of each, w. two ounces,
 liquorice root cut and bruised, w.
 half an ounce,
 distilled water one pint.

Boil them to two pints, and strain the li-
 quor.

DECOCTUM SARSAPARILLÆ.

DECOCTION OF SARSAPARILLA.

Take of sarsaparilla sliced, w. six ounces,
 distilled water one gallon.

Macerate the root in the water for two
 hours in a heat of about 195° ; take it out of
 the liquor, bruise it, and when bruised, re-
 turn it into the same liquor, and macerate it
 again for two hours; then boil the liquor to
 four pints, press it out, and strain it.*

DECOC-

* Sarsaparilla is one of those medicines which hardly
 yields its efficacious parts to any menstruum, and has
 therefore been exhibited in powder; particularly lately in
 hectic cases. But the great bulk of its powder renders it
 disagreeable;

DECOCTUM SARSAPARILLÆ COM-
POSITUM.

COMPOUND DECOCTION OF SARSAPA-
RILLA.

Take of sarsaparilla sliced and bruised, w. six
ounces,
bark of the root of saffras,
the wood of guaiacum rasped,
liquorice root bruised, of each, w:
one ounce,
bark of the root of mezereon, w.
three drachms,
distilled water ten pints.

Macerate all but the mezereon in a gentle
heat for six hours, then boil them down to
five pints, adding the bark of the mezereon
root towards the end. Strain the liquor.*

DECOC-

disagreeable; it is on this account, I suppose, that the
College have adopted this mode of trying to dissolve it,
which is nearly the same as the prescription found in the
Dispensatory of St. Thomas's hospital.

* I cannot understand the use of this complicated pre-
scription, but suppose the College must have left it as one
of

DECOCTUM ULMI.

DECOCTION OF ELM.

Take of the interior bark of fresh elm
bruised, w. four ounces,
distilled water four pints.

Boil them down to two pints, and strain
the liquor.

MUCILAGO AMYLI.

MUCILAGE OF STARCH.

Take of starch, w. three drachms,
distilled water one pint.

Rub the starch, pouring in the water by a
little at a time, then boil them a little.

MUCILAGO

of those which are to be thrown out in another edition.
The College indeed are rather to be praised for throwing
out so many absurd mixtures, than blamed for leaving a
few.

MUCILAGO ARABICI GUMMI.

MUCILAGE OF GUM ARABIC.

Take of gum arabic in powder, w. four
ounces,

distilled water boiling hot, m. eight
ounces.

Rub the gum with the water until it is
dissolved.

MUCILAGO SEMINIS CYDONII
MALI.

MUCILAGE OF QUINCE SEED.

Take of quince seed, w. one drachm,
distilled water, m. eight ounces.

Boil them with a moderate fire till the
water becomes thick; then strain the liquor
through a linen cloth.

L INFUSUM

INFUSUM GENTIANÆ COMPOSITUM.

COMPOUND INFUSION OF GENTIAN.

Take of gentian root, w. one drachm,
the exterior rind of fresh lemons,
w. half an ounce,
the exterior rind of Seville oranges
dried, w. one ounce and an
half,
water boiling hot, M. twelve
ounces.

Macerate them for an hour, and strain the
liquor.*

INFUSUM

* It is a question that does not seem to have been sufficiently agitated, whether medicines having nearly the same virtues are better exhibited singly, or mixed together. There was among the ancients a very apparent ground on which such mixtures were often made. They wanted a medicine of a peculiar degree of heat or cold, moisture or dryness, which if it did not reside in any one of the ingredients, might nevertheless be acquired by
mixing

distilled water boiling hot one pint.

Macerate the fenna and the ginger in the water for an hour in a covered vessel; let them cool, and strain the liquor.

INFUSUM SENNÆ TARTARISATUM.

TARTARISED INFUSION OF SENNA.

Take of fenna, w. one ounce and an half,
coriander seeds bruised, w. half an
ounce,

crystals of tartar, w. two drachms,
distilled water one pint.

Boil the crystals of tartar in the water until they are dissolved; pour the boiling hot liquor upon the fenna and the seeds; let them stand for an hour in a covered vessel; strain the liquor when cold.

AQUA CALCIS.

LIME WATER.

Take of lime half a pound,

distilled

distilled water boiling hot one gallon and an half.

Mix them, and set them by in a covered vessel for an hour; pour off the liquor, and preserve it in a stopped vessel.*

INFUSUM ROSÆ.

INFUSION OF THE ROSE.

Take the flower leaves of the red rose before they are full blown, the yellow points being removed, w. half an ounce,

dilute vitriolic acid, w. three drachms,

distilled water boiling hot two pints and an half,

double refined sugar, w. one ounce and an half.

L 3

First

* Care must be taken to let the air come at the liquor as little as possible, as the gas which is in the air would unite with the lime, and precipitate it from the water, if it was exposed.

First pour the water upon the flower leaves in a glass vessel, then add the dilute vitriolic acid, and macerate them for half an hour. Let the liquor cool, strain it, and add the sugar.

ACETUM SCILLÆ.

VINEGAR OF THE SQUILL.

Take of squills fresh dried one pound,
vinegar six pints,
weak spirits of wine half a pint.

Macerate the squills in the vinegar, in a gentle heat, in a glass vessel for four and twenty hours. Squeeze out the liquor, set it by till the dregs subside. Lastly pour off the liquor, and add the spirit.*

VINA

* Vinegar has been supposed by some to correct the disposition in squills to produce sickness, without hurting its other virtues; while others have supposed that it only did this by diminishing its virtues altogether, and I believe on better ground.

This preparation does not keep well, tincture of squills is therefore preferable to it.

VINA MEDICATA.
MEDICATED WINES.

VINUM ALOES.

ALOETIC WINE.

TAKE of socotorine aloës, w. eight
 ounces,
 canella alba, w. two ounces,
 mountain wine six pints,
 weak spirits of wine two pints.

Powder the aloës and the canella alba separately, then mix them together, and pour on the wine and the spirits. Digest them together for fourteen days,* shaking them now and then; and afterwards strain them.

L 4

It

* I am informed that the time of digestion of these wines was taken, but I do not know whether principally or wholly, from experiments made by the Committee of the Company of Apothecaries appointed to meet the Committee of the College.

It is not amifs to mix clean white fand with the powder, it prevents the aloës from sticking together when they are moistened.

VINUM ANTIMONII.

ANTIMONIAL WINE.

Take glafs of antimony in powder, w. one ounce,
mountain wine one pint and an half.

Digest them for twelve days, frequently stirring them, and strain them through paper.

VINUM ANTIMONII TARTARISATI.
WINE OF TARTARISED ANTIMONY.

Take of tartarised antimony, w. two scruples,
distilled water boiling hot, M. two ounces,
mountain wine, M. eight ounces.

Diffolve

Dissolve the tartarified antimony in the boiling distilled water, and add the wine.*

VINUM

* Tartarified antimony, formerly called emetic tartar, requires often to be exhibited in so small a dose as a quarter of a grain even to adults. Solutions of it have therefore been made in order for its division, scales used in apothecaries shops being seldom in such good order as to weigh very small weights with accuracy, independent of the want of attention which sometimes happens in their journeymen and apprentices.

Antimonial wine affords a solution of this or some other salt of antimony, which may be divided so as to give a very small dose. But the strength of that medicine, depending upon the quantity of tartar or some other salt contained in the wine, is very unequal.

I was formerly acquainted with the late Dr. Huck, who afterwards took the name of Saunders, of whom I enquired the reason of the tartarified antimony not being dissolved altogether in water, or in wine. He informed me, that if it was altogether dissolved in water it would not keep. And I have somewhere read in the Philosophical Transactions, that all metallic salts dissolved in a very large proportion of water were decomposed if they had any surface exposed to the air; and that the solution in wine took place with great difficulty. Therefore the tartarified antimony was first dissolved in the water, and the wine added to prevent its decomposition. He informed me
likewise

VINUM FERRI.

WINE OF IRON.

Take filings of iron, w. four ounces,
mountain wine four pints.

Digest them for a month, shaking them
frequently, and strain the liquor.

VINUM IPECACUANHÆ.

IPECACUAN WINE.

Take of ipecacuan root bruised, w. two
ounces,
mountain wine two pints.

Digest them for ten days, and strain the
liquor.

VINUM

likewise that it was a criterion by which the goodness of
the tartarised antimony might be judged of; as, if the tar-
tar was not saturated with the antimony, it would be pre-
cipitated by the wine in white powder.

This preparation has stood for some time in the Dis-
pensatory of St. Thomas's Hospital.

VINUM RHABARBARI.

RHUBARB WINE.

Take of rhubarb sliced, w. two ounces and
an half,

leffer cardamom seeds, taken out
of the husk and bruised, w. half
an ounce,

saffron, w. two drachms,

mountain wine two pints,

weak spirit of wine, M. eight
ounces.

Digest them for ten days, and strain the
liquor.*

* I apprehend that the spirit is added here, and in the
former case, to prevent the wine from fermenting. The
saffron can only give colour.

TINCTURÆ.

T I N C T U R Æ.
T I N C T U R E S.

TINCTURA ALOES.

TINCTURE OF ALOES.

TAKE of socotorine aloës powdered, w.
half an ounce,
extract of liquorice, w. an ounce
and an half,
distilled water,
weak spirit of wine, of each half
a pint.

Digest them in a sand heat, shaking the
vessel now and then until the extract be dis-
solved, and strain the liquor.

TINCTURA

TINCTURA ALOES COMPOSITA.
COMPOUND TINCTURE OF ALOES.

Take of tincture of myrrh two pints,
saffron,
socotorine aloës, of each, w. three
ounces,

Digest them for eight days, and strain the
liquor.

TINCTURA ASÆ FOETIDÆ.
TINCTURE OF ASA FŒTIDA.

Take of asa fœtida, w. four ounces,
rectified spirit of wine two pints.

Digest them for six days in a gentle heat,
and strain the liquor.

TINCTURA BALSAMI PERUVIANI.
TINCTURE OF BALSAM OF PERU.

Take balsam of Peru, w. four ounces,
I rectified

rectified spirit of wine one pint.

Digest them till the balsam be dissolved.

TINCTURA BALSAMI TOLUTANI.

TINCTURE OF BALSAM OF TOLU.

Take balsam of Tolu, w. one ounce and
an half,

rectified spirit of wine one pint.

Digest them till the balsam be dissolved,
and strain the liquor.*

TINCTURA BENZOËS COMPOSITA.

COMPOUND TINCTURE OF BENZOES.

Take of benzoës, w. three ounces,

strained styrax, w. two ounces,

balsam of Tolu, w. one ounce,

focotorine

* If either of these three tinctures be poured into water,
the gums or balsams will be separated, and mix with the
water in very small particles, forming a milky liquor;
which may be exhibited immediately, and by this means
equally diffused in the stomach.

focotorine aloës, w. half an ounce,
rectified spirit of wine two pints.

Digest them in a gentle heat for three days,
and strain the liquor.*

TINCTURA CANTHARIDIS.

TINCTURE OF CANTHARIDIS.

Take of cantharidis bruised, w. two-drachms,
cochineal powdered. w. half a
drachm,
weak spirit of wine one pint and
an half.

Digest them for eight days, and strain the
liquor.†

TINCTURA CARDAMOMI.

TINCTURE OF CARDAMOMS.

Take of lesser cardamom seeds bruised, after
having

* This seems to be one of those compound medicines
which the College have retained, whose compositions do
not appear to be founded on any reasonable ground.

† The cochineal can be of no use but to give colour,
which is not wanted.

having taken away the husks,
w. three ounces,
weak spirit of wine two pints.

Digest them for eight days, and strain the liquor.

TINCTURA CARDAMOMI COMPO-
SITA.

COMPOUND TINCTURE OF CARDA-
MOMS.

Take of lesser cardamom seeds, the husks
being taken away,
caraway seeds,
cochineal, of each of them pow-
dered, w. two drachms,
the bark of cinnamon bruised, w.
half an ounce,
raisins, the stones taken out, w.
four ounces,
weak spirit of wine two pints.

Digest

Digest them for fourteen days, and strain
the liquor.*

TINCTURA CASCARILLÆ.

TINCTURE OF CASCARILLA.

Take of the bark of cascarilla powdered,
w. four ounces,
weak spirit of wine two pints.

Digest them in a gentle heat for eight days,
and strain the liquor.

TINCTURA CASTOREI.

TINCTURE OF CASTOR.

Take of Ruffian castor, in powder, w. two
ounces,
M weak

* Mixtures of spices are evidently more agreeable to
the stomach than a larger dose of one taken singly; as is
well known not only in medicine, but even in cookery.
Sugar likewise renders spices and spirits more grateful to
the stomach. We find such mixtures not only in this,
but in several other prescriptions which come afterwards.

weak spirit of wine two pints.

Digest them for ten days, and strain the liquor.

TINCTURA CATECHU.

TINCTURE OF CATECHU.

Take of catechu, w. three ounces,

bark of cinnamon bruised, w. two ounces,

weak spirit of wine two pints.

Digest them for three days, and strain the liquor.

TINCTURA CINNAMOMI.

TINCTURE OF CINNAMON.

Take of cinnamon bark bruised, w. one ounce and an half,

weak spirit of wine one pint.

Digest them for ten days, and strain the liquor.

TINCTURA

TINCTURA CINNAMOMI COMPO-
SITA.COMPOUND TINCTURE OF CINNA-
MON.

Take of cinnamon bark bruised, w. six
drachms,

lesser cardamom seeds without the
husks, w. three drachms,

long pepper,

ginger, of each powdered, w. two
drachms,

weak spirit of wine two pints.

Digest them for eight days, and strain the
liquor.

TINCTURA COLOMBÆ.

TINCTURE OF COLOMBA.

Take of colomba root powdered, w. two
ounces and an half,

M 2 weak

weak spirit of wine two pints.

Digest them for eight days, and strain the liquor.

TINCTURA CORTICIS AURANTII.

TINCTURE OF THE RIND OF
ORANGES.

Take of the fresh rind of Seville oranges,
w. three ounces,
weak spirit of wine two pints.

Digest them for three days, and strain the liquor.

TINCTURA CORTICIS PERUVIANI.

TINCTURE OF PERUVIAN BARK.

Take of Peruvian bark in powder, w. four
ounces,
weak spirit of wine two pints.

Digest them with a gentle heat for eight
days, and strain the liquor.

TINCTURA

TINCTURA CORTICIS PERUVIANI
COMPOSITA.COMPOUND TINCTURE OF PERUVIAN
BARK.

Take of Peruvian bark powdered, w. two
ounces,

dried rind of Seville oranges, w.
one ounce and an half,

Virginian snake-root bruised, w.
three drachms,

saffron, w. one drachm,

cochineal powdered, w. two scru-
ples,

weak spirit of wine, M. twenty
ounces.

Digest them for fourteen days, and strain
the liquor.*

M 3 TINCTURA

* This seems to me an injudicious mixture. There
is certainly no stimulating power in cochineal; and the
saffron, at least in the dose in which it can be exhibited
here, seems to me to be incapable of producing any effect.

Some

TINCTURA FERRI MURIATI.
MURIATED TINCTURE OF IRON.

Take of iron rust half a pound,
muriatic acid, three pounds,
rectified spirit of wine three pints.

Pour the muriatic acid on the iron rust in a glass vessel; let them remain together for three days, shaking them now and then. Set them by till the dregs subside, and pour off the liquor. Evaporate it to a pint, and when cold, add the spirit of wine.*

TINCTURA

Some have thought Virginian snake-root particularly adapted to Peruvian bark as a stimulant. But I could never find that this opinion was grounded on any accurate observation.

* Iron in its metallic form will readily dissolve in muriatic acid. I conceived therefore that the College might as well employ iron filings. But as chemical theories, as has been before observed, are not to be admitted into medicine without trial, I made the two preparations, and found very considerable difference in them. Why the College made this change from their former edition, or which is the best medicine, I am not informed.

TINCTURA GALBANI.

TINCTURE OF GALBANUM.

Take of galbanum cut in small pieces, w.
two ounces,
weak spirit of wine two pints.

Digest them with a gentle heat for eight
days, and strain the liquor.

TINCTURA GENTIANÆ COMPOSI-
TA.

COMPOUND TINCTURE OF GENTIAN.

Take of gentian root, sliced and bruised,
w. two ounces,
the exterior rind of Seville oranges,
w. one ounce,
lesser cardamom seeds bruised, the
husks being taken away, w. half
an ounce,
weak spirit of wine two pints.

M 4

Digest

Digest them for eight days, and strain the liquor.

TINCTURA GUAIACI.

TINCTURE OF GUAIAACUM.

Take of gum guaiacum, w. four ounces,
compound spirit of ammonia one
pint and an half.

Digest them for three days, and strain the liquor.

TINCTURA HELLEBORI NIGRI.

TINCTURE OF BLACK HELLEBORE.

Take of the root of black hellebore in gross
powder, w. four ounces,
cochineal powdered, w. two scrup-
ples,
weak spirit of wine two pints.

Digest them in a gentle heat for eight days,
and strain the liquor.

TINCTURA

TINCTURA JALAPII.

TINCTURE OF JALAP.

Take root of jalap powdered, w. eight
ounces,
weak spirit of wine two pints.

Digest them in a gentle heat for eight days,
and strain the liquor.

TINCTURA LAVENDULÆ COMPO-
SITA.COMPOUND TINCTURE OF LAVEN-
DER.

Take spirit of lavender three pints,
spirit of rosemary one pint,
bark of cinnamon bruised,
nutmegs bruised, of each, w. half
an ounce,
red saunders, w. one ounce.

Digest them for ten days, and strain the
liquor.

TINCTURA

TINCTURA MYRRHÆ.

TINCTURE OF MYRRH.

Take of myrrh bruised, w. three ounces,
weak spirit of wine one pint and
an half,
rectified spirit of wine half a pint.

Digest them in a gentle heat for eight days,
and strain the liquor.

TINCTURA OPII.

TINCTURE OF OPIUM.

Take of hard purified opium powdered, w.
ten drachms,
weak spirit of wine one pint.

Digest them for ten days, and strain the
liquor.*

TINCTURA

* In the remarks which I have taken the liberty to
make on this edition of the Dispensatory, I have avoided
any comparison of it with former editions, or with other
Dispensatories.

TINCTURA OPII CAMPHORATA.
CAMPHORATED TINCTURE OF OPIUM.

Take hard purified opium,
flowers

Dispensatories. But it may be necessary, however, to enquire into the cause of the alteration of a medicine so very much in use as what has commonly been called liquid laudanum, *i. e.* a solution of opium which has been made in wine, in spirits, and in mixtures of wine and spirits. In the last edition of the London Dispensatory it was made in wine, and two ounces of opium was ordered to a pint. In this edition there are only ten drachms to a pint. Yet on trial, I find that the strength and narcotic power is nearly equal. If any thing the narcotic power of the tincture of opium is rather greater. The wine not being capable of dissolving the quantity of opium ordered in the tinctura thebaica, the appearance of the tincture of opium is thicker, more glutinous, and apter to dry upon the mouth of the phial, which renders it difficult to pour it out by drops, which I suppose is Sydenham's reason for employing wine rather than any other liquor. The solution in wine has two inconveniencies; one is that it is apt to ferment, and by that means the opium is partly destroyed, and the dose altered; wine also dissolves a larger proportion of opium in heat than in cold, and alters its strength in winter and summer.

The spices that were added to the thebaic tincture were not in sufficient quantity to have any effect.

flowers of benzoës, of each, w.
one drachm,

camphor, w. two scruples,
essential oil of aniseed, w. one
drachm,

weak spirit of wine two pints.

Digest them for three days.*

TINCTURA RHABARBARI.

TINCTURE OF RHUBARB.

Take of rhubarb sliced, w. two ounces,
lesser cardamom seeds bruised, and
freed

* It has been supposed that camphor joined to opium added to its power. But if this was the case, the quantity is here a great deal too small to have any effect. The same may be said of the flowers of benzoës. This mixture therefore, if it was good in principle, of which I do not know any ground, would have its purposes defeated by the proportions.

This preparation, as well as some of the others containing opium which follow, is in some degree calculated to be divided by measure or weight, half an ounce containing nearly a grain of opium,

TINCTURES.

173

freed from the husks, w. half
an ounce,

saffron, w. two drachms,
weak spirit of wine two pints.

Digest them for eight days, and strain the
liquor.

TINCTURA RHABARBARI COMPO-
SITA.

COMPOUND TINCTURE OF RHUBARB.

Take of rhubarb sliced, w. two ounces,
ginger powdered,
saffron, of each, w. two drachms,
liquorice root bruised, w. half an
ounce,
distilled water one pint,
weak spirit of wine, M. twelve
ounces.

Digest them for fourteen days, and strain
the liquor.

TINCTURA

TINCTURA SABINÆ COMPOSITA.

COMPOUND TINCTURE OF SAVIN.

Take of extract of favin, w. one ounce,
tincture of castor one pint,
tincture of myrrh half a pint.

Digest them together till the extract of favin is dissolved, and strain the liquor.*

TINCTURA SCILLÆ.

TINCTURE OF SQUILLS.

Take of squills fresh dried, w. four ounces,
weak spirit of wine two pints.

Digest

* Anti-spasmodics are generally disagreeable to the stomach. It is much better, therefore, to mix them together, that so each may be taken in a small quantity, and greater effect may be produced on the whole, upon a principle that has already been explained.

Digest them for eight days, and pour off
the liquor.*

TINCTURA SENNÆ.

TINCTURE OF SENNA.

Take of fenna one pound,
caraway feeds bruised, w. one ounce
and an half,
lesser cardamom feeds bruised, and
freed from the husks, w. half
an ounce,
raisins, the stones taken out, w.
sixteen ounces,
weak spirit of wine one gallon.

Digest them for fourteen days, and strain
the liquor.

TINCTURA

* This solution of squills keeps better than that which
is made in vinegar. I do not know whether the College
have taken it from experiment; I find however that as an
emetic, and probably therefore for other purposes, it is
equally efficacious.

TINCTURA SERPENTARIÆ:

TINCTURE OF VIRGINIAN SNAKE-
ROOT.

Take of Virginian snake-root, w. three
ounces,

weak spirit of wine two pints.

Digest them for eight days, and strain the
liquor.

TINCTURA VALERIANÆ.

TINCTURE OF VALERIAN.

Take of the root of wild valerian in gross
powder, w. four ounces,

weak spirit of wine two pints.

Digest them in a gentle heat for eight days,
and strain the liquor.

TINCTURA

TINCTURA VALERIANÆ VOLATILIS.

VOLATILE TINCTURE OF VALERIAN.

Take of the root of wild valerian, w. four
 ounces,
 compound spirit of ammonia two
 pints.

Digest them for eight days, and strain the
 liquor.*

NOTE.

All these tinctures should be made in stopped
 vessels, excepting muriated tincture of iron.†

N MISTURÆ.

* There being only one drachm of the valerian root in
 an ounce measure of the liquor, I cannot conceive that it
 can add any thing to the volatile alkali, if we even give
 two drachms for a dose.

† Those of them which are ordered to be made with
 artificial heat should be corked close, but not firm, that if
 the heat should be raised to too great a degree, the cork
 may rather fly out than the bottle burst. Those which
 are not ordered to be made with artificial heat, but are
 left to the accidental heat of the atmosphere, may be
 made in vessels stopped close with glass stoppers, or corked
 tight.

M I S T U R Æ.

M I X T U R E S.

MISTURA CAMPHORATA.

CAMPHORATED MIXTURE.

TAKE of camphor, w. one drachm,
rectified spirit of wine ten drops,
double-refined sugar, w. half
an ounce,
distilled water boiling hot one
pint.

Rub the camphor first in the spirit of wine,
then with the sugar, lastly with the water
poured in by a little at a time, and strain the
liquor.

MISTURA

MISTURA CRETACEA.

CHALK MIXTURE.

Take of prepared chalk, w. one ounce,
double-refined sugar, six drachms,
gum arabic powdered, w. two
ounces,
distilled water two pints.

Mix them together.

MISTURA MOSCHATA.

MUSK MIXTURE.

Take of rose water, w. six ounces,
musk, w. two scruples,
gum arabic powdered,
sugar, of each, w. one drachm.

Rub the musk first with the sugar, then
with the gum, and lastly with the rose water
added by a little at a time.

LAC AMYGDALÆ.

ALMOND MILK.

Take of sweet almonds, w. one ounce and
an half,
double-refined sugar, w. half an
ounce,
distilled water two pints.

Beat the almonds with the sugar; then,
rubbing them together, pour in the water by
a little at a time, and strain the liquor.*

LAC AMMONIACI.

MILK OF AMMONIAC.

Take of ammoniac, w. two drachms,
distilled water half a pint.

Rub the gum resin with the water, added
by a little at a time, until it is formed into a
milky liquor.

In

* It is to be strained through a sieve to take out the
husks.

In the same manner milk of asa foetida, and of other gum resins, may be formed.

SPIRITUS ÆTHERIS VITRIOLICI
COMPOSITUS.

COMPOUND SPIRIT OF VITRIOLIC
ÆTHER.

Take of vitriolic æther two pounds,
oil of wine, w. three drachms.

Mix them together.

SPIRITUS AMMONIÆ COMPOSITUS.
COMPOUND SPIRIT OF AMMONIA.

Take of spirit of ammonia two pints,
essential oil of lemons,
essential oil of nutmegs, of each,
w. two drachms.

Mix them together.

SPIRITUS AMMONIÆ SUCCINATUS.
SPIRIT OF AMMONIA WITH AMBER.

Take of alkohol, w. one ounce,
water of pure ammonia, m. four
ounces,
rectified oil of amber, w. one scruple,
soap ten grains.

Digest the soap and the oil of amber in the
alkohol until they are dissolved. Then pour
into them the water of pure ammonia, and
shake them till they are mixed.

SPIRITUS CAMPHORATUS.
CAMPHORATED SPIRIT.

Take of camphor, w. four ounces,
rectified spirit of wine two pints.

Mix them together that the camphor may
be dissolved.

SYRUPI

S Y R U P I.

S Y R U P S.

IN making syrups, where neither the prescription for the weight of the sugar, nor the manner of dissolving it are ordered, the following is the process.

Take of double-refined sugar, w. twenty-nine ounces,

the liquor, whatever it is, one pint,

Dissolve the sugar in the liquor, in a water bath; let the solution stand for twenty-four hours. Take off the scum, and pour off the syrup from the dregs, if there are any.

SYRUPUS ALTHÆÆ.

SYRUP OF MARSH-MALLOWS.

Take of fresh marsh-mallow roots, bruised,
one pound,
double-refined sugar four pounds,
distilled water one gallon.

Boil the marsh-mallow root in the water till there remains half a gallon. Let it stand to cool, and press out the liquor. Put it by for twelve hours, that the dregs may subside, and pour off the liquor. Add the sugar, and boil the whole till there are six pounds weight left.

SYRUPUS CARYOPHILLI RUBRI.

SYRUP OF CLOVE JULY-FLOWER.

Take of the fresh flowers of clove july-
flowers, having cut off the points,
two pounds,
distilled water boiling hot six pints.

Let

Let them stand together in a glass vessel for twelve hours, strain the liquor, and dissolve in it double-refined sugar so as to make a syrup.

SYRUPUS CORTICIS AURANTII.

SYRUP OF ORANGE-PEEL.

Take of the fresh exterior rind of Seville oranges, w. eight ounces,
distilled water, boiling hot, five pints.

Let them stand together in a covered vessel for twelve hours; strain the liquor, and make it into a syrup with sugar.

SYRUPUS CROCI.

SYRUP OF SAFFRON.

Take of saffron, w. one ounce,
distilled water boiling hot one pint.

Let them stand together for twelve hours in a covered vessel; strain the liquor, and
make

186 THE LONDON DISPENSATORY.

make it into a fyrup with double-refined
fugar.

SYRUPUS SUCCI LIMONIS.

SYRUP OF LEMON JUICE.

Take of lemon juice which has stood till
the dregs have subfided, and has
been strained off, two pints,
double - refined fugar, w. fifty
ounces.

Make them into a fyrup.

In the fame manner may be made fyrup of
the juice of mulberries, raspberries, and black
currants.

SYRUPUS PAPAVERIS ALBI.

SYRUP OF WHITE POPPY.

Take of the dried capsules of white poppy,
after having taken out the feed,
three pounds and an half,
double-

double-refined fugar six pounds,
distilled water eight gallons.

Slice and bruise the capsules, then boil them in the water, in a bath of water saturated with common salt, till three gallons are evaporated. Squeeze out the liquor; evaporate it to about four pints, which strain while it is yet hot, first through a sieve, and then through fine flannel, and set it by for twelve hours that the dregs may subside. Pour off the clear liquor, boil it down to three pints, and make it into a syrup with the fugar.

SYRUPUS PAPAVERIS ERRATICI.

SYRUP OF WILD POPPIES.

Take of the fresh flower leaves of wild
poppies four pounds,
distilled water boiling hot four pints
and an half.

Put the boiling water in a vessel placed in a
water bath; stir in the flower leaves by a few
at

at a time. Then take the vessel out of the water bath, and let it stand for twelve hours; press out the liquor, and set it by until the dregs subside. Make the clear part into a syrup with double-refined sugar.

SYRUPUS ROSÆ.

SYRUP OF THE ROSE.

Take of the flower leaves of damascene roses dried, w. seven ounces,
double-refined sugar six pounds,
distilled water boiling hot four pints.

Pour the water upon the flower leaves, and let them stand together for twelve hours, and strain the liquor. Evaporate it to two pints and an half, and make it into a syrup with the sugar.

SYRUPUS SPINÆ CERVINÆ.

SYRUP OF BUCKTHORN.

Take of the juice of fresh and ripe buckthorn berries one gallon,
bruised

bruised ginger, w. one ounce,
pimento powdered, w. one ounce
and an half,
double-refined sugar seven pounds.

Set by the juice for some days that the dregs may subside; then strain it. Mix the ginger and pimento with a pint of it; let them stand together four hours, and strain the liquor. In the mean time boil down the remaining liquor to three pints; then add that part of the juice in which the ginger and pimento were steeped, and make the whole into a syrup with the sugar.*

SYRUPUS TOLUTANUS.

SYRUP OF TOLU.

Take of balsam of Tolu, w. eight ounces,
distilled water three pints.

Boil them for two hours. Let the liquor
stand

* The juice must be set by in a cold place, otherwise a fermentation will arise which will prevent the dregs from subsiding.

stand to cool, strain it, and make it into a syrup with double-refined sugar.

SYRUPUS VIOLÆ.

SYRUP OF VIOLETS.

Take of recent flower leaves of violets two pounds,
distilled water boiling hot five pints.

Let them stand together for twenty-four hours; let the liquor drain off through a fine linen cloth without pressure. Then make it into a syrup with double-refined sugar.

SYRUPUS ZINGIBERIS.

SYRUP OF GINGER.

Take of ginger bruised, w. four ounces,
distilled water boiling hot three pints.

Let them stand together four hours, strain off

off the liquor, and make it into a fyrup with double-refined fugar.*

* The form of a fyrup is very far from being proper, where any thing more is meant than to give sweetness, flavour, or colour to medicines. In the first place, the juices of fruits, such as buckthorn berries, are very unequal in the proportion of water to the medicated part. In the second place, they are very apt to spoil by fermenting; especially the fyrup of white poppies and the fyrup of violets, which should therefore be kept in as cold a place as possible.

M E L L A

M E L L A M E D I C A T A .

M E D I C A T E D H O N I E S .

M E L R O S Æ .

H O N E Y O F T H E R O S E .

TAKE of the flower leaves of red roses,
the yellow points being taken
off, and dried, w. four ounces,
distilled water boiling hot three
pints,
purified honey five pounds.

Let the rose leaves and water stand together
for six hours; strain off the liquor, mix it
with the honey, and boil it to the thickness
of a syrup.

M E L

MEL SCILLÆ.

HONEY OF THE SQUILL.

Take of clarified honey three pounds,
tincture of squills two pints.

Boil them in a glass vessel to the thickness
of a syrup.

OXYMEL ÆRUGINIS.

OXYMEL OF VERDIGRIS.

Take of prepared verdigris, w. one ounce,
vinegar, m. seven ounces,
clarified honey, w. fourteen ounces.

Dissolve the verdigris in the vinegar. Strain
the liquor through a linen cloth, pour it into
the honey, and boil it to a proper thickness.

OXYMEL COLCHICI.

OXYMEL OF COLCHICUM.

Take of fresh root of colchicum cut in thin
slices, w. one ounce,

O

distilled

distilled vinegar one pint,
clarified honey two pounds.

Let the root stand with the vinegar in a glass vessel in a gentle heat for eight and forty hours. Squeeze out the liquor in a strong press, strain it, and mix the honey with it. Boil it to the thickness of a syrup, stirring it with a wooden spoon.

OXYMEL SCILLÆ.

OXYMEL OF THE SQUILL.

Take of clarified honey three pounds,
vinegar of the squill two pints.

Boil them in a glass vessel with a moderate heat to the thickness of a syrup.

OXYMEL SIMPLEX.

SIMPLE OXYMEL.

Take of clarified honey two pounds,
distilled vinegar one pint.

Boil

Boil them in a glass vessel with a moderate heat to the thickness of a syrup.*

* These medicated honies (excepting honey of the rose, and oxymel of verdigris) being employed as expectorants, the honey was supposed to assist them as such more than sugar. It must be remembered, however, that honey is even more apt to ferment than sugar; they ought therefore to be kept in a cold place.

It is very difficult to make honies or syrups in glass vessels. The heat of boiling syrup being above 230° of Fahrenheit's thermometer, it can hardly be obtained in a water bath. A wide mouthed open vessel fit for evaporation can hardly be well managed in a sand heat. A silver vessel is too expensive. But an iron vessel plated with silver, or an iron vessel kept very clean, might be employed, provided the syrup or honey be poured out as soon as it is made. A porcelain or stone-ware vessel may be employed in a sand-heat, or even over a gentle charcoal fire, at the distance of six or eight inches, especially porcelain and common brown stone earthen-ware.

No stone ware glazed with lead, or any glazing which has in its composition more lead than flint-glass has, can be used with safety.

Powder the aloës and the guaiacum separately, then mix the whole together.*

PULVIS ALOETICUS CUM FERRO.

ALOETIC POWDER WITH IRON.

Take of focotorine aloës powdered, w. one
ounce and an half,
myrrh powdered, w. two ounces,
dried extract of gentian,
vitriolated iron, of each powdered,
w. one ounce.

Mix them together.

PULVIS AROMATICUS.

AROMATIC POWDER.

Take of the bark of cinnamon, w. two
ounces,

O 3 lesser

* I have never been able to find good evidence of any use arising from a mixture of guaiacum and aloës. Purgatives generally require spices, or other stimulants, to prevent them from giving pain, or rendering the patient sick; especially those which have been called drastic purges.

leffer cardamom seeds, freed from
the husks,

ginger,

long pepper, of each, w. one
ounce.

Powder them together.*

PULVIS ASARI COMPOSITUS.

COMPOUND POWDER OF ASARUM.

Take of asarum leaves dried,

marjoram,

marum Syriacum,

flowers of lavender dried, of each,
w. one ounce.

Powder them together.

PULVIS E CERUSSA.

POWDER OF WHITE LEAD.

Take of white lead, w. five ounces,

farcocolla,

* I have already observed the advantages of mixing
spices.

farcocolla, w. one ounce and an
half,

tragacanth, w. half an ounce.

Powder them together.

PULVIS E CHELIS CANCRORUM
COMPOSITUS.

COMPOUND POWDER OF CRABS
CLAWS.

Take of prepared crabs claws one pound,
prepared chalk,
red coral prepared, of each, w.
three ounces.

Mix them together.*

PULVIS CONTRAYERVÆ COMPOSI-
TUS.

COMPOUND POWDER OF CONTRA-
YERVA.

Take of contrayerva powdered, w. five
ounces,

O 4 compound

* I do not know any reason for mixing mild calcareous
earth procured from different sources.

200 THE LONDON DISPENSATORY.

compound powder of crabs claws
one pound and an half.

Mix them together.

PULVIS E CRETA COMPOSITUS.
COMPOUND POWDER OF CHALK.

Take of prepared chalk half a pound,
cinnamon bark, w. four ounces,
tormentilla,
gum arabic, of each, w. three
ounces,
long pepper, w. half an ounce.

Powder them separately, then mix them
together.

PULVIS E CRETA COMPOSITUS CUM
OPIO.

COMPOUND POWDER OF CHALK WITH
OPIUM.

Take of compound powder of chalk, w.
eight ounces,

powder

powder of hard purified opium one
drachm and an half.

Mix them together.

PULVIS IPECACUANHÆ COMPOSITUS.

COMPOUND POWDER OF IPECACUAN.

Take ipecacuan powdered,

hard purified opium powdered, of
each, w. one drachm,

vitriolated kali powdered, w. one
ounce.

Mix them together.

PULVIS E MYRRHA COMPOSITUS.

COMPOUND POWDER OF MYRRH.

Take myrrh,

favin dried,

rue dried,

Ruffian

Ruffian castor, of each, w. one
ounce.

Powder them together.*

PULVIS OPIATUS.

OPIATE POWDER.

Take powder of hard purified opium, w.
one drachm,

burnt and prepared hartshorn, w.
nine ounces.

Mix them together.

PULVIS

* These substances are difficultly powdered separately, and therefore most probably ordered to be mixed before they are powdered. Care therefore should be taken that the whole mass is passed through the sieve.

In general when powders are to be mixed together, a sufficient quantity of each is passed through the sieve, leaving the grosser part to be employed with a fresh quantity of the matter for a new operation; which in many cases is a bad practice, as the most efficacious remedy is sometimes easiest, sometimes the most difficult of being powdered. In either case the powder will be various in its powers. The whole therefore of any mass of matter should be entirely passed through the sieve in all cases.

PULVIS E SCAMMONIO COMPOSITUS.

COMPOUND POWDER OF SCAMMONY.

Take scammony,

hard extract of jalap, of each, w:
two ounces,

ginger, w. half an ounce.

Powder them separately, and mix them together.

PULVIS E SCAMMONIO CUM ALOE.

POWDER OF SCAMMONY WITH ALOES.

Take of scammony, w. six drachms,

hard extract of jalap,

focotorine aloës, of each, w. one
ounce and an half,

ginger, w. half an ounce.

Powder

Powder them separately, and mix them together.*

PULVIS E SCAMMONIO CUM CALOMELANE.

POWDER OF SCAMMONY WITH CALOMEL.

Take of scammony, w. half an ounce,
calomel,

double-

* Of all mixtures of medicines similar in their action, or employed with the same intention, that of purgatives has the most evident good effect, performing the operation easier, with less pain and sickness, and with greater certainty. And this often goes even to mixing ten or twelve; such a mixture often having much greater effect than could be expected from the doses of each of the ingredients.

This probably depends on what I have already remarked, that the stomach will bear a larger quantity of two medicines of the same intention without their becoming simple stimulants, than it will of either of them alone.

It is probable that the great composition which had crept into antient, as well as modern prescriptions, arose from this quality being remarked in purgatives, in which it is so evident; especially as purgatives were much used in Grecian practice,

double-refined sugar, of each, w.
two drachms.

Powder them separately, and mix them together.

PULVIS E SENNA COMPOSITUS.

COMPOUND POWDER OF SENNA.

Take senna,

crystals of tartar, of each, w. two
ounces,

scammony, w. half an ounce,
ginger, w. two drachms.

Powder them together, except the scammony, which is to be separately powdered, and afterwards mixed with the other ingredients.

PULVIS E TRAGACANTHA COMPOSITUS.

COMPOUND POWDER OF TRAGACANTH.

Take tragacanth powdered,

gum

gum arabic,
starch, of each, w. one ounce and
an half,
double-refined sugar, w. three
ounces.

Powder them together.*

* I have not been able to find proof of any advantage arising from the mixture of inspid mucilages with one another. In stiffening linen, such a mixture of farinaceous matter and gum gives a different consistence from either of them alone; the tragacanth being to be considered rather as farinaceous matter than gum. But I can find no authority of any advantage gained by it in medicine.

TROCHISCI.

T R O C H I S C I.
L O Z E N G E S.

TROCHISCI EX AMYLO.

STARCH LOZENGES.

TAKE of starch, w. one ounce and an
half,
liquorice, w. six drachms,
orris, w. half an ounce,
double-refined sugar, one pound
and an half.

Powder them, and make them into lozenges
with a solution of tragacanth in water.

They may also be made without the orris.

TROCHISCI

TROCHISCI E CRETA.

CHALK LOZENGES.

Take of prepared chalk, w. four ounces,
prepared crabs claws, w. two
ounces,
cinnamon bark, w. half an ounce,
double-refined sugar, w. three
ounces.

Powder them, and make them into lozenges
with mucilage of gum arabic.

TROCHISCI E GLYCYRRHIZA.

LIQUORICE LOZENGES.

Take extract of liquorice,
double-refined sugar, of each, w.
ten ounces,
powder of tragacanth, w. three
ounces.

Make them into lozenges with water.

TROCHISCI

TROCHISCI E MAGNESIA.

LOZENGES OF MAGNESIA.

- Take of burnt magnesia, w. four ounces,
double-refined sugar, w. two
ounces,
ginger powdered, w. one scruple.

Make them into lozenges with mucilage of
gum arabic.

TROCHISCI E NITRO.

LOZENGES OF NITRE.

- Take of purified nitre powdered, w. four
ounces,
double-refined sugar powdered,
one pound,
powder of tragacanth, w. six
ounces,

Make them into lozenges with water.

P

TROCHISCI

TROCHISCI E SULPHURE.

LOZENGES OF SULPHUR.

Take of washed flowers of sulphur, w. two
ounces,

double - refined sugar, w. four
ounces,

Rub them together, and make them into
lozenges with mucilage of quince seed.

PILULÆ.

P I L L U L Æ.

P I L L S.

PILULÆ EX ALOE.

ALOETIC PILLS.

TAKE of focotorine aloës in powder,
 w. one ounce,
 extract of gentian, w. half an
 ounce,
 fyrup of ginger, as much as is
 necessary to give the consistence
 of pills.

Beat them together.

PILULÆ EX ALOE CUM MYRRHA.

ALOETIC PILLS WITH MYRRH.

Take of focotorine aloes, w. two ounces,

P 2

myrrh,

myrrh,
 saffron, of each, w. one ounce,
 fyrup of saffron a sufficient quantity
 to give the consistence of pills.

Powder the aloës and myrrh separately, then
 beat the whole thoroughly together.*

PILULÆ E GUMMI.

GUM PILLS.

Take Galbanum,
 opopanax,
 myrrh,
 sagapenam, of each, w. one ounce,
 asa fætida, w. half an ounce,
 fyrup

* Saffron is retained by the College in several compositions, although many practitioners now consider it as a medicine of no efficacy; it being used in feasts in Indostan in much larger quantity than ever was employed in medicine, without any apparent effect. I cannot however put my judgment in competition with the College, only in pills encreasing the bulk is a great inconvenience. Unless therefore the saffron should be thought essential, I should be led to believe that it was better without it.

fyrup of faffron, as much as is
 necessary to give the consistence
 of pills.

Beat them thoroughly together.

PILULÆ EX HYDRARGYRO.

QUICKSILVER PILLS.

Take purified quicksilver,

extract of liquorice of the consistence
 of honey, of each, w. two
 drachms,

liquorice in fine powder, w. one
 drachm.

Rub the quicksilver with the extract of
 liquorice until no globules are to be seen; then
 add the liquorice powder, and mix the whole
 together.*

P 3

PILULÆ

* Many viscid fluids, both oily and watery, have been
 employed to divide mercury so as to render it active, or
 according to the vulgar phrase to kill it. It does not seem
 to be of consequence which is used, provided the quick-
 silver

PILULÆ EX OPIO.

OPIUM PILLS.

Take of hard purified opium, w. two
drachms,

extract of liquorice, w. one ounce.

Beat them thoroughly together.

PILULÆ E SCILLA.

SQUILL PILLS.

Take of fresh dried squills in powder, w.
one drachm,

ginger in powder,

soap, of each, w. three drachms,

gum ammoniac, w. two drachms,

fyrup

silver be divided with sufficient minuteness; which it is not, if it can be seen with a microscope that magnifies the diameter fifteen or twenty times.

Care is to be taken not to rub or beat the mass after putting in the powder too much, or till it becomes at all dry, as that would occasion the globules to unite again and become inactive.

fyryp of ginger, as much as is
necessary to give the consistence
of pills.

Beat them thoroughly together.*

* Care should be taken that the soap be perfectly in-
fipid, as, if there be any superfluous alkali, it will spoil
the squills.

ELECTUARIA.

ELECTUARIES.

ELECTUARIUM E CASSIA.

ELECTUARY OF CASSIA.

TAKE of pulp of cassia recently extracted,
w. half a pound,
manna, w. two ounces,
pulp of tamarinds, w. one ounce,
syrup of roses half a pound.

Bruise the manna, and dissolve it in the
syrup in a moderate heat; then mix in the
pulp, keep up the heat, and evaporate until
the electuary has acquired a proper consistence.

ELEC-

ELECTUARIUM E SCAMMONIO.

ELECTUARY OF SCAMMONY.

Take of scammony in powder, w. one ounce
and an half,

cloves,

ginger, of each six drachms,

essential oil of caraway, w. half a
drachm,

syrop of the rose, a sufficient quan-
tity to form an electuary, by stirring into it
first the spices mixed together, then the scam-
mony, and lastly the oil of caraway.

ELECTUARIUM E SENNA.

ELECTUARY OF SENNA.

Take of senna, w. eight ounces,

figs one pound,

pulp of tamarinds,

caffia,

French

French prunes, of each
half a pound,
coriander seeds, w. four ounces,
liquorice, w. three ounces,
double-refined sugar two pounds
and an half.

· Powder the fenna and coriander seeds together, sift out ten ounces. Boil what remains in the sieve in four pints of distilled water till half is consumed, then press out the liquor and strain it. Afterwards evaporate it till there remains about one pound and an half. Add to it the sugar, so as to make a syrup. Mix this syrup by a little at a time with the pulps, and lastly stir in the powder obtained from the fenna and coriander seeds.

CON-

CONFECTIONES.
CONFECTIONS.

CONFECTIO AROMATICA.
AROMATIC CONFECTION.

TAKE zedoary grosly powdered,
saffron, of each half a pound,
distilled water three pints.

Let them stand together for four and twenty hours, then press out and strain the liquor. Evaporate it until a pint and a half is left, and mix it with the following ingredients rubbed into fine powder.

Compound powder of crabs claws,
w. sixteen ounces,

bark of cinnamon,
nutmeg, of each, w. two ounces,
cloves, w. one ounce,

leffer

lesser cardamom seeds, freed from
the husk, w. half an ounce,
double-refined sugar two pounds,
So as to make a confection.*

CONFECTIO OPIATA.

OPIATE CONFECTION.

Take powder of hard purified opium, w.
six drachms,
long pepper,
ginger,
caraway seeds, of each, w. two
ounces,
syrup of white poppies, boiled
down to the consistence of ho-
ney, three times the weight of
all the other ingredients.

Warm the syrup, and mix in the opium as
equally as possible, and afterwards the other
ingredients powdered.

AQUÆ

* I have not noticed in general the improvements
which the College have made, where they have been so
obvious as avoiding the throwing away the essential oil
from Raleigh's confection.

AQUÆ MEDICATÆ.

MEDICATED WATERS.

AQUA ALUMINIS COMPOSITA.

COMPOUND ALUM WATER.

TAKE alum,
vitriolated zinc, of each, w. half
an ounce,
distilled water boiling hot two
pints.

Pour the water upon the salts in a glass ves-
sel, and strain the liquor.

AQUA CUPRI AMMONIATI.

WATER OF AMMONIATED COPPER.

Take of lime water one pint,
sal ammoniac, w. one drachm.

Let

Let them stand together in a copper vessel till the ammoniac is saturated.

AQUA LITHARGYRI ACETATI COM-
POSITA.

COMPOUND WATER OF ACETATED
LITHARGE.

Take of water of acetated litharge, w. two
drachms,

distilled water two pints,

weak spirit of wine, M. two
drachms.

Mix the spirit of wine with the water of
acetated litharge; then add the distilled water.

AQUA ZINCI VITRIOLATI CUM
CAMPORA.

WATER OF VITRIOLATED ZINC WITH
CAMPHOR.

Take of vitriolated zinc, w. half an ounce,
camphorated

MEDICATED WATERS. 223

camphorated spirit, M. half an
ounce,
water boiling hot two pints.

Mix them together, and strain them through
paper.

EMPLASTRA.

E M P L A S T R A.
P L A I S T E R S.

EMPLASTRUM AMMONIACI CUM
HYDRARGYRO.

AMMONIAC PLAISTER WITH QUICK-
SILVER.

TAKE of strained gum ammoniac one
pound,
purified quicksilver, w. three
ounces,
fulphurated oil, w. one drachm,
or more if necessary.

Rub the quicksilver with the fulphurated
oil until its globules are no longer visible.
Then melt the gum ammoniac, pour it in by a
little at a time, and mix them thoroughly to-
gether.*

EMPLAS-

* Care should be taken to employ as little heat as pos-
sible in melting the ammoniac.

EMPLASTRUM CANTHARIDIS.

PLAISTER OF THE CANTHARIS.

Take of cantharides one pound,
wax plaister two pounds,
hogs lard prepared half a pound.

Reduce the cantharides to a very fine powder; melt the plaister and the lard together, remove them from the fire, let them stand till they are just fluid, then sprinkle and stir into them the powder of the cantharides.*

Q

EMPLAS-

* It is of great consequence that blisters should be certain in their operation, it happening frequently in acute diseases that a loss even of a few hours is very material. Their failure is more frequently owing to the cantharides not being good, than to any other cause. There are two insects which prey upon them, and not only destroy part of them, but by inducing moisture occasion the remaining part to putrify. These insects attack the heads and extremities, so that such cantharides should be chosen as are most perfect, dry and light, and the apothecary should try each parcel in some chronic case, where it is not of much consequence if they should fail, such as in chronic rheumatism.

EMPLASTRUM CERÆ.

WAX PLAISTER.

Take yellow wax,
prepared mutton suet, of each three
pounds,
yellow rosin one pound.

Melt them together, remove them from
the fire, and strain the mass while it is yet
fluid.

EMPLASTRUM CUMINI.

CUMIN PLAISTER.

Take cumin seeds,
caraway seeds,
laurel berries, of each, w. three
ounces,
Burgundy pitch three pounds,
yellow wax, w. three ounces.

Melt the pitch and the wax together, and
fir

stir in the other ingredients powdered, so as to form the whole into a plaister.

EMPLASTRUM LADANI.

LADANUM PLAISTER.

Take of ladanum, w. three ounces,
frankincense, w. one ounce,
cinnamon bark in powder,
the expressed oil called oil of mace,
of each, w. half an ounce,
essential oil of mint, w. one drachm.

Melt the frankincense, and mix it first with the ladanum heated till it has become of a soft texture, then the oil of mace, and lastly beat the mixture with the cinnamon and oil of mint in a warm mortar into a plaister. Keep it in a covered vessel.*

Q 2

EMPLAS-

* I have seen in Mesue, and other authors, a number of ointments and plaisters applied externally with a view to affect the stomach and intestines, even some to act as purgatives. Most of these have fallen out of use, as it is easy

to

EMPLASTRUM LYTHARGYRI.

PLAISTER OF LITHARGE.

Take of litharge in fine powder five pounds,
oil of olives one gallon.

Boil them with a moderate fire in about two pints of water, stir them constantly, adding more water when any considerable portion has evaporated, until the oil and litharge unite, and become of the consistence of a plaister.

EMPLASTRUM LYTHARGYRI CUM
GUMMI.

PLAISTER OF LITHARGE WITH GUM.

Take of litharge plaister three pounds,
strained

to see that there is no passage from the skin to the intestines for any solid, fluid, or vapour, excepting by the ordinary course of absorption and circulation. I suppose, however, that the College had ground from experience for believing that good effects might be produced in cases of flatulency by this application, it being the only one they have left with this intention.

strained galbanum, w. eight ounces,
turpentine, w. ten drachms,
frankincense, w. three ounces.

Melt the galbanum and turpentine together with a moderate heat; stir in the frankincense, having first powdered it; then melt the litharge plaister with as little heat as possible, and mix them all together so as to form a plaister.

EMPLASTRUM LITHARGYRI CUM
HYDRARGYRO.

LITHARGE PLAISTER WITH QUICK-
SILVER.

Take of litharge plaister one pound,
purified quicksilver, w. three ounces,
fulphurated oil, w. one drachm, or
more if necessary.

Make this plaister in the manner that ammoniac plaister with quicksilver is ordered to be made.

Q 3

EMPLAS-

EMPLASTRUM LITHARGYRI CUM
RESINA.

LITHARGE PLAISTER WITH ROSIN.

Take of litharge plaister three pounds,
yellow rosin half a pound.

Powder the rosin, and stir it into the li-
tharge plaister, melted with as little heat as
possible.

EMPLASTRUM PICIS BURGUNDICÆ.
BURGUNDY PITCH PLAISTER.

Take of Burgundy pitch two pounds,
ladanum one pound,
yellow rosin,
yellow wax, of each, w. four
ounces,
expressed oil commonly called oil
of mace, w. one ounce.

Melt the pitch, rosin, and wax together;
then

then mix in with them first the ladanum, then the oil of mace.

EMPLASTRUM SAPONIS.

SOAP PLAISTER.

Take of soap half a pound,
litharge plaister three pounds.

Melt the litharge plaister, stir the soap into it, and boil it to the consistence of a plaister.

EMPLASTRUM THURIS.

FRANKINCENSE PLAISTER.

Take of frankincense half a pound,
dragon's blood, w. three ounces,
litharge plaister two pounds.

Melt the litharge plaister, and add to it the other ingredients in powder.

UNGUENTA ET LINI-
MENTA.

OINTMENTS AND LINIMENTS.

UNGUENTUM ADIPIS SUILLÆ.

OINTMENT OF HOGS LARD.

TAKE of prepared hogs lard two pounds,
rose water, M. three ounces.

Beat the lard with the rose water until they are thoroughly mixed, then melt the whole with a gentle heat, and set them by till the water subsides. Lastly pour off the lard from the water, and stir it continually until it becomes cold.

UNGUENTUM

UNGUENTUM CALCIS HYDRARGY-
RI ALBÆ.

OINTMENT OF WHITE CALX OF
QUICKSILVER.

Take of white calx of quicksilver, w. one
drachm,
ointment of hogs lard, w. one
ounce and an half.

Mix them together so as to form an oint-
ment.

UNGUENTUM CANTHARIDIS.

OINTMENT OF CANTHARIDES.

Take of cantharides powdered, w. two
ounces,
distilled water, M. eight ounces,
ointment of yellow rosin, w. eight
ounces.

Boil the cantharides and the water till half
of it is evaporated, and strain it. Mix the
strained

strained liquor with the ointment, and evaporate it in a water bath saturated with salt, to the consistence of an ointment.

UNGUENTUM CERÆ.

OINTMENT OF WAX.

Take of white wax, w. four ounces,
sperma ceti, w. three ounces,
olive oil one pint.

Melt them together with a gentle heat, and stir them constantly and quickly until they become cold.

UNGUENTUM CERUSSÆ ACETATÆ.

OINTMENT OF ACETATED CERUSSE.

Take of acetated cerusse, w. two drachms,
white wax, w. two ounces,
olive oil half a pint.

Rub the acetated cerusse to a powder, which is to be rubbed into part of the oil. Then
melt

melt the wax with the remaining part of the oil. Mix the whole together, and stir them till they cool.

UNGUENTUM ELEMI.

OINTMENT OF ELEMI.

Take of elemi one pound,
 turpentine, w. ten ounces,
 prepared mutton suet two pounds,
 olive oil, M. two ounces.

Melt the elemi and suet together, take them off the fire, and mix in the turpentine and oil immediately; lastly strain the mixture.

UNGUENTUM HELLEBORI ALBI.

OINTMENT OF WHITE HELLEBORE.

Take of white hellebore powdered, w. one
 ounce,
 ointment of hogs lard, w. four
 ounces,

essence

essence of lemon, w. half a scruple.

Mix them together so as to form an ointment.

UNGUENTUM HYDRARGYRI FORTIUS.

STRONG QUICKSILVER OINTMENT.

Take of purified quicksilver two pounds,
prepared hogs lard, w. twenty-
three ounces,
prepared mutton suet, w. one
ounce.

Rub the quicksilver first with the suet, and a little of the lard, until the globules are no longer visible, then add the remaining part of the lard so as to form an ointment.*

UNGUEN-

* I cannot help remarking again the necessity of dividing the quicksilver into very fine particles, where it is rubbed down with any viscid fluid. This is still more essentially necessary when employed in the form of an ointment. The College have here preferred mutton suet to turpentine, or sulphurated oil, which have been formerly

UNGUENTUM HYDRARGYRI MI-
TIUS.

WEAK QUICKSILVER OINTMENT.

Take of strong quicksilver ointment one
part,

prepared hogs lard two parts.

Mix them together.*

UNGUENTUM

merly used, I suppose partly on account of their stimulating the skin, and perhaps also of their smell. It requires, however, more care in the rubbing with the fuet. Suet that has been kept for some years is more efficacious.

* In this instance the College have ordered proportional parts, instead of specific quantities of the different ingredients, which is the practice commonly with Celsus, and many other great authorities. But it is liable to several objections, particularly that it is difficult to specify each ingredient either by weight or by measure. If one be taken by weight, and others by measure, the prescription will be very vague.

UNGUENTUM HYDRARGYRI NI-
TRATI.

OINTMENT OF NITRATED QUICK-
SILVER.

Take of purified quicksilver, w. one ounce,
nitrous acid, w. two ounces,
prepared hogs lard one pound.

Dissolve the quicksilver in the nitrous acid,
melt the suet with as little heat as possible,
pour it into the solution while it is yet hot,
stir them together till they are cold.

UNGUENTUM PICIS.

TAR OINTMENT.

Take tar,
prepared mutton suet, of each half
a pound.

Melt them together, and strain them.

UNGUENTUM

UNGUENTUM RESINÆ FLAVÆ.

OINTMENT OF YELLOW ROSIN.

Take yellow rosin,

yellow wax, of each one pound,
oil of olives one pint.

Melt the rosin and the wax with a moderate heat, then pour in the oil, and strain the whole while they are yet hot.

UNGUENTUM SAMBUCI.

OINTMENT OF ELDER.

Take of elder flowers four pounds,

prepared mutton suet three pounds,
olive oil one pint.

Melt the suet with the oil, throw in the flowers, and boil them until they are crisp; press out the ointment, and strain it.

UNGUEN-

UNGUENTUM SPERMATIS CETI.

SPERMA CETI OINTMENT.

Take of sperma ceti, w. six drachms,
white wax, w. two drachms,
olive oil, M. three ounces.

Melt them together with a moderate heat,
remove them from the fire, stir them con-
stantly and briskly until they become cold.

UNGUENTUM SULPHURIS.

OINTMENT OF SULPHUR.

Take of ointment of hogs lard half a pound,
flowers of sulphur, w. four ounces.

Mix them together so as to form an oint-
ment.

UNGUENTUM TUTIÆ.

OINTMENT OF TUTTY.

Take prepared tutty,

spermaceti

spermaceti ointment a sufficient
quantity to form a soft ointment.

Mix them together.

LINIMENTUM AMMONIÆ.

LINIMENT OF AMMONIA.

Take of water of ammonia, M. half an
ounce,
oil of olives, M. one ounce and
an half.

Shake them together in a phial till they are
well mixed.

LINIMENTUM AMMONIÆ FOR-
TIUS.

STRONG LINIMENT OF AMMONIA.

Take of water of pure ammonia, M. one
ounce,
olive oil, M. two ounces.

Shake them together in a phial.

R LINIMEN-

LINIMENTUM CAMPHORÆ.

CAMPHOR LINIMENT.

Take of camphor, w. two ounces,
water of ammonia, M. six ounces,
spirit of lavender, w. sixteen ounces.

Mix the water of ammonia with the spirit,
and distill over sixteen ounces in glass vessels,
with a moderate heat. Then dissolve the
camphor in the distilled liquor.

LINIMENTUM SAPONIS.

LINIMENT OF SOAP.

Take of soap, w. three ounces,
camphor, w. one ounce,
spirit of rosemary one pint.

Digest the soap in the spirit of rosemary, and
afterwards the camphor, till they are dissolved.

CERATA.

C E R A T A.

C E R A T E S.

CERATUM CANTHARIDIS.

CERATE OF THE CANTHARIS.

TAKE of cerate of sperma ceti, rendered soft by heat, w. six drachms,
 cantharides in fine powder, w. one drachm.

Mix them together.

CERATUM LAPIDIS CALAMINARIS.

CERATE OF CALAMINE.

Take prepared calamine,
 yellow wax, of each half a pound,
 oil of olives one pint.

R 2

Melt

Melt the wax and oil together, remove them from the fire, and as soon as the mixture begins to be of a tough consistence, stir in the calamine, and continue stirring till the whole becomes cold.

CERATUM LYTHARGYRI ACETATI.
CERATE OF ACETATED LITHARGE.

Take of water of acetated litharge M. two
ounces and an half,
yellow wax, w. four ounces,
oil of olives, M. nine ounces,
camphor, w. half a drachm.

Rub the camphor into a powder with a few drops of the oil. Melt the wax with the rest of the oil; remove the mixture from the fire, and as soon as it begins to acquire a viscid consistence, pour in by a little at a time the water of acetated litharge; stir the mass until it becomes cold, and lastly mix in the camphubbered down with the oil.

CERATUM

CERATUM RESINÆ FLAVÆ. ▲

CERATE OF YELLOW ROSIN.

Take ointment of yellow rosin half a pound,
yellow wax, w. one ounce.

Melt them together so as to form a cerate.

CERATUM SAPONIS.

CERATE OF SOAP.

Take of soap, w. eight ounces,
yellow wax, w. ten ounces,
litharge powdered one pound,
oil of olives one pint,
vinegar one gallon.

Boil the vinegar and the litharge with a moderate fire, stirring them frequently, until the mixture unites and becomes viscid; then stir in the other ingredients so as to form a cerate.

CERATUM

CERATUM SPERMATIS CETI.

CERATE OF SPERMA CETI.

Take of sperma ceti, w. half an ounce,
white wax, w. two ounces,
olive oil, M. four ounces.

Melt them together, remove them from
the fire, and stir them till the cerate becomes
cold,

EPITHE-

E P I T H E M A T A.
P O U L T I C E S.

CATAPLASMA CUMINI.

CUMIN POULTICE.

TAKE of cumin seed one pound,
laurel berries,
the leaves of scordium dried,
Virginian snake-root, of each, w.
three ounces,
cloves, w. one ounce.

Rub all of them together, and add three
times the weight of honey so as to make a
poultice.

CATAPLASMA SINAPEOS.

MUSTARD POULTICE.

Take mustard seed powdered,

R 4

crumb

248 THE LONDON DISPENSATORY.

crumb of bread, of each half a
pound,

vinegar a sufficient quantity to
form a poultice.

Mix them together.

COAGULUM ALUMINIS.

ALUM POULTICE.

Take the whites of two eggs.

Rub them with a piece of alum until they
are coagulated.

THE END OF THE DISPENSATORY.

TABLE

T A B L E,

Showing the Proportion of QUICKSILVER
and OPIUM in the compound Medicines
which contain them.

COMPOUND powder of chalk, with
opium, in forty-three grains contains
one grain of opium.

Compound powder of ipecacuan, in ten
grains contains one grain of opium.

Opiate powder, in ten grains contains one
grain of opium.

Powder of scammony with calomel, in
four grains contains one grain of calomel.

Opium pills, in five grains, contain one
grain of opium.

Quicksilver

Quicksilver pills, in ten grains contain four grains of quicksilver.

Opiate confection, in thirty six grains contains one grain of opium.

Ammoniac plaister with quicksilver, in five ounces contains one ounce of quicksilver.

Litharge plaister with quicksilver, in five ounces contains one ounce of quicksilver.

Strong quicksilver ointment, in two drachms contains one drachm of quicksilver.

Weak quicksilver ointment, in six drachms contains one drachm of quicksilver.

Ointment of nitrated quicksilver, in one drachm contains twelve grains of nitrated quicksilver.

Ointment of white calx of quicksilver, in one drachm contains four grains and an half of white calx of quicksilver.

T A B L E

T A B L E

Showing the former and present NAMES
where they have been changed.

FORMER NAMES.

PRESENT NAMES.

A

A CETUM scilliticum.	Acetum scillæ.
Æthiops mineralis.	Hydrargyrus cum sulphure.
Aqua aluminosa bateana.	Aqua aluminis composita.
calcis simplex.	calcis.
cinnamomi simplex.	cinnamomi.
spirituosa.	Spiritus cinnamomi.
hordeata.	Decoctum hordei.
juniperi composita.	Spiritus juniperi compositus.
menthæ piperitidis	Aqua menthæ piperitidis.
simplex.	
spirituosa.	Spiritus menthæ piperitidis
vulgaris simplex.	Aqua menthæ sativæ.
spirituosa.	Spiritus menthæ sativæ.
nucis moschataæ.	nucis moschataæ.
piperis jamaicensis.	Aqua pimento.
pulegii simplex.	pulegii.
spirituosa.	Spiritus pulegii.

raphani composita.	raphani compositus.
Aqua rosarum damascenarum.	Aqua rosæ.
sapphirina.	cupri ammoniati.
feminum anethi.	anethi.
anisi composita.	Spiritus anisi compositus.
carui.	carui.
vitriolica camphorata.	Aqua zinci vitriolati cum camphora.
Argenti vivi purificatio.	Hydrargyri purificatio.
Axungix porcinæ curatio.	Adipis fuillæ præparatio.

B.

Balsamum sulphuris barbadense.	Petroleum sulphuratum.
simplex.	Oleum sulphuratum.
traumaticum.	Tinctura benzoës composita.

C.

Calx antimonii.	Antimonium calcinatum.
Cauticum antimoniale.	muriatum.
commune fortius.	Calx cum kali puro.
lunare.	Argentum nitratum.
Ceratum album.	Ceratum spermatis ceti.
citrinum.	resinæ flavæ.
epuloticum.	lapidis calaminaris.
Chalybis rubigo præparata.	Ferri rubigo.
Cinnabaris factitia.	Hydrargyrus sulphuratus ruber.
	Confectio

WHICH HAVE BEEN CHANGED. 253

Confectio cardiaca.	Confectio aromatica.
Cornu cervi calcinatio.	Cornu cervi ustio.
D.	
Decoctum album.	Decoctum cornu cervi.
commune pro	pro enemate.
clystere.	horde compo-
pectorale.	tum.
E.	
Electarium lenitivum.	Electarium e fenna.
Elixir aloës.	Tinctura aloës composita.
myrrhæ compositum.	fabinæ composita.
paregoricum.	opii camphorata.
Emplastrum exammoniaco	Emplastrum ammoniaci
cum mercurio.	cum hydrargyro.
Emplastrum attrahens.	Emplastrum ceræ.
cephalicum.	picis burgundicæ.
commune.	lithargyri.
adhæsivum.	cum resina.
cum gummi.	cum gummi.
cum mercurio.	cum hydrat-
e cymino.	gyro.
roborans.	eumini.
e sapone.	thuris.
stomachicum.	saponis.
vesicatorium.	ladani.
Emulsiio communis.	cantharidis.
Extractum catharticum.	Lac amygdalæ.
	Extractum e colocynthide
	compositum.
	thebæicum.

thebäicum.

Opium purificatum.

F.

Flores benzöini.

Flores benzoës.

martiales.

Ferrum ammoniacale.

Fotus communis.

Decoctum pro formento.

H.

Hiera picra.

Pulvis aloëticus.

I.

Infusum amarum simplex.

Infusum gentianæ compo-
situm.

senæ commune.

fennæ tartarifatum.

Julepum e camphora.

Mistura camphorata.

e creta.

cretacea.

e moscho.

moschata.

L.

Linimentum album.

Unguentum spermatis ceti

saponaceum.

Linimentum saponis.

volatile.

ammoniæ.

Lixivium saponarium.

Aqua kali puri.

tartari.

kali.

M.

Mel ægyptiacum.

Oxymel æruginis.

rosaceum.

Mel rosæ

Mercurius calcinatus.

Hydrargyrus calcinatus

corrosivus sublimatus.

muriatus.

ruber.

nitratus ruber.

dulcis

WHICH HAVE BEEN CHANGED. 255

dulcis sublimatus.	Calomelas.
Mercurius emeticus flavus.	Hydrargyrus vitriolatus.
præcipitatus albus.	Calx hydrargyri alba.

N.

Nitrum vitriolatum.	Kali vitriolatum.
---------------------	-------------------

O.

Oleum petrolei barbadensis	Oleum petrolei.
terebinthinæ æthé-	terebinthinæ recti-
reum.	ficatum.
Opium colatum.	Opium purificatum.
Oxymel scilliticum.	Oxymel scillæ.

P.

Philonium londinense.	Confectio opiata.
Pilulæ aromaticæ.	Pulvis aloëticus cum guaia-
	co.
rufi.	Pilulæ ex aloë cum myrrha.
Pulvis e bolo compositus.	Pulvis e creta compositus.
cum opio.	cum opio.
e cerussa compositus.	e cerussa.
sternutatorius.	asarî compositus.

R.

Rob baccaram sambuci.	Succus baccæ sambuci
	spisatus.

S.

Saccharum faturni.	Cerussa acetata.
Sal absinthii.	Kali præparatum.
catharticus glauberi.	Natron vitriolatum.

Sal

Sal diureticus.	Kali acetatum.
martis.	Ferrum vitriolatum.
tartari.	Kali præparatum.
vitrioli.	Zincum vitriolatum.
volatilis falis ammoniaci.	Ammonia præparata.
Species aromaticæ.	Pulvis aromaticus.
Spiritus cornu cervi.	Liquor volatilis cornu cervi
lavendulæ compositus.	Tinctura lavendulæ composita.
simplex.	Spiritus lavendulæ.
nitri dulcis.	ætheris nitrosi.
glauberi.	Acidum nitrosum.
falis ammoniaci.	Aqua ammoniæ.
falis ammoniaci dulcis.	Spiritus ammoniæ.
falis marini glauberi.	Acidum muriaticum.
vinosus camphoratus.	Spiritus camphoratus.
vitrioli dulcis.	ætheris vitriolici.
volatilis aromaticus.	Spiritus ammoniæ compositus.
	fetidus.
	fetidus.
fetidus.	Succus cochleariæ compositus.
Succi scorbutici.	
Syrupus ex althæa.	Syrupus althææ.
e corticibus aurantium.	corticis aurantii.
balsamicus.	tolutanus.
e meconio.	papaveris albi.
rofarum solutivus.	rosæ.

T. Tabellæ

T.


Tabellæ cardialgicæ.	Trochisci e creta.
Tartarum emeticum.	Antimonium tartarifatum.
solubile.	Kali tartarifatum.
vitriolatum.	vitriolatum.
Tinctura amara.	Tinctura gentianæ compo-
	fita.
aromatica.	cinnamomi composita.
foetida.	asæ foetidæ.
guaiacina volatilis.	guaiaci.
japonica.	catechu.
martis in spiritu falis.	ferri muriati.
melampodii.	hellebori nigri.
rhabarbari spirituosa.	rhabarbari.
vinosa.	Vinum rhabarbari.
rosarum.	Infusum rosæ.
sacra.	Vinum aloës.
stomachica.	Tinctura cardamomi com-
	posita.
Trochisci bechici albi.	Trochisci ex amylo.
nigri.	e glycyrrhiza.

V.

Vinum antimoniale.	Vinum antimonii.
chalybeatum.	ferri.
Unguentum album.	Unguentum ceræ.
basilicum flavum.	resinæ flavæ.
cæruleum fortius.	hydrargyri fortius.
mitius.	mitius.
e mercuriopræcipitato.	calcis hydrargyri albæ.
saturinum.	cerussæ acetatæ.
simplex.	adipis suillæ.

S

INDEX



I N D E X
M E D I C A M E N T O R U M.

Under the names in the Latin Index the doses are set down in Roman characters. The quantities are meant for adults. It generally requires about a sixth part of the quantity for a child about one year old, and a full dose after fifteen. Where the dose for an infant is considerably different, it is marked by an I. put before the quantity. A. is placed before the quantity, when the medicine is not meant to vomit, or to act as a purgative; and E. when it is meant to vomit or to purge. The numbers 1, 2, 3, &c. in Arabic characters, denominate the number of times they are to be exhibited in the twenty-four hours. Those prescriptions, to whose names there is no dose affixed, are generally for external use.

A,		
A	CETUM distillatum ——— Page 68 scillæ, <i>olim</i> Acetum scilliticum 150 A. gr. xxx ad ʒj—4.	
	Acidum acetosum ——— 70 muriaticum, <i>olim</i> Spiritus falis marini glauberi 66 A. gr. xx ad gr. xxx—3.	Acidum
	S 2	

Acidum nitrosum, <i>olim</i> Spiritus nitri glauberi	65
———— dilutum, <i>vice</i> Aquæ fortis	66
vitriolicum dilutum, <i>vice</i> Elixir vitrioli acidum	65
A. gr. xxx ad 3 j—3.	
Adipis fuillæ preparatio, <i>olim</i> Axungiaë porcinaë curatio	39
Æruginis præparatio	37
Æther vitriolicus	129
A. gr. xx ad 3 j—4.	
Alkohol	127
Aluminis purificatio	89
A. gr. v ad gr. x—4 ad 6.	
Alumen ustum	89
A. gr. iv ad gr. viii—4 ad 6.	
Ammonia præparata, <i>olim</i> Sal volatilis falsi ammoniaci	78
A. gr. v ad gr. xv—4.	
Ammoniaci gummi purificatio	39
A. gr. v ad gr. x—4.	
Antimonii præparatio	38
A. gr. xx ad 3 j—3.	
Antimonium calcinatum, <i>olim</i> Calx antimonii	96
A. gr. iii—gr. xx—gr. xxx—4.	
muriatum, <i>vice</i> Caustici antimonialis	97
tartarisatum, <i>olim</i> Tartarum emeticum	100
A. gr. $\frac{1}{4}$ ad gr. $\frac{1}{3}$ —4 ad 6, E. gr. i ad gr. iii.	
vitrificatum	101
Aqua aluminis composita, <i>olim</i> Aqua aluminosa bateana	221
ammoniaë, <i>olim</i> Spiritus falsi ammoniaci	81
A. gr. xx ad gr. xxx—4. 3 fs ad 3 vi—1.	
ammoniaë acetataë, <i>vulgo</i> Spiritus mindereri	87
A. 3 ji ad 3 fs—4	
	Aqua

INDEX MEDICAMENTORUM. 261

Aqua ammoniæ puræ, <i>vulgo</i> Spiritus salis ammoniaci		
	cum calce	79
anethi, <i>olim</i> Aqua feminum anethi	—	123
I. 3j. P. R. N.		
calcis, <i>olim</i> Aqua calcis simplex	—	148
A. ℞s and ℞j—2.		
cinnamomi, <i>olim</i> Aqua cinnamomi simplex		123
A. ℞j.		
cupri ammoniati, <i>olim</i> Aqua sapphirina		221
distillata	—————	122
fœniculi	—————	124
A. ℞ij.		
kali, <i>olim</i> Lixivium tartari	—————	74
A. gr. xx ad xxx—3.		
kali puri, <i>olim</i> Lixivium saponarium	—	75
A. gr. x ad gr. xx ex infuso lini ℞iv—2.		
lithargyri acetati	—————	117
lithargyri acetati composita, <i>vulgo</i> Extractum		
	saturni	222
menthæ piperitidis, <i>olim</i> Aqua menthæ piperi-		
	tidis simplex	124
A. 3ij ad ℞s.		
menthæ sativæ, <i>olim</i> Aqua menthæ vulgaris		
	simplex	124
A. ℞j cum fs.		
pimento, <i>olim</i> Aqua piperis jamaicensis		125
A. ℞j.		
pulegii, <i>olim</i> Aqua pulegii simplex	—	125
A. ℞j cum fs.		
rosarum, <i>olim</i> Aqua rosarum damascenarum		126
zinci violati cum camphora, <i>olim</i> Aqua vi-		
	triolica camphorata	222
S 3	Argentum	

Argentum nitratum, <i>olim</i> Causticum lunare	102
Ase foetidæ purificatio	39
A. gr. v ad gr. x.	

C.

Calomelas, <i>olim</i> Mercurius dulcis sublimatus	111
A. gr. ii ad gr. iv—i. vel gr. i—2 ad 4. E. gr. x ad gr. xx. I. gr. fs—2.	
Calx cum kali puro, <i>olim</i> Causticum commune fortius	76
hydrargyri alba, <i>olim</i> Mercurius præcipitatus albus	114
Cassiae Fistularis pulpæ extractio	42
E. ʒj ad ʒij.	
Cataplasma cumini	247
sinapeos	247
Ceratum cantharidis	243
lapidis calaminaris, <i>olim</i> Ceratum epuloticum	243
Ceratum lithargyri acetati	244
resinæ flavæ, <i>olim</i> Ceratum citrinum	245
saponis	245
spermatis ceti, <i>olim</i> Ceratum album	246
Cerussa acetata, <i>olim</i> Saccharum saturni	117
A. gr. fs ad gr. iii—4.	
Chelarum cancrorum præparatio	38
A. ʒj ad ʒiv.	
Coagulum aluminis	248
Confectio aromatica, <i>vice</i> Confectionis cardiacæ	217
A. ʒj ad ʒj.	
opiata, <i>olim</i> Philonium londinense	220
A. gr. ix—3. gr. xx ad gr. xxx—7	

Conserva

INDEX MEDICAMENTORUM. 263

Conserva absinthii maritimi	—————	44
A. ʒ ss—3.		
aris	—————	45
A. ʒ ss—3.		
corticis exterioris aurantii hispalensis		44
A. ʒ ij—4.		
cynobati	—————	45
A. Ad libitum.		
lujulæ	—————	44
A. Ad libitum.		
pruni sylvestris	—————	45
E. ʒ j ad ʒ ij.		
rosæ rubræ	—————	44
scillæ	—————	46
A. ʒ ss ad ʒ j.		
Corallii præparatio	—————	38
A. ʒ j ad ʒ iv.		
Cornu cervi ustio	—————	40
A. ʒ j—4.		
Cretæ præparatio	—————	38
A. ʒ j ad ʒ iv.		
Crocus antimonii	—————	97

D.

Decoctum cornu cervi, olim Decoctum album		138
A. ʒ ss—4.		
corticis peruviani	—————	139
A. ʒ ij—4.		
pro enemate, olim Decoctum commune pro clystere		139
E. ʒ xii.		

S 4

Decoctum

Decoctum pro fomento, <i>olim</i> Fetus communis	140
hellebori	140
hordei, <i>olim</i> Aqua hordeata	141
Pro potu.	
hordei compositum, <i>olim</i> Decoctum pectorale	141
Pro potu.	
sarsaparillæ	142
A. ℞j—2.	
sarsaparillæ compositum	143
A. ℞j—2.	
ulmi	144
A. ℞j—2.	
E.	
Elaterium	51
E. gr. ii ad gr. vi.	
Electuarium e cassia	216
E. ℥j ad ℥ij	
e scammonio	217
A. gr. xv ad gr. xxx.	
e fenna, <i>olim</i> Electuarium lenitivum	217
A. ℥iss ad ℥ij.	
Emplastrum ammoniaci cum hydrargyro, <i>olim</i> Emplastrum ex ammoniaco cum mercurio	224
cantharidis, <i>vice</i> Emplastri vesicatorii	225
ceræ, <i>olim</i> Emplastrum attrahens	226
cumini, <i>olim</i> Emplastrum e cymino	226
ladani, <i>olim</i> Emplastrum stomachicum	227
lithargyri, <i>olim</i> Emplastrum commune	228
cum gummi, <i>olim</i> Emplastrum commune cum gummi	228
Emplastrum	

INDEX MEDICAMENTORUM. 265

Emplastrum lithargyri cum hydrargyro, <i>olim</i> Empla-		
strum commune cum mercurio	229	
cum refina, <i>olim</i> Emplastrum		
adhæfivum	230	
picis burgundicæ, <i>olim</i> Emplastrum ce-		
phalicum	230	
saponis, <i>olim</i> Emplastrum e sapone	231	
thuris, <i>olim</i> Emplastrum roborans	231	
Extractum cacuminis geniftæ	—	49
A. ʒ fs ad ʒ j—3.		
cafcarrillæ	—	55
A. gr. xx ad gr. xxx—3.		
chamæmeli	—	49
A. ʒ j ad ʒ ij—2 ad 4.		
colocynthidis compositum, <i>vice</i> Extracti		
cathartici		50
E. gr. xx ad gr. xxx.		
corticis peruviani	—	53
A. gr. xx—4.		
cum refina	—	54
A. gr. xx—4.		
gentianæ	—	49
A. gr. xx ad gr. xl—4.		
glycyrrhizæ	—	49
hellebori nigri	—	49
E. gr. x ad gr. xx.		
jalapii	—	55
E. gr. vi ad gr. x.		
ligni campechenfis	—	52
A. gr. xxx ad gr. xl—4.		
		Extractum

Extractum rutæ	—	—	49
A. gr. x ad gr. xx—4.			
fabinæ	—	—	49
A. gr. xxx ad gr. xl—4.			
fennæ	—	—	55
E. gr. xl ad lx.			

F.

Ferri rubigo, <i>olim</i> Chalybis rubigo præparata			104
A. gr. x ad gr. xx—3.			
Ferrum ammoniacale, <i>olim</i> Flores martiales			103
A. gr. v ad gr. x—3.			
tartarifatum	—	—	105
A. gr. xxx—3.			
vitriolatum, <i>olim</i> Sal martis		—	105
A. gr. v—3.			
Flores benzoës, <i>olim</i> Flores benzöini		—	72
A. gr. xx—4.			
fulphuris loti	—	—	93
E. 3j.			
Florum exsiccatio	—	—	40

G.

Galbani purificatio	—	—	39
E. gr. xx ad xxx.			

H.

Herbarum exsiccatio	—	—	40
Hydrargyrus purificatus, <i>olim</i> Argenti vivi purificatio			107
A. ʒj ad ʒiv.			

Hydrargyrus

INDEX MEDICAMENTORUM. 267

Hydrargyrus acetatus	— — —	108
A. gr. ii ad gr. iv—1 ad 3.		
calcinatus, <i>olim</i> Mercurius calcinatus		109
A. gr. fs ad gr. i—1 ad 3.		
cum creta, <i>vulgo</i> Mercurius alkalifatus		109
A. gr. xxx—3.		
muriatus, <i>olim</i> Mercurius corrosivus sublimatus		110
A. gr. fs—2 ad 4.		
mitis, Mercurius præcipitatus dulcis, Pharm. Lond. 1721		111
A. gr. i ad gr. ii—1 ad 2.		
nitratus ruber, <i>olim</i> Mercurius corrosivus ruber		113
cum sulphure, <i>olim</i> Æthiops mineralis		114
A. gr. xxx—3.		
sulphuratus ruber, <i>olim</i> Cinnabaris factitia		115
vitriolatus, <i>olim</i> Mercurius emeticus flavus		116
A. gr. fs ad gr. i—1 ad 3. E. gr. v ad gr. viii.		

I.

Infusum gentianæ compositum, <i>olim</i> Infusum amarum simplex		146
A. ℥ ij—3.		
rosæ, <i>olim</i> Tinctura rosarum	— — —	149
A. ℥ ij—4 ad 6.		
fennæ simplex	— — —	147
E. ℥ iv.		
tartarifatum, <i>olim</i> Infusum fennæ commune		148
E. ℥ iv.		

K.

K.

Kali præparatum, <i>vice</i> Salis abinthii, &c.	73
A. gr. v ad gr. xv—3.	
acetatum, <i>olim</i> Sal diureticus	86
A. ʒj ad ʒij—3.	
purum, <i>vulgo</i> Alkali vegetabile fixum causticum	75
fulphuratum, <i>vulgo</i> Hepar sulphuris	93
tartarifatum, <i>olim</i> Tartarum solubile	87
E. ʒij ad ʒvj.	
vitriolatum, <i>olim</i> Tartarum vitriolatum, &c.	84
A. gr. xxx ad gr. lx—4.	

L.

Lac ammoniaci	180
A. ʒss ad ʒj—4.	
amygdalæ, <i>vice</i> Emulsionis communis	180
A. ʒiv—4 ad 8.	
Lapidis calaminaris præparatio	38
Linimentum ammoniæ, <i>olim</i> Linimentum volatile	241
fortius	241
camphoræ	242
saponis, <i>olim</i> Linimentum saponaceum	242
Liquor volatilis cornucervi, <i>olim</i> Spiritus cornucervi	82
A. gr. xxx ad gr. lx—4. ʒiv and ʒvi—1.	

M.

Magnesia alba	91
E. ʒj ad ʒij.	
	Magnesia

INDEX MEDICAMENTORUM. 269

Magnesia usta	_____	_____	92
E. ʒ ss ad ʒ j cum fs.			
Mel rosæ, <i>olim</i> Mel rosaceum	_____	_____	192
scillæ	_____	_____	193
A. ʒ ss ad ʒ j cum fs—3. E. ʒ ss.			
Mellis despumatio	_____	_____	41
Millepedæ præparatio	_____	_____	41
A. ʒ ij ad ʒ iv—2.			
Mistura camphorata, <i>olim</i> Julepum e camphora			178
A. ʒ ij—4.			
cretacea, <i>olim</i> Julepum e creta	_____		179
A. ʒ ij—4.			
moschata, <i>olim</i> Julepum e moscho			179
A. ʒ ij—4.			
Mucilago amyli	_____	_____	144
arabici gummi	_____	_____	145
A. ʒ ij—4.			
feminis cydonii mali	_____	_____	145

N.

Natron præparatum, <i>vulgo</i> Sal sodæ	_____		77
A. gr. xx ad gr. xl—3.			
tartarifatum, <i>vulgo</i> Sal rupellensis	_____	_____	88
E. ʒ ss ad . j.			
vitriolatum, <i>olim</i> Sal catharticus glauberi			84
E ʒ j ad ʒ j cum fs.			
Nitrum purificatum	_____	_____	85
A. gr. xl ad gr. lx—4.			

O.

O.

Oleum amygdalæ	—	—	57
A. ℥ ij—4. E. ℥ j ad ℥ ij.			
animale	—	—	61
A. gr. x ad gr. xx—3.			
cornu cervi	—	—	82
petrolei	—	—	69
fuccini	—	—	71
rectificatum	—	—	62
A. gr. x and gr. xx—3.			
fulphuratum, <i>olim</i> Balsamum sulphuris simplex			94
A. gr. iv ad gr. vi—3.			
terebinthinæ	—	—	60
A. gr. x ad gr. xv—3.			
rectificatum	—	—	61
A. gr. viii ad gr. xii—3.			
vini	—	—	62
e fiminibus lini	—	—	57
A. ℥ ss—6 ad 8.			
ricini	—	—	57
E. ℥ ss ad ℥ j.			
sinapeos	—	—	57
essentiale anisi	—	—	58
A. gr. ii ad gr. iv.			
baccæ junepери	—	—	58
A. gr. iii ad gr. v.			
carui	—	—	58
A. gr. i. ad gr. iii.			

Oleum

INDEX MEDICAMENTORUM, 275

Oleum effentiale lavendulæ	————	58
A. gr. ii. ad gr. iv.		
menthæ piperitidis	————	58
A. gr. fs ad gr. i.		
fativæ	————	58
A. gr. i ad gr. iii.		
origani	————	58
A. gr. ii ad gr. iv.		
pulegii	————	58
A. gr. ii ad gr. iv.		
radicis fassafras	————	58
A. gr. i ad gr. iii.		
rorismarini	————	58
A. gr. ii ad gr. iv.		
Opium purificatum, <i>olim</i> Extractum thebaicum		56
A. gr. fs ad gr. i—1. gr. $\frac{1}{2}$ ad $\frac{1}{2}$ —3 ad 6. gr. i—4. I.		
		gr. $\frac{1}{25}$.
Oxymel æruginis, <i>vice</i> Mellis ægyptiaci	————	193
colchici	————	193
A. 3 fs ad 3 j cum fs—3.		
scillæ, <i>olim</i> Oxymel scilliticum	————	194
A. 3 fs ad 3 j cum fs—3. E. 3 fs.		
simplex	————	194
A. 3 j—3.		

P.

Petroleum sulphuratum, <i>olim</i> Balsamum sulphuris bar-		
badense	————	94
Pilulæ ex aloë	————	211
E. gr. x ad gr. xxx.		
		Pilulæ

Pilulæ ex aloë cum myrrha, <i>olim</i> Pilulæ rufi	211
E. gr. x ad gr. xv.	
e gummi, <i>olim</i> Pilulæ gummosæ	212
E. gr. xx.	
ex hydrargyro, <i>vice</i> Pilularum mercurialium	213
A. gr. x ad gr. xx—1 ad 3.	
ex opio	214
Vidi OPIUM.	
e scilla	214
A. gr. viii—3.	
Pulparum extractio	42
Pulvis aloëticus, <i>olim</i> Hieræ picra	196
E. gr. x ad gr. xv ad gr. xxx.	
cum ferro, <i>vice</i> Pilularum ephracti-	
carum	197
E. gr. xx ad gr. xxx.	
cum guaiaco, <i>vice</i> Pilularum aroma-	
ticarum	196
E. gr. xv ad gr. xx	
antimonialis	98
A. gr. iv ad vi—4.	
aromaticus, <i>olim</i> Species aromaticæ	197
A. gr. iv.	
afari compositus, <i>olim</i> Pulvis sternutatorius	198
e cerussa	198
e chelis cancerorum compositus	199
A. 3j. I. gr. x—4 ad 12.	
contrayervæ compositus	199
A. gr. xxx—4.	
e creta compositus, <i>vice</i> Pulveris e bolo com-	
positi	200
A. gr. xxx ad gr. xl.	
	Pulvis

Succus limonis spissatus	— — —	48
ribis nigri spissatus	— — —	48
Sulphur Antimonii præcipitatum	— — —	99
A. gr. v ad gr. viii—4.		
præcipitatum	— — —	94
A. ʒ j ad ʒ ij.		
Syrupus althææ	— — —	184
caryophylli rubri	— — —	184
corticis aurantii	— — —	185
croci	— — —	185
mori	— — —	186
papaveris albi, <i>olim</i> Syrupus e meconio erratici	— — —	186
A. ʒ ij ad ʒ vj.		
erratici	— — —	187
ribis nigri	— — —	186
rosæ	— — —	188
rubi idæi	— — —	186
spinæ cervinæ	— — —	188
E. ʒ vj ad ʒ xij.		
succu limonis	— — —	186
tolutanus, <i>vice</i> syrui balsamici	— — —	189
violæ	— — —	190
zingiberis *	— — —	190

* The Syrups, whose doses are not set down, are used to sweeten other medicines, and the dose is about a drachm,

T.

Tinctura cinnamomi composita, <i>olim</i> Tinctura aro-	matica	163
A. gr. xxx ad gr. xl.		
colombæ	—————	163
A. gr. xxx ad gr. xc—4.		
corticis aurantii	—————	164
A. 3j.		
peruviani	—————	164
A. 3j ad ʒij—4.		
composita	—————	165
A. 3j ad ʒij—4.		
ferri muriati, <i>vice</i> Tincturæ martis in spiritu	falis	166
A. gr. xx ad gr. xl—3.		
galbani	—————	167
A. 3j ad ʒij.		
gentianæ composita, <i>olim</i> Tinctura amara		167
A. 3j ad ʒij—3.		
guaiaci, <i>olim</i> Tinctura guaiacina volatilis		168
A. 3j—3.		
hellebori nigri, <i>olim</i> Tinctura melampodii		168
jalapii	—————	169
E. 3j ad ʒij.		
lavendulæ composita, <i>olim</i> Spiritus laven-	dulæ compositus	169
A. gr. xxx ad gr. xl.		
myrrhæ	—————	170
opii, <i>vice</i> Tinctura thebaïcæ	—————	170
A. gr. v ad gr. x—4. gr. xv ad gr. xxv—1. vide opium.		
camphorata, <i>vice</i> Elixir paregorici		171
A. 3j ad ʒij.		
	Tinctura	

INDEX MEDICAMENTORUM.		279
Tinctura rhabarbari	— —	172
E. ʒj ad ʒiv.		
composita	— —	173
E. ʒj ad ʒiv.		
fabinae composita, <i>olim</i> Elixir myrrhae compositum	— —	174
A. ʒj.		
scillae	— —	174
A. gr. xxx ad gr. lx—3.		
fennae	— —	175
E. ʒij ad ʒss ad ʒj—4.		
serpentariae	— —	176
A. ʒj ad ʒij—4.		
valerianae	— —	176
A. ʒj ad ʒij—4.		
volatilis	— —	177
E. gr. xxx ad lx—4.		
Trochisci ex amylo, <i>olim</i> Trochisci bechici albi		207
e creta, <i>olim</i> Tabellae cardialgicae		208
e glycyrrhiza, <i>olim</i> Trochisci bechici nigri		208
e magnesia	— —	209
e nitro	— —	209
e sulphure	— —	210
Tutiae praeparatio	— —	38

V.


Vinum aloës, <i>olim</i> Tinctura sacra	— —	151
E. ʒss ad ʒj.		
antimonii	— —	152
A. gr. xxx ad gr. lx. E. ʒij.		
T 4		Vinum

Vinum antimonii tartarifati	—————	152
A. gr. xxv ad gr. xl. E. ʒj ad ʒij.		
ferri, <i>olim</i> Vinum chalybeatum	———	154
A. gr. xxx and gr. lx—3.		
ipecacuanhæ	———	154
A. gr. xxx ad gr. lx—4. E. ʒij ad ʒiv.		
rhabarbari, <i>olim</i> Tinctura rhabarbari vinosa		155
A. ʒfs ad ʒj.		
Unguentum adipis suillæ, <i>olim</i> Unguentum simplex		232
calcis hydrargyri albæ, <i>olim</i> Unguentum		
e mercurio præcipitato		233
cantharidis, <i>vice</i> Unguentied vesicatoria		233
ceræ, <i>olim</i> Unguentum album		234
cerussæ acetatæ, <i>olim</i> Unguentum satur-		
ninum		234
elemi	———	235
hellebori albi	———	235
hydrargyri fortius, <i>vice</i> Unguenti cæru-		
lei fortioris		236
mitius, <i>vice</i> Unguenti ceru-		
lei mitioris		237
nitрати	———	238
picis	———	238
resinæ flavæ, <i>olim</i> Unguentum basilicum		
flavum		239
sambuci	———	239
spermatis ceti, <i>olim</i> Linimentum album		240
fulphuris	———	240
tutiæ	———	240

Z.

Z.

- Zincum calcinatum, *vulgo* Flores zinci — 120
A. gr. iii ad gr. v—2.
vitriolatum purificatum, *vice* Salis vitrioli 121
A. gr. ii ad gr. v—2. E. gr. x.



ENGLISH INDEX.

A.

A C E T A T E D	ceruffe	_____	Page 117
	kali	_____	86
	quickfilver	_____	108
Acetous acid	_____	_____	70
Alkohol	_____	_____	127
Almond milk	_____	_____	180
Aloëtic pills	_____	_____	211
	with myrrh	_____	211
	powder	_____	196
	with guaiacum	_____	196
	with iron	_____	197
	wine	_____	151
Alum poultice	_____	_____	248
Ammoniacal iron	_____	_____	103
Ammoniac plaifter with quickfilver	_____	_____	224
Animal oil	_____	_____	61
Antimonial powder	_____	_____	98
	wine	_____	152
Aromatic confection	_____	_____	219
	powder	_____	197
			B.

B.

Barley water	—————	—————	141
Burgundy pitch plaister	— — —	—————	230
Burning of hartshorn	—————	—————	40
sponge	—————	—————	43
Burnt alum	—————	—————	89
magnesia	—————	—————	92

C.

Calcined antimony	—————	—————	96
quicksilver	—————	—————	109
zinc	—————	—————	120
Calomel	—————	—————	111
Camphorated mixture	—————	—————	178
spirit	—————	—————	182
tincture of opium	—————	—————	171
Camphor liniment	—————	—————	242
Cerate of acetated litharge	—————	—————	244
calamine	—————	—————	243
the cantharis	—————	—————	243
soap	—————	—————	245
sperma ceti	—————	—————	246
yellow rofin	—————	—————	245
Chalk lozenges	—————	—————	208
mixture	—————	—————	179
Cinnamon water	—————	—————	123
Clarification of honey	—————	—————	41
			Common

ENGLISH INDEX.

285

Common mint water	— —	124
Compound allum water	—	221
decoction of barley	—	141
sarsaparilla	—	143
extract of colocynthida	—	50
infusion of gentian	—	146
juice of scurvy gras	—	47
powder of asarum	—	198
chalk	—	200
chalk with opium	—	200
contrayerva	—	199
crabs claws	—	199
ipecacuan	—	201
myrrh	—	201
scammony	—	203
fenna	—	205
tragacanth	—	205
spirit of ammonia	—	181
anise	—	132
horse-radish	—	137
juniper	—	134
vitriolic æther	—	181
tincture of aloës	—	157
benzoës	—	158
cardamoms	—	160
cinnamon	—	163
gentian	—	167
lavender	—	169
peruvian bark	—	165
rhubarb	—	173
savin	—	174

Compound

Compound water of acetated litharge	—	—	222
Conserve of arum	—	—	45
the exterior rind of the Seville orange			44
the hip	—	—	45
lujula	—	—	44
the red rose	—	—	44
sea wormwood	—	—	44
the floe	—	—	45
the squill	—	—	46
Crocus of antimony	—	—	97
Cumin plaister	—	—	226
poultice	—	—	247

D.

Decoction of elm	—	—	144
for an enema	—	—	139
for fomentation	—	—	140
of hartshorn	—	—	138
of hellebore	—	—	140
of peruvian bark	—	—	139
of sarsaparilla	—	—	142
Dill water	—	—	123
Dilute nitrous acid	—	—	66
vitriolic acid	—	—	65
Distilled vinegar	—	—	68
water	—	—	122
Drying of herbs and flowers	—	—	40
of squills	—	—	42

E.

Elaterium	—	—	51
Electuary of cassia	—	—	216
scammony	—	—	217
fenna	—	—	217
Essential oil of aniseed	—	—	58
carraway	—	—	58
common mint	—	—	58
the juniper berry	—	—	58
lavender	—	—	58
origanum	—	—	58
pennyroyal	—	—	58
pepper mint	—	—	58
rosemary	—	—	58
sassafras root	—	—	58
Extract of black hellebore	—	—	49
broom tops	—	—	49
camomile	—	—	49
cascaquilla	—	—	55
gentian	—	—	49
jalap	—	—	55
liquorice	—	—	49
logwood	—	—	52
peruvian bark	—	—	53
rue	—	—	49
savin	—	—	49
fenna	—	—	55
pulps	—	—	42

F.

F.

Fennel water	—	—	124
Flowers of benzoës	—	—	72
Fœtid spirit of ammonia	—	—	132
Frankincense plaifter	—	—	231

G.

Gum pills	—	—	212
-----------	---	---	-----

H.

Honey of the rose	—	—	192
squill	—	—	193

I.

Jamaica pepper water	—	—	125
Infusion of the rose	—	—	149
Insuffated juice of black currants		—	48
the elder berry		—	47
hemlock	—	—	48
lemons	—	—	48
Ipecacuan wine	—	—	154

L.

L.

Ladanum plaister	_____	227
Lime mixed with pure kali	_____	76
water	_____	148
Liniment of ammonia	_____	241
soap	_____	242
Litharge plaister with rosin	_____	230
quickfilver	_____	229
Liquorice lozenges	_____	208
Lozenges of magnesia	_____	209
nitre	_____	209
fulphur	_____	210

M.

Mild muriated quickfilver	_____	111
Milk of ammoniac	_____	180
Mucilage of gum arabic	_____	145
quince feed	_____	145
starch	_____	144
Muriatic acid	_____	66
Muriated antimony	_____	97
quickfilver	_____	110
tincture of iron	_____	166
Musk mixture	_____	179
Mustard poultice	_____	247

U

N.

N.

Nitrated silver	—	—	102
Nitrous acid	—	—	65

O.

Oil of the almond	—	—	57
amber	—	—	71
hartshorn	—	—	82
mustard seed	—	—	57
petroleum	—	—	60
ricinus	—	—	57
turpentine	—	—	60
wine	—	—	62
Ointment of acetated cerusse	—	—	234
cantharides	—	—	233
elder	—	—	239
elemi	—	—	235
hogs lard	—	—	232
nitrated quicksilver	—	—	238
sulphur	—	—	240
tutty	—	—	240
wax	—	—	234
white calx of quicksilver	—	—	233
hellebore	—	—	235
yellow rosin	—	—	239
Opiate confection	—	—	220

Opiate

ENGLISH INDEX.

297

Opiate powder	—	—	202
Opium pills	—	—	214
Oxymel of colchicum		—	193
the squill	—	—	194
verdigris	—	—	193

P.

Pennyroyal water	—	—	125
Peppermint water	—	—	124
Plaiſter of the cantharis		—	225
litharge	—	—	228
with gum		—	228
Powder of ſcammony with aloës		—	203
calomel		—	204
white lead		—	198
Powdered tin	—	—	119
Precipitated ſulphur	—	—	94
of antimony		—	99
Preparation of amber	—	—	38
antimony		—	38
calamine		—	38
chalk	—	—	38
coral		—	38
crabs claws		—	38
hogs lard	—	—	39
millepedes		—	41
mutton ſuet		—	39
oyster ſhells		—	38
tutty	—	—	38
verdigris	—	—	38

U 2

Prepared

Prepared ammonia	—	—	78
kali	—	—	73
natron	—	—	77
Pure kali	—	—	75
Purification of ammoniac		—	39
azæ foetida		—	39
galbanum		—	39
styrax	—	—	43
Purified alum	—	—	89
nitre	—	—	85
opium	—	—	56
quickfilver	—	—	107
salt of amber	—	—	71

Q.

Quickfilver with chalk	—	—	109
fulphur	—	—	114
pills	—	—	213

R.

Rectified oil of amber	—	—	62
turpentine	—	—	61
Red nitrated quickfilver	—	—	113
fulphurated quickfilver	—	—	115
Resinous extract of bark		—	54
Rhubarb wine		—	155
Rose water	—	—	126
Rust of iron	—	—	104
			S.


		S.	
78			
73			
77			
75	Salt of amber	— —	71
39	hartshorn	— —	82
39	Simple infusion of fenna	— —	147
39	oxymel	— —	194
39	Soap plaister	— —	231
43	Spermaceti ointment	— —	240
89	Spirit of ammonia	— —	131
85	with amber	— —	182
56	carraway	— —	133
107	cinnamon	— —	133
71	garden mint	— —	135
	lavender	— —	134
	nitrous æther	— —	130
	nutmeg	— —	136
109	pennyroyal	— —	136
114	peppermint	— —	135
213	pimento	— —	136
	rosemary	— —	137
	vitriolic æther	— —	128
	Squill pills	— —	214
62	Starch lozenges	— —	207
61	Strong liniment of ammonia	— —	241
	quicksilver ointment	— —	236
113	Sulphurated kali	— —	93
115	oil	— —	94
54	petroleum	— —	94
155	Syrup of buckthorn	— —	188
126	clove July flowers	— —	184
104			Syrup
S.			


Syrup of ginger	—————	199
lemon juice	—————	186
marsh mallows	—————	184
orange peel	—————	185
the rose	—————	188
saffron	—————	185
tolu	—————	189
violets	—————	190
the white poppy	—————	186
wild poppies	—————	187

T.

Tar ointment	—————	238
Tartarised antimony	—————	100
infusion of fenna	—————	148
iron	—————	105
kali	—————	87
natron	—————	88
Tincture of aloës	—————	156
afa foetida	—————	157
balsam of peru	—————	157
tolu	—————	158
black hellebore	—————	168
cantharides	—————	159
cardamoms	—————	159
cascarilla	—————	161
castor	—————	161
catechu	—————	162
cinnamon	—————	162

Tincture





17. Jan 1871

18. Jan 1871

19. Jan 1871

20. Jan 1871

