

MATERIA MEDICA TABLE

See Notes

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Acaciæ Cortex.	{Acacia Arabica and Acacia decurrens . . .}	{Leguminosæ (Fabaceæ) . . .}	Africa, Australia and India .
Acaciæ Gummi . . .	{Acacia Senegal, and other species . . .}	" . . .	Khordofan, in Eastern Africa
Acalypha . . .	Acalypha Indica . . .	Euphorbiaceæ . . .	India . . .
Aconiti Folia . . .	Aconitum 'Napellus' . . .	Ranunculaceæ . . .	Britain . . .
Aconiti Radix . . .	" . . .	" . . .	" . . .
Aconitina . . .	" . . .	" . . .	" . . .
Adeps Lanæ . . .	Ovis Aries . . .	Ungulata . . .	Domesticated everywhere .
Adeps . . .	Sus Scrofa . . .	" . . .	" . . .
Adhatoda . . .	{Adhatoda Vasica (Justicia Adhatoda) . . .}	Acanthaceæ . . .	India . . .
Agropyrum . . .	Agropyrum repens . . .	Graminaceæ . . .	{Europe, North America, and Asia . . .}
Ajowan Oleum . . .	Carum Copticum . . .	{Umbelliferae (Apiaceæ) . . .}	India, Egypt and Persia . . .
Aloe Barbadensis . . .	{Aloe Chinensis, A. vera, and probably other species . . .}	Liliaceæ . . .	{Dutch West Indian Islands and Barbados . . .}
Aloe Socotrina . . .	{Aloe Perryi and probably other species . . .}	" . . .	{Socotra (shipped by way of Bombay and Zanzibar) . . .}
Alstonia . . .	{Alstonia scholaris . . . Alstonia constricta . . .}	Apocynaceæ . . .	India, the Philippine Islands, Australia . . .
Ammoniacum . . .	{Dorema Ammoniacum and other species . . .}	Umbelliferae . . .	Persia . . .
Amygdala Amara . . .	{Prunus Amygdalus (var. amara) . . .}	Rosaceæ . . .	{South of France, Sicily, and Morocco . . .}
Amygdala Dulcis . . .	{Prunus Amygdalus (var. dulcis) . . .}	" . . .	{Spain, Portugal, and South of France . . .}
Amygdalæ Oleum . . .	Both of the above . . .	" . . .	" . . .
Amylum . . .	{Triticum sativum (wheat) . . . Zea 'Mays' (maize) . . . Oryza sativa (rice) . . .}	Graminaceæ . . .	{Cultivated in various parts of the world . . .}
Andrographis . . .	{Andrographis paniculata . . .}	Acanthaceæ . . .	India, Ceylon, Java . . .
Anethi Fructus . . .	Peucedanum graveolens . . .	Umbelliferae . . .	{England, Middle and Southern Europe . . .}
Anisi Fructus . . .	Pimpinella Anisum . . .	" . . .	{Central and Southern Europe, etc. . .}
Anisi Oleum . . .	{Illicium anisatum . . .}	{Magnoliaceæ . . .}	{Distilled in Europe and in China . . .}
Anthemidis Flores . . .	Anthemis nobilis . . .	{Compositæ (Asteraceæ) . . .}	{Britain and Belgium, cultivated . . .}
Anthemidis Oleum . . .	" . . .	" . . .	{Britain and Belgium, and France, etc. . .}
Apomorphinæ Hydrochloridum . . .	Morphina or Codeina . . .	see Opium . . .	see Opium . . .
Arachis Oleum . . .	Arachis hypogæa . . .	Leguminosæ . . .	{Tropical Africa, China, India, etc., cultivated . . .}

OF THE ORGANIC KINGDOM.

on p. xxxvi.

Parts used.	Preparations into which it enters.
Dried bark . . . . .	Decoctum Acaciæ Corticis.
{Gum exuded from stem and branches . . . . .}	{Mucilago Acaciæ; Pulv. Amygdalæ Co. and Pulv. Tragacanth Co. All Trochisci.
Fresh and dried herb . . . . .	Extractum Acalyphæ Liquidum, and Succus Acalyphæ.
Fresh leaves and flowering tops . . . . .	
Dried root . . . . .	Linimentum, and Tinctura Aconiti, and Aconitina.
Dried root . . . . .	Unguentum Aconitinæ.
{The purified cholesterin-fat of sheep's wool . . . . .}	Adeps Lanæ Hydrosus.
Purified fat of the abdomen . . . . .	{Adeps Benzoatus; Empl. Canthar.; Pilula Phosphori; Unguenta Aconitinæ, Atropinæ, Cocainæ, Hydrargyri, Hydrarg. Nitratis, Iodi, Resinæ, and Veratrinæ.
Fresh and dried leaves . . . . .	Extractum Adhatodæ Liquidum, Succus Adhatodæ, and Tinctura Adhatodæ.
Dried rhizome . . . . .	Decoctum Agropyri, and Extractum Agropyri Liquidum.
Distilled from fruit.	
{Juice of the leaf evaporated to dryness . . . . .}	{Aloin; Decoct. Aloes Co.; Ext. Aloes Barb.; Ext. Coloc. Co.; Pil. Aloes Barb.; Pil. Aloes et Ferri; Pil. Cambogiæ Comp.; Pil. Colocynthidis Co.; Pil. Colocynthidis et Hyoseyami; Tinctura Aloes.
{Juice of the leaf evaporated to dryness . . . . .}	{Aloin; Ext. Aloes Socotrinæ; Pil. Aloes et Asafetidæ; Pil. Aloes et Myrrhæ; Pil. Aloes Socot.; Pil. Rhei Co.; Tinct. Benzoini Co.
Dried bark . . . . .	Infusum Alstoniæ, and Tinctura Alstoniæ.
{Gum-resinous exudation from the flowering and fruiting stem . . . . .}	{Emplastrum Ammoniaci c. Hydrargyro; Emplast. Galbani; Mistura Ammoniaci; Pilula Scillæ Co.; Pil. Ipecac. c. Scillâ.
Ripe seed . . . . .	Oleum Amygdalæ.
Ripe seed . . . . .	{Oleum Amygdalæ; Pulvis Amygdalæ Compositus; Mistura Amygdalæ.
Expressed oil from seeds . . . . .	Ol. Phosphoratum; Unguenta Aquæ Rosæ, and Cetacei.
Starch . . . . .	Glycerinum Amyli; Pulvis Tragacanth. Co.
Dried plant . . . . .	{Infusum Andrographidis, Liquor Andrographidis Concentratus, and Tinctura Andrographidis.
Dried ripe fruit . . . . .	Aqua Anethi; Oleum Anethi.
Dried ripe fruit . . . . .	Aqua Anisi; Oleum Anisi.
Oil distilled from fruits . . . . .	Spiritus Anisi; Tinct. Camph. Co.; Tinct. Opii Ammoniata.
{Dried expanded flower-heads or capitula . . . . .}	{Extractum and Oleum Anthemidis. The latter is also used in the preparation of the Extract.
Volatile oil . . . . .	Extractum Anthemidis.
Salt of the alkaloid . . . . .	Injectio Apomorphinæ Hypodermica.
Expressed oil, without heat.	

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Araroba . . . .	Andira 'Araroba' . . . .	Leguminosæ . . . .	Brazil (Bahia) . . . .
Areca . . . . .	Areca Catechu . . . . .	Palmaceæ . . . . .	{Malayan Archipelago, India, Ceylon and the Philippines}
Aristolochia . . . .	Aristolochia Indica . . . .	Aristolochiaceæ . . . .	India and Holland . . . .
Armoraciæ Radix.	Cochlearia Armoracia . . . .	{Cruciferae (Brassicaceæ)} . . . .	Britain, cultivated . . . .
Arnicae Flores . . . .	Arnica montana . . . . .	Compositæ . . . . .	{Mountainous parts of Mid- dle and Southern Europe.}
Arnicae Rhizoma . . . .	" " . . . . .	" " . . . . .	" " " " . . . . .
Asafetida . . . . .	{Ferula foetida, and probably other species}	Umbelliferae . . . . .	Afghanistan and Persia . . . .
Atropina . . . . .	{Belladonnae Radix or Folia . . . . .}	Solanaceæ . . . . .	{Britain, Germany, Austria, etc. . . . .}
Aurantii Floris Aqua . . . . .	{Citrus Aurantium var. 'Bigaradia' . . . .}	Rutaceæ . . . . .	South of Europe . . . . .
Aurantii Cortex . . . .	" " . . . . .	" " . . . . .	" " " " . . . . .
Aurantii Cortex Indicus . . . . .	Citrus Aurantium . . . . .	" " . . . . .	India and Ceylon . . . . .
Azadirachta Indica	Melia Azadirachta . . . . .	Meliaceæ . . . . .	Southern India, Ceylon . . . .
Balsamum Peruvianum	Myroxylon Pereiræ . . . . .	Leguminosæ . . . . .	{San Salvador, in Central America . . . . .}
Balsamum Tolutanum	Myroxylon Toluifera . . . . .	" " . . . . .	New Granada . . . . .
Bebeeru Cortex	Nectandra Rodiaei . . . . .	Lauraceæ . . . . .	Guiana . . . . .
Bela Fructus . . . . .	Ægle Marmelos . . . . .	Rutaceæ . . . . .	India . . . . .
Belladonnae Folia . . . .	Atropa Belladonna . . . . .	Solanaceæ . . . . .	Britain . . . . .
Belladonnae Radix	" " . . . . .	" " . . . . .	Britain, Germany, etc. . . . .
Benzoinum . . . . .	{Styrax Benzoin and other species . . . . .}	Styraceæ . . . . .	Siam and Sumatra . . . . .
Berberidis Cortex.	Berberis vulgaris . . . . .	Berberidaceæ . . . . .	Britain . . . . .
Berberis . . . . .	Berberis aristata . . . . .	" " . . . . .	India and Ceylon . . . . .
Betel . . . . .	Piper Betle . . . . .	Piperaceæ . . . . .	India, Ceylon, etc. . . . .
Bryoniae Radix . . . . .	{Bryonia dioica . . . . . Bryonia alba . . . . .}	Cucurbitaceæ . . . . .	{England and Central and Southern Europe . . . . .}
Buchu Folia . . . . .	Barosma betulina . . . . .	Rutaceæ . . . . .	Cape Colony . . . . .
Butea Gummi . . . . .	Butea frondosa . . . . .	Leguminosæ . . . . .	India . . . . .
Butea Seminum	" " . . . . .	" " . . . . .	" " " " . . . . .
Cadinum Oleum . . . . .	{Juniperus Oxycedrus and other species . . . .}	{Coniferae (Pinaceæ)} . . . . .	Southern Europe . . . . .
Caffeina . . . . .	{Camellia Thea . . . . . Coffea Arabica . . . . .}	{Camelliaceæ . . . . . Rubiaceæ . . . . .}	{China, Japan, and Upper India . . . . . Cultivated throughout the Tropics (native of Abyssinia)

Parts used.	Preparations into which it enters.
{A substance found in cavities in the trunk of Andira Araroba, freed as much as possible from fragments of wood, dried and powdered . . . .}	Chrysarobin; of Chrysarobin, Ung. Chrysarobin.
Seeds.	
Dried stem and root . . . . .	Liquor Aristolochiæ Concentratus, and Tinctura Aristolochiæ.
Fresh root . . . . .	Spiritus Armoraciæ Compositus.
Dried flower-heads . . . . .	Tinctura Arnicae Florum.
Dried rhizome and roots . . . .	Tinctura Arnicae.
Gum-resin . . . . .	{Pil. Aloes et Asafetida; Pil. Galbani Co.; Spiritus Ammonia Fetida; Tinctura Asafetida.
Alkaloid . . . . .	{Unguentum Atropinae, Atropinae Sulphas, Lamellæ Atropinae and Liquor Atropinae Sulphatis.
Distilled water of flowers . . . .	{Syrupus Aurantii Floris, Mistura Olei Ricini, Syrupus Calcii Lactophosphatis.
Dried outer part of the rind . . . .	{Fresh Peel: Tinctura Aurantii, Vinum Aurant., Syrupus Aurantii, Tinct. Quininae, Syrupus Aromaticus, and Syrup. Cascara Aromat. Dried Peel: Inf. Aurantii, Inf. Aurantii Comp.; Inf. Gentian. Comp.; Tinct. Cinchon. Comp.; Tinct. Gentian. Comp.; Spirit. Armoraciæ Comp.
{The fresh and dried outer part of the rind.	
Dried bark of the stem . . . . .	Infusum Azadirachtæ Indicae, and Tinctura Azadirachtæ.
Balsam, from the trunk.	
Balsam, from the trunk . . . . .	{Mist. Ammoniaci; Syrupus Tolutanus; Tinct. Tolutana; Tinct. Benzoini Co.; Tolu Basis; Troch. Acidi Carbolici; Troch. Morphinae; Troch. Morphinae et Ipecacuanhæ.
Dried bark.	
Fresh half-ripe fruit . . . . .	Extractum Bela Liquidum.
Fresh leaves and branches . . . .	{Extract. Belladonnae Viride; Succus Belladonnae; Atropina. Atropina; Emplast. Belladonnae; Extract. Belladonnae Alco- holicum; Extractum Belladonnae Liquidum; Linimentum Belladonnae; Tinct. Belladonnae; Ung. Belladonnae; Suppos. Belladonnae.
Dried root . . . . .	{Acidum Benzoicum; Adeps Benzoatus (and Ointments con- taining it); Tinct. Benzoini Co.; Ung. Cetacei.
Balsamic resin . . . . .	{Extractum Berberidis Fluidum, Berberinae Phosphas. Liquor Berberidis Concentratus, and Tinctura Berberidis.
Dried bark . . . . .	
Dried stem . . . . .	
Leaves . . . . .	
Fresh and dried root . . . . .	Tinctura Bryoniae.
Dried leaves . . . . .	Infusum Buchu, Tinctura Buchu.
Inspissated juice from stem . . . .	
Seeds . . . . .	Pulveris Butea Seminum.
{An empyreumatic oily liquid ob- tained by destructive distilla- tion of the branches and wood}	
{Alkaloid from leaves of tea or seeds of coffee. . . . .}	Caffeina Citras; Caffeina Citras Effervescens.

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Cajuputi Oleum .	Melaleuca leucadendron	Myrtaceæ . .	{Imported from Batavia and Singapore . . . . .}
Calendulæ Flores.	Calendula officinalis . .	Compositæ . .	Levant and Southern Europe
Calotropis . . .	{Calotropis procera . . . Calotropis gigantea . . .}	Asclepiadaceæ .	India . . . . .
Calumbæ Radix .	Jateorhiza 'Calumba' .	Menispermaceæ	{Eastern Africa, between Ibo and the Zambesi . . . . .}
Cambogia . . .	Garcinia Hanburii . . .	Clusiaceæ . . .	Siam . . . . .
Cambogia Indica .	Garcinia Morella . . .	" . . . . .	India . . . . .
Camphora . . .	Cinnamomum <i>Camphora</i>	Lauraceæ . . .	{China (Formosa) and Japan (purified in Britain and elsewhere) . . . . .}
Canellæ Cortex .	Canella alba . . . . .	Canellaceæ . .	West Indies . . . . .
Cannabis Indica .	Cannabis sativa . . . . .	Urticaceæ . . .	India . . . . .
Cantharis . . .	Cantharis vesicatoria .	Coleoptera . . .	{Spain, France, Sicily, Hungary, and Southern Russia}
Caoutchouc . . .	Hevea Brasiliensis . . .	Euphorbiaceæ .	Brazil (Para) . . . . .
Capsici Fructus .	Capsicum minimum . . .	Solanaceæ . . .	Zanzibar, Sierra Leone, etc.
Carbo Ligni . . .	Wood . . . . .	Various . . . .	Britain . . . . .
Cardamomi Semina	Elettaria <i>Cardamomum</i>	Scitaminaceæ .	India and Ceylon . . . . .
Carrageen . . .	Chondrus crispus . . . .	{Algæ (Gigartinaceæ)}	{North Cape to Gibraltar . . . Eastern Coasts of N. America}
Carui Fructus . .	Carum Carvi . . . . .	Umbellifere . .	Europe . . . . .
Carui Oleum . . .	" " . . . . .	" . . . . .	Central and Northern Europe
Caryophylli Oleum	Eugenia caryophyllata .	Myrtaceæ . . .	{Molucca Islands, Zanzibar, and Pemba . . . . .}
Caryophyllum . .	" " . . . . .	" . . . . .	{Zanzibar, Penang, Bencoolen, etc. . . . .}
Cascara Sagrada .	Rhamnus Purshianus . . .	Rhamnaceæ . .	California . . . . .
Cascarilla . . .	Croton <i>Eluteria</i> . . . . .	Euphorbiaceæ .	Bahama Islands . . . . .
Cassie Pulpa . . .	Cassia fistula . . . . .	Leguminosæ . .	East and West Indies . . .
Castoreum . . .	Castor Fiber . . . . .	Rodentia . . . .	Siberia and Canada . . . .
Catechu . . . . .	Uncaria 'Gambier' . . . .	Rubiaceæ . . .	{Singapore, and other places in the Eastern Archipelago}
Catechu Nigrum (Pegu Catechu and Cutch) . . .	Acacia Catechu . . . . .	Leguminosæ . .	India and Burma . . . . .
Cera Alba . . . .	Apis mellifica . . . . .	Hymenoptera .	Britain . . . . .
Cera Flava . . . .	" " . . . . .	" . . . . .	Britain, etc. . . . .
Cetaceum . . . .	Physeter macrocephalus	Cetacea . . . .	Pacific and Indian Oceans .
Cetraria . . . . .	Cetraria Islandica . . . .	{Discomycetes or Disco-lichenes}	North of Europe . . . . .

Parts used.	Preparations into which it enters.
{Volatile oil distilled from the leaves . . . . .}	Spiritus Cajuputi, Lin. Crotonis.
Dried florets . . . . .	Tinctura Calendulæ Florum.
Dried root-bark . . . . .	Tinctura Calotropis.
{Dried transversely cut slices of the root . . . . .}	{Infusum, Liquor Calumbæ Concentratus, and Tinct. Calumbæ.
Gum-resin . . . . .	Pilula Cambogiæ Composita.
Gum-resin . . . . .	{Aqua Camph.; Linimenta Aconiti, Belladonnæ, Camph., Camph. Ammoniatum, Chloroformi, Hydrargyri, Opii, Saponis, Sinapis, Terebinthinæ, and Tereb. Aceticum; Spiritus Camph.; Tinct. Camph. Co.; Ung. Hydrarg. Co.
{A white crystalline substance obtained from the wood. . . . .}	{Aqua Camph.; Linimenta Aconiti, Belladonnæ, Camph., Camph. Ammoniatum, Chloroformi, Hydrargyri, Opii, Saponis, Sinapis, Terebinthinæ, and Tereb. Aceticum; Spiritus Camph.; Tinct. Camph. Co.; Ung. Hydrarg. Co.
Dried bark . . . . .	{Extractum Cannabis Ind.; Tinctura Cannab. Ind.; Tinct. Chlorof. et Morphine Comp.
{Dried flowering or fruiting tops of female plants . . . . .}	{Acetum, Emplastrum, Tinctura and Unguentum Cantharidis; Colloidium Vesicans; Liquor Epispast.; Emplast. Calefaciens.
The dried beetle . . . . .	Liquor Caoutchouc, Charta Sinapis.
Prepared milk juice . . . . .	{Tinctura Capsici; Tinct. Chlorof. et Morphine Comp.; Ung. Capsici.
Dried ripe fruit . . . . .	{Carbonaceous residue from wood.
{Carbonaceous residue from wood.	{Tinct. Cardam. Co.; Ext. Coloc. Co.; Pulv. Cinnam. Co.; Pulv. Cretæ Arom.; Tinct. Gentian. Co.; Tinct. Rhei Co.; Decoct. Aloes Comp.; Mist. Sennæ Comp.
The dried ripe seeds . . . . .	Saccharum, and Gelatina Carrageen.
Dried seaweed . . . . .	{Aqua and Oleum Carui; Conf. Piperis; Pulvis Opii Compositus; Tinct. Cardam. Co.; Tinci. Sennæ Co.; Pil Aloes Barb. Pilula Aloes Barbadosis.
Dried fruit . . . . .	{Pilula Colocynthidis, Composita, Pilula Colocynthidis et Hyoscyami.
Volatile oil . . . . .	Infusum and Oleum Caryophylli; Inf. Aurant. Co.; Pulv. Cretæ Aromat.; Pil. Coloc. Co.; Pil. Coloc. et Hyoscyami.
Volatile oil . . . . .	Ext. Cascariæ Sagradæ; Ext. Cascar. Sagrad. Liquid.; Syrup. Cascariæ Aromaticus.
Dried flower-buds . . . . .	Infusum and Tinctura Cascarilla.
Dried bark . . . . .	Confectio Sennæ.
Dried bark . . . . .	Tinctura Castorei.
Pulp from the pods . . . . .	{Infusum Catechu, Pulvis Catechu Comp., Tinctura Catechu, Trochiscus Catechu.
{Dried preputial follicles and secretion . . . . .}	{Pulvis Catechu Compositus, Tinctura Catechu, Trochiscus Catechu.
{An extract of the leaves and young shoots . . . . .}	{Pilula Phosphori, Suppositoria Acidi Carbolicum, Unguentum Aqua Rosæ, and Unguentum Cetacei.
Extract from the wood . . . . .	Emplast. Calefaciens, Cantharidis, Picis; Unguenta Hydrarg. Co., Menthol, Picis Liquid., Resinæ, and Staphisagriæ. Cera Alba:—Pilula Phosphori, Suppos. Acidi Carbolicum, Ung. Aquæ Rosæ, Ung. Cetacei.
Honeycomb; wax, bleached . . . . .	Unguentum Cetacei, Ung. Aquæ Rosæ and Ung. Capsici.
Honeycomb . . . . .	{A concrete fatty substance, mixed with oil, obtained from the head of the sperm whale, purified . . . . .}
Dried lichen . . . . .	Decoctum Cetrariæ, Saccharum Cetrariæ, Gelatina Cetrariæ.

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Chirata . . . . .	Swertia 'Chirata' . . .	Gentianaceæ .	Northern India . . . . .
Chrysarobinum . . . . .	See Araroba.		
Cimicifugæ Rhizoma . . . . .	Cimicifuga racemosa . . .	Ranunculaceæ .	Canada and U.S. . . . .
Cinchonæ Flav. Cortex . . . . .	Cinchona Calisaya . . . . .	Rubiaceæ . . . . .	{Bolivia, Southern Peru, and India . . . . .}
Cinchonæ lancifoliæ Cortex . . . . .	Cinchona lancifolia . . . . .	" . . . . .	New Granada . . . . .
Cinchonæ Rubræ Cortex . . . . .	Cinchona succirubra . . . . .	" . . . . .	{South America, cultivated in East Indies, Ceylon, Java, etc. . . . .}
Cinnamomi Cortex . . . . .	{Cinnamomum Zeylanicum . . . . .}	Lauraceæ . . . . .	Ceylon . . . . .
Cinnamomi Oleum . . . . .	" . . . . .	" . . . . .	" . . . . .
Cissampelos . . . . .	Cissampelos 'Pareira' . . . . .	Menispermaceæ . . . . .	{India, East and West Indies, and South Central America . . . . .}
Cocæ Folia . . . . .	Erythroxylon 'Coca' . . . . .	Linaceæ . . . . .	Peru, Java, and Bolivia . . . . .
Cocainæ Hydrochloridum . . . . .	" . . . . .	" . . . . .	" . . . . .
Coccus . . . . .	Coccus Cacti . . . . .	Hemiptera . . . . .	Mexico and Teneriffe . . . . .
Colchici Cormus . . . . .	Colchicum autumnale . . . . .	Liliaceæ . . . . .	Indigenous . . . . .
Colocynthidis Pulpa . . . . .	Citrullus <i>Colocynthis</i> . . . . .	Cucurbitaceæ . . . . .	{Northern Africa, Syria, and Spain . . . . .}
Condurango . . . . .	Gonolobus 'Condurango' . . . . .	Asclepiadaceæ . . . . .	Ecuador . . . . .
Conii Folia . . . . .	Conium maculatum . . . . .	Umbellifere . . . . .	Britain . . . . .
Conii Fructus . . . . .	" . . . . .	" . . . . .	" . . . . .
Copaiba . . . . .	{Copaifera Lansdorffii and other species . . . . .}	Leguminosæ . . . . .	Valley of the Amazon, etc. . . . .
Copaibæ Oleum . . . . .	" . . . . .	" . . . . .	" . . . . .
Coriandri Fructus . . . . .	Coriandrum sativum . . . . .	Umbellifere . . . . .	Germany and Britain, etc. . . . .
Coriandri Oleum . . . . .	" . . . . .	" . . . . .	" . . . . .
Coscinium . . . . .	Coscinium fenestratum . . . . .	Menispermaceæ . . . . .	Ceylon and Southern India . . . . .
Coto . . . . .	{Botanical source uncertain . . . . .}	Lauraceæ ? . . . . .	New Granada . . . . .
Crocus . . . . .	Crocus sativus . . . . .	Iridaceæ . . . . .	Spain and France, etc. . . . .
Crotonis Oleum . . . . .	Croton <i>Tiglium</i> . . . . .	Euphorbiaceæ . . . . .	{Hindustan, Ceylon, and Indian Archipelago . . . . .}
Cubebæ Fructus . . . . .	Piper <i>Cubeba</i> . . . . .	Piperaceæ . . . . .	Java . . . . .
Cucurbitæ Semina Preparata . . . . .	{Cucurbita maxima (Cucurbita Pepo) . . . . .}	Cucurbitaceæ . . . . .	{Italy, West Indies, Burma, and South America, cultivated . . . . .}
Curara . . . . .	Strychnos toxifera, etc. . . . .	Loganiaceæ . . . . .	South America . . . . .
Cuspariæ Cortex . . . . .	Cusparia febrifuga . . . . .	Rutaceæ . . . . .	Tropical South America . . . . .
Cusso . . . . .	Brayera anthelmintica . . . . .	Rosaceæ . . . . .	Abyssinia . . . . .
Cydonium . . . . .	Pyrus <i>Cydonia</i> . . . . .	" . . . . .	Western Asia and Europe . . . . .
Damiana . . . . .	{Turnera aphrodisiaca, etc. . . . .}	Turneraceæ . . . . .	Mexico and California . . . . .

Parts used.	Preparations into which it enters.
Dried plant . . . . .	Infusum, Liquor Chiratæ Concentratus, and Tinctura Chiratæ.
Dried rhizome and roots . . . . .	Ext. Cimicifugæ Liquid.; Tinct. Cimicifugæ.
Dried bark . . . . .	Used in the preparation of the official Salts of Quinine.
Dried bark . . . . .	{Its use is permitted for the manufacture of the official Salts of Quinine.
{Dried bark of stem and branches of cultivated plants}	{Ext. Cinchon. Liquid.; Inf. Cinchon. Acid.; Tinct. Cinchonæ; Tinct. Cinchon. Comp.; Quinine.
{The dried inner bark of shoots from the truncated stocks . . . . .}	{Aqua, Oleum, Pulvis Co., and Tinct. Cinnamomi; Pulvis Catechu Comp. Tinct. Catechu; Decoct. Hæmatoxyli; Pulv. Cretæ Arom.; Pulv. Kino Co.; Tinct. Cardam. Co.; Tinct. Lavand. Co.; of the Water, Mist. Cretæ, Mist. Guaiaci, Mist. Olei Ricini, Mist. Spiritus Vin. Gallici, Syrupus Aromaticus and Syrupus Cascaræ Aromaticus. Spiritus Cinnamomi; of the Compound Powder, Pil. Aloes et Ferri, and Pil. Cambogiæ Composita; of the Spirit, Acidum Sulphuricum Aromaticum.
Volatile oil from bark . . . . .	
Dried root . . . . .	Decoctum Cissampeli, and Extractum Cissampeli Liquidum.
Dried leaves . . . . .	Ext. Cocæ Liquid.; Cocainæ Hydrochloridum; Cocaine.
Salt of the alkaloid . . . . .	Lamellæ Cocainæ and Injectio Cocainæ Hypodermica.
Dried fecundated female insect	Tinctura Cocci; Tinct. Cardam. Co.; Tinct. Cinchonæ Co.
Fresh corn dried	Extractum and Vinum Colchici.
Dried ripe seeds	Tinctura Colchici Seminum.
{Dried pulp of the fruit freed from seeds . . . . .}	{Extractum Coloc. Co.; Pil. Coloc. Co.; Pil. Coloc. et Hyoscyam.
Dried bark . . . . .	
{Fresh leaves and young branches . . . . .}	Succus Conii; Unguentum Conii (from Succus).
Dried full-grown unripe fruits	Tinctura Conii.
Oleo-resin from the trunk . . . . .	Oleum Copaibæ.
Distilled oil from oleo-resin.	
Dried ripe fruit . . . . .	{Oleum Coriandri; Conf. Sennæ; Syrupus Rhei et Tinct. Rhei Co.; Syrupus Sennæ et Tinct. Sennæ Co.
Volatile oil . . . . .	
Dried stem . . . . .	{Infusum Coscinii, Liquor Coscinii Concentratus, and Tinctura Coscinii.
Bark . . . . .	
{The dried stigmas and tops of the styles . . . . .}	Tinctura Croci; Decoct. Aloes Co.; Tinct. Cinchon. Co.
Expressed oil from the seeds . . . . .	Linimentum Crotonis.
Dried full-grown unripe fruits.	Oleum Cubebæ; Tinctura Cubebæ.
Fresh ripe seeds.	
Extract from plant.	
Dried bark . . . . .	
{Dried panicles of pistillate flowers . . . . .}	Infusum Cuspariæ, and Liquor Cuspariæ Concentratus.
Seeds . . . . .	
Leaves . . . . .	

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Daturæ Folia . . .	{ <i>Datura fastuosa</i> var. <i>alba</i> <i>Datura Metel</i> . . . }	Solanaceæ . . .	India . . . . .
Daturæ Semina . . .	<i>Datura fastuosa</i> var. <i>alba</i>	" . . . . .	" . . . . .
Digitalis Folia . . .	<i>Digitalis purpurea</i> . . .	Scrophulariaceæ	Britain . . . . .
Elaterium . . . . .	<i>Ecballium Elaterium</i> . . .	Cucurbitaceæ . . .	Britain (cultivated), Malta . . .
Elemi . . . . .	{ <i>Canarium commune</i> and other species . . . }	Burseraceæ . . .	Manila, Brazil, and Mexico . . .
Embelia . . . . .	{ <i>Embelia ribes</i> , <i>Embelia</i> <i>robusta</i> . . . . . }	Myrsinaceæ . . .	India and East Indies . . .
Ergota . . . . .	<i>Secale cereale</i> . . . . .	Graminaceæ . . .	Europe and the Canaries . . .
Erythrophloeum . . .	{ <i>Erythrophloeum Guine-</i> <i>ense</i> . . . . . }	Leguminosæ . . .	Western Africa . . . . .
Eucalypti Oleum . . .	{ <i>Eucalyptus globulus</i> , and other species . . . }	Myrtaceæ . . . . .	Australia, etc. . . . .
Eucalypti Gummi . . .	{ <i>Eucalyptus rostrata</i> and other species . . . }	" . . . . .	" . . . . .
Euonymi Cortex . . .	<i>Euonymus atropur-</i> <i>pureus</i> . . . . .	Celastraceæ . . .	United States . . . . .
Euphorbia piluli- fera . . . . .	<i>Euphorbia pilulifera</i> . . .	Euphorbiaceæ . . .	{Queensland and Tropical America . . . . . }
Euphorbium . . . . .	<i>Euphorbia resinifera</i> . . .	" . . . . .	Morocco . . . . .
Fel Bovinum Puri- ficatum . . . . .	<i>Bos Taurus</i> . . . . .	Ungulata . . . . .	Domesticated everywhere . . .
Ficus . . . . .	<i>Ficus Carica</i> . . . . .	Urticaceæ . . . . .	Smyrna . . . . .
Filix-Mas . . . . .	<i>Aspidium Filix-mas</i> . . .	Filices . . . . .	Britain . . . . .
Fœniculi Fructus . . .	<i>Fœniculum capillaceum</i>	Umbelliferae . . .	{Central and Southern Europe, also India and Japan, etc. . . . . }
Fucus . . . . .	<i>Fucus vesiculosus</i> . . .	{Algae (Fucaceæ) . . . }	Britain . . . . .
Galbanum . . . . .	{ <i>Ferula galbaniflua</i> and probably other species }	Umbelliferae . . .	Persia . . . . .
Galla . . . . .	<i>Quercus infectoria</i> . . .	{Cupuliferae (Corylaceæ) . . . }	Asia Minor . . . . .
Gaultheriæ Oleum . . .	{ <i>Gaultheria procumbens</i> (leaves) . . . . . }	Ericaceæ . . . . .	United States and Canada . . .
Gelsemii Radix . . .	<i>Betula lenta</i> (bark) . . .	Betulaceæ . . . . .	" . . . . .
Gentianæ Radix . . .	<i>Gelsemium nitidum</i> . . .	Loganiaceæ . . .	{Southern part of the United States of America . . . }
Glycyrrhizæ Radix . . .	<i>Gentiana lutea</i> . . . . .	Gentianaceæ . . .	{Central and Southern Eu- rope (mountains) . . . }
Gossypii Radicis Cortex . . . . .	{ <i>Glycyrrhiza glabra</i> and other species . . . . . }	Leguminosæ . . .	{England, France, Spain, Sicily, Russia, and Persia }
Gossypium . . . . .	<i>Gossypium herbaceum</i> . . .	Malvaceæ . . . . .	{Imported from the United States . . . . . }
Graminis Citrati Oleum . . . . .	{ <i>Gossypium Barbadosense</i> and other species . . . }	" . . . . .	Warm and Tropical regions . . .
Granati Cortex . . .	<i>Andropogon citratus</i> . . .	Graminaceæ . . .	{India, Malay Peninsula, Ceylon, etc. (cultivated) . . . }
Grindelia . . . . .	<i>Punica Granatum</i> . . . . .	Lythraceæ . . . . .	{Shores of the Mediterranean and Central Asia . . . . . }
Guaiaci Lignum . . .	<i>Grindelia camporum</i> . . .	Compositæ . . . . .	{Southern part of the United States of America . . . . . }
Guaiaci Resina . . .	{ <i>Guaiacum officinale</i> or <i>Guaiacum sanctum</i> . . . }	Zygophyllaceæ . . .	St. Domingo and Jamaica . . .

Parts used.	Preparations into which it enters.
Dried leaves.	
Dried seeds . . . . .	Tinctura Daturæ Seminum.
Dried leaves . . . . .	Infusum and Tinctura Digitalis.
Nearly ripe fruit . . . . .	Elaterinum, Pulvis Elaterini Compositus.
Oleo-resinous exudation.	
Fruit.	
{The sclerotium of <i>Claviceps</i> <i>purpurea</i> , originating in the ovary of <i>Secale cereale</i> . . . }	{Extractum Ergotæ, Extractum Ergotæ Liquidum, Infusum Ergotæ, and Tinctura Ergotæ Ammoniata, Ergotinum, Inject. Ergotæ Hypoderm. . . }
Bark.	
{Distilled oil from the fresh leaves . . . . . }	Unguentum Eucalypti.
A ruby-coloured exudation . . .	Trochiscus Eucalypti Gummi.
Dried root bark . . . . .	Extractum Euonymi Siccum.
Dried herb.	
Concrete resinous juice.	
The purified ox bile.	
Dried fleshy receptacles . . . . .	Confectio Sennæ.
Dried rhizome . . . . .	Extractum Filicis Liquidum.
Dried ripe fruit . . . . .	Aqua Fœniculi; Pulv. Glycyrrhizæ Co.
Dried seaweed.	
Gum-resin . . . . .	Pil. Galbani Co
{Excrescences caused by the punctures and deposition of an egg or eggs of <i>Cynips</i> <i>Galle tinctoria</i> . . . . . }	{Acidum Gallicum and Tannicum; Tinct. Gallæ, Ung. Gallæ, and Ung. Gallæ c. Opio. . . }
Distilled oil.	
Dried rhizome and roots . . . . .	Tinct. Gelsemii.
Dried rhizome and roots . . . . .	Extractum, Infusum Co., and Tinct. Gentianæ Co.
{The peeled root and peeled subterranean stem . . . . . }	{Extract., Ext. Liquid., Liquor Sarsæ Comp. Conc., and Pulv. Gycyrrh. Co.; Pil. Hydr. Of the Extract, Conf. Sennæ, and Decoct. Aloes Comp. Of the Liquid Extract, Mistura Sennæ Comp., and Tinctura Aloes. . . }
Dried root-bark . . . . .	Decoctum Gossypii Radicis Corticis, and Extractum Radicis Corticis Liquidum.
Hairs of the seed . . . . .	Pyroxylin.
Distilled oil: 'Oil of Verbena.'	
Dried bark of the stem and root	Decoctum Granati Corticis.
Dried leaves and flowering tops	Extractum Grindeliæ Liquidum.
Heart-wood . . . . .	Liquor Sarsæ Compositus Concentratus.
Resin from the stem . . . . .	{Mist. Guaiaci; Pil. Hydrarg. Subchlor. Co., Tinct. Guaiaci Ammon.; Trochiscus Guaiaci Resinæ. . . }

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Gummi Indicum .	Anogeissus latifolia . .	Combretaceæ .	India and Ceylon . . . . .
Gynocardia Oleum	Taraktogenos Kurzii . .	Bixaceæ . . . .	{Malay Peninsula and Eastern India . . . . .}
Hæmatoxyli Lig- num . . . . .	{Hæmatoxylon Campe- chianum . . . . .}	Leguminosæ . .	{Campeachy, Honduras, and Jamaica . . . . .}
Hamamelidis Cor- tex . . . . .	Hamamelis Virginiana .	Hamamelidaceæ	United States and Canada .
Hamamelidis Folia	" " " "	" " " "	" " " "
Hemidesmi Radix	Hemidesmus Indicus . .	Asclepiadaceæ	India and Ceylon . . . . .
Hirudo . . . . .	Sanguisuga {medicinalis (speckled) officinalis (green)}	Hirudinea . . .	Spain, France, Italy, Hungary
Hirudo Australis .	{Hirudo quinquestriata (Limnobia quin- questriata)}	" " " "	Australia . . . . .
Hydrastis Rhizoma	Hydrastis Canadensis . .	Ranunculaceæ	United States and Canada .
Hygrophila . . .	Hygrophila spinosa . . .	Acanthaceæ . .	India . . . . .
Hyoscyami Folia .	Hyoscyamus niger . . . .	Solanaceæ . . .	Britain and Germany, etc. .
Ignatia amara . .	Strychnos Ignatii . . . .	Loganiaceæ . . .	Philippines . . . . .
Ipecacuanhæ Radix	Psychotria Ipecacuanha	Rubiaceæ . . . .	Brazil and Selangor . . . . .
Iris . . . . .	Iris versicolor . . . . .	Iridaceæ . . . .	United States . . . . .
Ispaghula . . . .	Plantago ovata . . . . .	Plantaginaceæ	India and Persia . . . . .
Jaborandi Folia .	Pilocarpus 'Jaborandi'	Rutaceæ . . . . .	Brazil (Pernambuco).
Jalapa . . . . .	Ipomœa 'Purga' . . . . .	Convolvulaceæ	{Mexico (cultivated in Ja- maica and India) . . . . .}
Juniperi Oleum . .	Juniperus communis . . . .	Coniferae . . . .	North of Europe, indigenous
Kaladana . . . . .	Ipomœa hederacea . . . .	Convolvulaceæ	India and Persia . . . . .
Kamala . . . . .	Mallotus Philippinensis	Euphorbiaceæ . .	{Ceylon, the Philippines, China, Australia, etc. . . .}
Kavæ Rhizoma . . .	Piper Methysticum . . . . .	Piperaceæ . . . .	Sandwich Islands . . . . .
Kino . . . . .	Pterocarpus marsupium	Leguminosæ . . .	Malabar . . . . .
Kino Eucalypti . .	{Eucalyptus calophylla, etc. . . . .}	Myrtaceæ . . . .	Western Australia . . . . .
Kola . . . . .	Cola acuminata . . . . .	Sterculiaceæ . .	West Coast of Africa . . . .
Krameria Radix . .	{Krameria triandra (Peruvian) . . . . . Krameria argentea (Pará) . . . . .}	Polygalaceæ . . .	Peru and Brazil . . . . .
Lactuca . . . . .	Lactuca virosa . . . . .	Compositæ . . . .	{Western, Central and South- ern Europe and Britain . . .}
Laricis Cortex . . .	Larix Europæa . . . . .	Coniferae . . . .	{England, Southern and Cen- tral Europe . . . . .}
Laurocerasi Folia .	Prunus Laurocerasus . . . .	Rosaceæ . . . . .	Britain, cultivated . . . . .
Lavandulæ Oleum	Lavandula vera . . . . .	Labiatae . . . . . (Lamiaceæ)	{England, cultivated (also the Western shores of the Mediterranean) . . . . .}

Parts used.	Preparations into which it enters.
{Gummy exudation from the wood . . . . .}	Mucilago Gummi Indici.
Fatty oil expressed from seeds	Unguentum Gynocardia.
Heart-wood . . . . .	Decoctum Hæmatoxyli.
Dried bark . . . . .	Tinctura Hamamelidis.
Leaves, fresh and dried . . .	{Of the Dried Leaves, Extractum Hamamelidis Liquidum; Unguentum Hamamelidis (from Liquid Extract). Of the Fresh Leaves, Liquor Hamamelidis.
Dried root . . . . .	Syrupus Hemidesmi.
Leech (annelid).	
Leech (annelid).	
Dried rhizome and roots . . .	Extractum Hydrastis Liquidum, Tinctura Hydrastis.
Dried herb including the root.	Decoctum Hygrophilæ.
{Fresh leaves and flowers, with branches . . . . .}	Extractum Viride, Succus, and Pil. Coloc. et Hyoscyam.
{Dried leaves and flowering tops . . . . .}	Tinctura Hyoscyami; Hyoscyin. Hydrobrom.; Hyoscyamine Sulphas.
Seeds (dried) . . . . .	{Acetum Ipecac.; Pil. Ipecac. c. Scillâ; Pulv. Ipecac. Co.; Trochiscus Ipecac.; Trochiscus Morphinae et Ipecac.; Vinum Ipecac.; Extractum Ipecacuanhæ Liquidum.
Dried root . . . . .	
Rhizome and roots . . . . .	
Dried seeds . . . . .	Decoctum Ispaghulæ.
Dried leaflets . . . . .	Extract. Liquidum, and Tinct. Jaborandi; Pilocarpinae Nitras. Extractum, Pulv. Co., Resina, and Tinctura Jalapæ; Pilula Scammonii Composita; Pulvis Scammonii Comp.
Dried tubercles . . . . .	
{Oil from the full-grown unripe green fruit. . . . .}	Spiritus Juniperi, and Misturi Creosoti.
Dried seeds . . . . .	{Pulvis Kaladanae Compositus, Resina Kaladanae, and Tinctura Kaladanae.
{Minute glands and hairs from the fruits. . . . .}	
Rhizome without the roots . .	Extractum Kavæ Liquidum.
{Juice from the trunk evapo- rated to dryness . . . . .}	Pulvis Compositus, Tinct. Kino; Pulv. Catechu Co.
Exudation from the stem.	
The seeds.	
Dried root . . . . .	{Extractum, Infusum, Tinctura Krameria; Pulv. Catechu Co.; Liquor Krameria Concentratus; Troch. Krameria; Troch. Krameria et Cocainæ.
Fresh herb.	
Bark (dried).	
Fresh leaves . . . . .	Aqua Laurocerasi.
Distilled oil from flowers . . .	{Spiritus and Tinctura Lavandulæ Comp.; Lin. Camph. Am- moniatum; Tinct. Lavand. Comp. is contained in Liquor Arsenicalis.

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Limonis Cortex	{Citrus medica var. 'Limonum' . . . .}	Rutaceæ . . .	South of Europe . . . .
Limonis Succus	" . . . . .	" . . . . .	" . . . . and West Indies
Linum . . . . .	Linum usitatissimum . . . .	Linaceæ . . .	Britain, Holland, Russia, etc.
Lobelia . . . . .	Lobelia inflata . . . . .	Campanulaceæ	North America
Lupulinum . . . . .	Humulus <i>Lupulus</i> . . . . .	Urticaceæ . . .	{England, imported from Europe, United States, and Canada . . . . .}
Lupulus . . . . .	" . . . . .	" . . . . .	" . . . . .
Lycopodium . . . . .	{Lycopodium "clavatum" and other species . . . .}	Lycopodiaceæ	{Great "Britain," Central and Northern Europe, etc. . . .}
Manna . . . . .	Fraxinus Ornus . . . . .	Oleaceæ . . .	Sicily and Calabria . . . .
Mastiche . . . . .	Pistacia Lentiscus . . . . .	Anacardiaceæ .	The Island of Scio . . . . .
Matico . . . . .	Piper angustifolium . . . . .	Piperaceæ . . .	Peru, etc. . . . .
Mel Depuratum	Apis mellifica . . . . .	Hymenoptera .	Universally domesticated
Menthæ piperitæ Oleum	Mentha piperita . . . . .	Labiata . . .	{Britain and the United States, etc. . . . .}
Menthæ viridis Oleum	Mentha viridis . . . . .	" . . . . .	Britain and Germany, etc. . . .
Menthol . . . . .	{Mentha arvensis vars. piperascens and glabrata . . . . . Mentha piperita . . . . .}	" . . . . .	{China and Japan . . . . . United States . . . . .}
Mezerei Cortex	Daphne {Mezereum . . . . . Laureola . . . . . Gnidium . . . . .}	Thymelæaceæ .	Mountainous parts of Europe
Morphinæ Acetas.	Opium . . . . .	see Opium . . .	see Opium . . . . .
Morphinæ Hydrochloridum . . . . .	" . . . . .	" . . . . .	" . . . . .
Morphinæ Tartras	" . . . . .	" . . . . .	" . . . . .
Morrhua Oleum . . . . .	Gadus <i>Morrhua</i> . . . . .	Teleostei . . .	{Coasts of Norway, France, and England, Newfoundland and Labrador
Moschus . . . . .	Moschus moschiferus . . . . .	Ungulata . . .	{Native of Central Asia; imported from China. . . . .}
Mylabris . . . . .	{Mylabris phalerata and M. Cichorii . . . . .}	Coleoptera . . .	{China, Southern Europe, and India . . . . .}
Myristica . . . . .	Myristica fragrans . . . . .	Myristicaceæ .	{Banda Islands of the Malayan Archipelago and Sumatra, etc. . . . .}
Myristicæ Oleum.	" . . . . .	" . . . . .	" . . . . .
Myrobalanum . . . . .	Terminalia Chebula . . . . .	Combretaceæ .	East Indies . . . . .
Myrrha . . . . .	{Balsamodendron Myrrha and probably other species . . . . .}	Burseraceæ . .	Somaliland and Arabia Felix
Nux Vomica . . . . .	Strychnos Nux-vomica . . . . .	Loganiaceæ . .	{East Indies, imported from India, Ceylon, and Cochinchina . . . . .}

Parts used.	Preparations into which it enters.
{Fresh outer part of the pericarp of the fruit . . . . .}	{Oleum, Syrupus, Tinctura Limonis; Inf. Aurant. Co.; Inf. Gentian. Co. The Oil is contained in Lin. Pot. Iodid. c. Sapone, Spir. Amm. Aromat., Tinct. Guaiac. Amm. and Tinct. Valer. Amm. . . . .}
{Freshly expressed juice of the ripe fruit . . . . . Dried ripe seeds, entire and reduced to a coarse powder. . . . . Dried flowering herb . . . . .}	Syrupus Limonis, Acidum Citricum. . . . . Linum Contusum, Oleum Lini. . . . . Tinct. Lobeliæ Ætherea. . . . .
Glands from the strobiles. . . . .	Infusum, Tinctura Lupuli, Lupulinum. . . . .
Dried strobiles . . . . .	" . . . . .
The minute yellow spores. . . . .	" . . . . .
Concrete saccharine exudation. . . . . Concrete resinous exudation. . . . . Dried leaves. . . . .	" . . . . .
{Saccharine secretion in honey-comb . . . . . Oil distilled from fresh flowering peppermint . . . . . Oil distilled from fresh flowering spearmint . . . . .}	Mel Boracis; Oxymel; Oxymel Scillæ; Conf. Piper. . . . . Aqua, and Spiritus Menthæ Piperitæ; Pil. Rhei Co.; Tinct. Chloroformi et Morphinæ Comp. . . . . Aqua Menthæ Viridis. . . . .
{A crystalline substance obtained by cooling the oil . . . . .}	Emplastrum Menthol. . . . .
Dried bark . . . . .	Liquor Sarsæ Compositus Concentratus. . . . .
Salt of the alkaloid . . . . .	Liquor Morphinæ Acetatis. . . . .
Salt of the alkaloid . . . . .	{Liquor Morph. Hydrochlor., Suppos. Morph., Tinct. Chlorof. et Morph. Comp., Trochiscus Morph., also et Ipecac. . . . . Injectio Morphinæ Hypodermica, and Liquor Morphinæ Tartratis. . . . .}
Oil of fresh liver of the cod. . . . .	" . . . . .
{Dried secretion from the preputial follicles. . . . .}	" . . . . .
The dried beetle. . . . .	{Acetum Mylabridis, Emplastrum Mylabridis, Emplastrum Calefaciens Mylabridis, Liquor Epispasticus Mylabridis, Unguentum Mylabridis. . . . . Oleum, and Oleum Myristicæ Expressum; Pulv. Catechu Co.; Pulv. Cretæ Aromat.; Sp. Armoraciæ Co.; Tinct. Lavand. Co. . . . . Pil. Aloes Socot., Spir. Ammon. Arom., Spir. Myristicæ; Tinct. Guaiac. Amm. and Tinct. Valer. Amm.; of the Spirit, Mist. Ferri Comp. . . . . Unguentum Myrobalani, Unguentum Myrobalani c. Opio. . . . .}
Dried seed divested of its testa . . . . .	" . . . . .
Oil distilled from seed . . . . .	" . . . . .
Dried immature fruits . . . . .	" . . . . .
Gum-resin (from the stem) . . . . .	{Tinct. Myrrh.; Pil. Aloes et Myrrh.; Decoct. Aloes Co.; Mist. Ferri Co.; Pil. Galbani Co.; Pil. Rhei Co. . . . .}
Dried ripe seeds. . . . .	{Extractum, Extractum Liquidum, and Tinctura Nucis Vomice; Strychnina. . . . .}

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Olivæ Oleum . . .	Olea Europæa . . .	Oleaceæ . . .	South of Europe . . .
Oliveri Cortex . . .	Cinnamomum Oliveri . . .	Lauraceæ . . .	{Australia, India, Burma, North America, and Southern United States . . .}
Opium . . . . .	Papaver somniferum . . .	Papaveraceæ . . .	Asia Minor (Smyrna, etc.) . . .
Papaveris Capsulæ . . .	" . . . . .	" . . . . .	Britain and Asia Minor . . .
Pareiræ Radix . . . . .	{Chondrodendron tomentosum . . . . .}	Menispermaceæ . . .	Brazil . . . . .
Pepsinum . . . . .	{Sus Scrofa . . . . . Ovis Aries . . . . . Bos Taurus . . . . .}	Ungulata . . . . .	Domesticated everywhere . . .
Physostigmatis Semen . . . . .	Physostigma venenosum . . .	Leguminosæ . . . . .	Western Africa . . . . .
Picrorhiza . . . . .	Picrorhiza 'Kurroa' . . . . .	Scrophulariaceæ . . .	Northern India . . . . .
Picrotoxinum . . . . .	Anamirta paniculata . . . . .	Menispermaceæ . . .	{Eastern India and the Malayan Islands . . . . .}
Pimenta . . . . .	Pimenta officinalis . . . . .	Myrtaceæ . . . . .	Jamaica . . . . .
Pimentæ Oleum . . . . .	" . . . . .	" . . . . .	{West Indies, Mexico, and Jamaica, etc. . . . .}
Pini Oleum . . . . .	Pinus pumilio . . . . .	Conifereæ . . . . .	Mountains of Central Europe . . .
Piper Nigrum . . . . .	Piper nigrum . . . . .	Piperaceæ . . . . .	East Indies . . . . .
Pix Burgundica . . . . .	Picea excelsa . . . . .	Conifereæ . . . . .	Germany . . . . .
Pix Liquida . . . . .	{Pinus sylvestris and other species . . . . .}	" . . . . .	{Scotland, Russia, Denmark, and Norway . . . . .}
Podophylli Indici Rhizoma . . . . .	Podophyllum Emodi . . . . .	Berberidaceæ . . . . .	Indigenous to the Himalayas . . .
Podophylli Rhizoma . . . . .	Podophyllum peltatum . . . . .	" . . . . .	North America . . . . .
Pruni Virginianæ Cortex . . . . .	Prunus serotina . . . . .	" . . . . .	" . . . . .
Prunum . . . . .	Prunus domestica . . . . .	Rosaceæ . . . . .	South of France . . . . .
Pterocarpæ Lignum . . . . .	Pterocarpus santalinus . . . . .	Leguminosæ . . . . .	Madras and Ceylon . . . . .
Pyrethri Flores . . . . .	{Pyrethrum cinerariæ- folium, P. roseum and P. carneum . . . . .}	Compositæ . . . . .	{California, Dalmatia, and the Caucasus . . . . .}
Pyrethri Radix . . . . .	Anacyclus Pyrethrum . . . . .	" . . . . .	Algeria . . . . .
Quassia Lignum . . . . .	Picraea excelsa . . . . .	Simarubaceæ . . . . .	Jamaica . . . . .
Quillaia Cortex . . . . .	Quillaia 'Saponaria' . . . . .	Rosaceæ . . . . .	Chili and Peru . . . . .
Quinina Hydrochlor . . . . .	{Various species of Cin- chona and Remijia . . . . .}	Rubiaceæ . . . . .	{South America, India, Ceylon, and Java . . . . .}
Quinina Sulphas . . . . .	" . . . . .	" . . . . .	" . . . . .
Resina . . . . .	Various species of Pinus . . . . .	Conifereæ . . . . .	North America . . . . .
Rhamni Purshiani Cortex . . . . .	See Cascara Sagrada . . . . .	" . . . . .	" . . . . .
Rhei Radix . . . . .	{Rheum palmatum, Rheum officinale, and probably other species . . . . .}	Polygonaceæ . . . . .	China and Tibet . . . . .
Rhœados Petala . . . . .	Papaver Rhœas . . . . .	Papaveraceæ . . . . .	Britain . . . . .

Parts used.	Preparations into which it enters.
{Expressed oil from the ripe fruit . . . . .}	{Emplastra Ammoniac. c. Hydrarg., Hydrarg. Picis, Plumbi, Linimenta Ammoniac, Calcis, and Camphoræ; Sapo Durus and Mollis; Unguenta Capsici, Hydrarg. Comp., Hydrarg. Nitratis, and Resinæ.
Dried bark . . . . .	Tinctura Oliveri Corticis.
{Inspissated juice from unripe capsules . . . . . Nearly ripe dried fruits.	Preparations many. Vide Opium.
Dried root . . . . .	Extractum Pareiræ Liquidum.
{An enzyme from the mucous lining of the fresh and healthy stomach of the pig, sheep or calf . . . . .}	Glycerinum Pepsinæ.
Ripe seeds . . . . .	{Extractum Physostigmatis; Physostigminæ Sulphas, from which is prepared Lamellæ Physostigminæ.
Dried rhizome . . . . .	Extractum Picrorhizæ Liquidum, Tinctura Picrorhizæ.
{A neutral principle from the fruits.	
Dried full-grown unripe fruit.	Aqua, Oleum Pimentæ.
Volatile oil from unripe fruit.	
{Oil distilled from fresh leaves and shoots.	
Dried unripe fruit . . . . .	Confectio Piperis, Pulv. Opii Co.
{Resinous exudation from the stem . . . . .}	Emplastrum Picis.
{Bituminous liquid obtained from the wood . . . . .}	Unguentum Picis Liquidæ.
Dried rhizome and roots . . . . .	{Resina Podophylli Indici, and from it Tinctura Podophylli Indici.
Dried rhizome and roots . . . . .	Resina Podophylli, and from it Tinctura Podophylli.
The bark . . . . .	Syrupus and Tinctura Pruni Virginianæ.
Dried ripe fruits . . . . .	Confectio Sennæ.
Heart-wood . . . . .	Tinctura Lavandulæ Composita.
Flower-heads in powder.	
Dried root . . . . .	Tinctura Pyrethri.
Wood of the trunk and branches . . . . .	Infusum, Liqueur Concentratus and Tinctura Quassia.
The inner part of the bark . . . . .	Tinctura Quillaia and Liqueur Picis Carbonis.
Salt of the alkaloid . . . . .	Tinctura Quinina and Vinum Quinina.
Salt of the alkaloid . . . . .	{Ferri et Quinina Citras, Tinct. Quinina Ammon., Pil. Quin. Sulphat., Syrup. Ferri Phosph. c. Quini. et Strych.
{The residue left after the dis- tillation of the Oil of Tur- pentine from the crude oleo- resin . . . . .}	{Emplastrum and Unguentum Resinæ. Emplastra Calefaciens, Cantharidis, Menthol, Picis, Plumbi Iodidi, and Saponis.
{The erect rhizome, deprived of more or less of its cortex and dried . . . . .}	Extractum Rhei, Infusum Rhei, Liqueur Rhei Concentratus, Pilula Rhei Co., Pulvis Rhei Co., Syrupus Rhei, Tinctura Rhei Comp.
Fresh petals . . . . .	Syrupus Rhœados.

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Ricini Oleum . . .	Ricinus communis . . .	Euphorbiaceæ . . .	India . . . . .
Rosæ Gallicæ Petala	Rosa Gallica . . . . .	Rosaceæ . . . . .	Britain and France, cultivated
Rosæ Oleum . . . .	Rosa Damascena . . . . .	" . . . . .	Bulgaria, etc. . . . .
Rosmarini Oleum.	Rosmarinus officinalis . . . . .	Labiatae . . . . .	South of Europe, cultivated in England . . . . .
Rutæ Oleum . . . .	Ruta graveolens . . . . .	Rutaceæ . . . . .	England and Southern Europe . . . . .
Sabine Cacumina	Juniperus Sabina . . . . .	Conifereæ . . . . .	Britain, Southern Europe, etc. . . . .
Saccharum Purificatum . . . . .	Saccharum officinarum . . . . .	Graminaceæ . . . . .	West Indies and British Guiana . . . . .
Saccharum Lactis	Bos Taurus . . . . .	Ungulata . . . . .	Domesticated everywhere . . . . .
Salicinum . . . . .	{From various species of Salix, chiefly Salix fragilis, and Populus.}	Salicaceæ . . . . .	{Temperate regions of the Northern Hemisphere}
Sambuci Flores . . . .	Sambucus nigra . . . . .	Caprifoliaceæ . . . . .	{Britain, Central and Southern Europe . . . . .}
Santali Oleum . . . .	Santalum album . . . . .	Santalaceæ . . . . .	India . . . . .
Santoninum . . . . .	{Artemisia maritima, var. Stechmanniana .}	Compositæ . . . . .	Asiatic Russia . . . . .
Sappan . . . . .	Cesalpinia Sappan . . . . .	Leguminosæ . . . . .	{Madras and Malayan Peninsulas . . . . .}
Sarsæ Radix . . . . .	Smilax ornata . . . . .	Liliaceæ . . . . .	{Costa Rica and South America . . . . .}
Sassafras Radix . . . .	Sassafras officinale . . . . .	Lauraceæ . . . . .	North America . . . . .
Scammonie Radix	{Convolvulus Scammonia . . . . .}	Convolvulaceæ . . . . .	Syria and Asia Minor . . . . .
Scammonie Resina	" " . . . . .	" " . . . . .	" " " . . . . .
Scammonium . . . . .	" " . . . . .	" " . . . . .	{Chiefly from Smyrna, in Asia Minor . . . . .}
Scilla . . . . .	Urginea Scilla . . . . .	Liliaceæ . . . . .	Mediterranean coasts . . . . .
Scoparii Cacumina	Cytisus Scoparius . . . . .	Leguminosæ . . . . .	Britain . . . . .
Senegæ Radix . . . . .	Polygala Senega . . . . .	Polygalaceæ . . . . .	North America . . . . .
Senna Alexandrina	Cassia acutifolia . . . . .	Leguminosæ . . . . .	{Soudan, imported from Alexandria . . . . .}
Senna Indica . . . . .	Cassia angustifolia . . . . .	" . . . . .	Southern India and Arabia . . . . .
Serpentariæ Rhizoma . . . . .	{Aristolochia Serpentaria or Aristolochia reticulata . . . . .}	Aristolochiaceæ . . . . .	{Southern parts of North America . . . . .}
Sesami Oleum . . . . .	Sesamum Indicum . . . . .	Pedaliaceæ . . . . .	{Indigenous to India, cultivated in Turkey and Greece}
Sevum Preparatum	Ovis Aries . . . . .	Ungulata . . . . .	Domesticated everywhere . . . . .
Sinapis . . . . .	Brassica {alba . . . . . {nigra . . . . .}	Crucifereæ . . . . .	Britain . . . . .
Staphisagriæ Semina . . . . .	Delphinium Staphisagria . . . . .	Ranunculaceæ . . . . .	South of Europe . . . . .
Stramonii Folia . . . . .	Datura Stramonium . . . . .	Solanaceæ . . . . .	Britain, cultivated . . . . .
Stramonii Semina	" " . . . . .	" " . . . . .	" " . . . . .
Strophanthi Semina	Strophanthus "Kombé" . . . . .	Apocynaceæ . . . . .	East Africa . . . . .
Strychnina . . . . .	{Strychnos Nuxvomica and other species of Strychnos . . . . .}	Loganiaceæ . . . . .	East Indies and Philippines . . . . .

Parts used.	Preparations into which it enters.
Oil expressed from the seeds . . . . .	{Collodium Flexile, Lin. Sinapis, Mistura Olei Ricini, Pil. Hydrarg. Subchlor. Co.
{Fresh and dried unexpanded petals . . . . .}	{Confectio Rosæ Gallicæ, Syrupus Rosæ, Infusum Rosæ Acidum. Of the Confection, Pilulæ Aloes Barb., Aloes et Asafetidæ, Aloes Socot., and Hydrarg.
Oil distilled from fresh flowers . . . . .	Unguentum Aqua Rosæ.
{Oil distilled from the flowering tops . . . . .}	{Spiritus Rosmarini; Lin. Saponis and Tinct. Lavand. Comp.
{Volatile oil distilled from fresh herb. . . . .}	
Fresh and dried tops. . . . .	
Juice of the sugar-cane. . . . .	All Syrups and Lozenges, and several other preparations.
Whey of milk, evaporated . . . . .	{Pulv. Elaterini Comp., Extract. Bellad. Alcohol., Nucis Vom., Physostigmatis, and Strophanthi.
{Crystalline glucoside obtained from the bark. . . . .}	
Flowers separated from stalks . . . . .	Aqua Sambuci.
Oil distilled from the wood. . . . .	
A crystalline principle . . . . .	Trochiscus Santonini.
Heart-wood . . . . .	Decoctum Sappan.
Dried root . . . . .	{Extractum Sarsæ Liquidum, and Liquor Sarsæ Compositus Concentratus.
Dried root . . . . .	Liquor Sarsæ Compositus Concentratus.
Dried root . . . . .	Resina Scammonie.
Resin from root . . . . .	{Pil. Scam. Co.; Pulv. Scam. Co.; Ext. Col. Co.; Pil. Col. Co.; Pil. Col. et Hyoscyam.
{Gum-resin obtained from living root. . . . .}	
Sliced and dried bulb . . . . .	{Acetum, Oxymel, Pilula Co., Syrupus and Tinctura Scillæ; Pil. Ipecac. c. Scillâ.
Fresh and dried tops . . . . .	Infusum Scoparii (from fresh); Succus Scoparii (from fresh).
Dried root . . . . .	Infusum and Tinctura Senegæ, Liquor Senegæ Concentratus.
Dried leaflets. . . . .	{Confectio, Infusum, Mistura Co., Syrupus, and Tinctura Sennæ; Pulvis Glycyrrhizæ Compositus, Liquor Sennæ Concentratus.
Dried leaflets. . . . .	May be used in the place of Alexandrian Senna.
Dried rhizome and roots . . . . .	{Infusum, Liquor Serpentariæ Concentratus, and Tinctura Serpentariæ; Tinct. Cinchon. Co.
Oil expressed from seeds. . . . .	
Internal fat of the abdomen . . . . .	Unguentum Hydrargyri.
{Dried ripe seeds of both, powdered and mixed . . . . .}	{Charta Sinapis, Oleum Sinapis Volatile (from Black Mustard only); Lin. Sinapis from the Oil.
Dried ripe seeds . . . . .	Unguentum Staphisagriæ.
Dried leaves . . . . .	Tinctura Stramonii.
Dried ripe seeds. . . . .	Extractum Stramonii.
Dried ripe seeds freed from awns . . . . .	Tinctura Strophanthi and Extractum Strophanthi.
The alkaloid. . . . .	

B.P. Name.	Obtained from.	Natural Order.	Geographical Source.
Styrax preparatus	Liquidambar orientalis	Hamamelidaceæ	South-west of Asia Minor
Succinum . . .	Pinites succinifer . . .	Coniferæ . . .	Shores of the Baltic . . .
Sumbul Radix . . .	Ferula 'Sumbul' . . .	Umbelliferae . . .	Asiatic Russia . . .
Tabaci Folia . . .	Nicotiana <i>Tabacum</i> . . .	Solanaceæ . . .	America . . .
Tamarindus . . .	Tamarindus Indica . . .	Leguminosæ . . .	West Indies . . .
Taraxaci Radix . . .	Taraxacum officinale . . .	Compositæ . . .	Britain . . .
Terebinthina Canadensis . . .	Abies balsamea . . .	Coniferæ . . .	Canada . . .
Terebinthinae Oleum . . .	{ Pinus Tæda, palustris, and other species . . . }	" . . .	United States of America . . .
Theobromatis Oleum . . .	Theobroma <i>Cacao</i> . . .	Sterculiaceæ . . .	Central America . . .
Thus Americanum	Pinus { Tæda . . . palustris . . . }	Coniferæ . . .	{ Southern States of North America . . . }
Thymol . . .	{ Thymus vulgaris . . . Monarda punctata . . . Carum Copticum . . . }	{ Labiatae . . . Umbelliferae . . . }	Manufactured in Britain . . .
Thyroideum Siccum . . .	Ovis Aries . . .	Ungulata . . .	. . . . .
Tinospora . . .	Tinospora cordifolia . . .	Menispermaceæ . . .	Tropical India . . .
Toddalia . . .	Toddalia aculeata . . .	Rutaceæ . . .	Madras Peninsula . . .
Tragacantha . . .	Astragalus gummifer . . .	Leguminosæ . . .	Asia Minor and Persia . . .
Turpethum . . .	Ipomea Turpethum . . .	Convolvulaceæ . . .	India and Ceylon . . .
Tylophora Folia . . .	Tylophora asthmatica . . .	Asclepiadaceæ . . .	{ Bengal, Madras Peninsula and other parts of India, and Ceylon . . . }
Ulmæ Cortex . . .	Ulmus campestris . . .	Urticaceæ . . .	{ Central and Southern Europe, and England . . . }
Urginea . . .	{ Urginea Indica and Scilla Indica . . . }	Liliaceæ . . .	India . . .
Uvæ Ursi Folia . . .	Arctostaphylos Uva-ursi . . .	Ericaceæ . . .	Britain . . .
Valerianæ Indicæ Rhizoma . . .	Valeriana Wallichii . . .	Valerianaceæ . . .	Himalayas . . .
Valerianæ Rhizoma . . .	Valeriana officinalis . . .	" . . .	Britain and Europe . . .
Vanilla . . .	Vanilla planifolia . . .	Orchidaceæ . . .	Mexico and Bourbon . . .
Veratri Viridis Rhizoma . . .	Veratrum viride . . .	Liliaceæ . . .	United States and Canada . . .
Veratrina . . .	Schœnocaulon officinale . . .	" . . .	Mexico . . .
Viburnum . . .	Viburnum prunifolium . . .	Caprifoliaceæ . . .	{ Middle and Southern United States . . . }
Zingiber . . .	Zingiber officinale . . .	Scitamineæ . . .	{ West Indies, India, and Cochinchina . . . }

In these Tables, Bentham and Hooker's *Genera Plantarum* has been followed in the names of the botanical sources of the drugs have been made, with the view of name is derived from the vernacular name; italics indicate that the name has been derived from a proper name, or from an old popular or commercial name, recommended by Engler, and in the few names without this termination that have been given in parentheses in the first instance where such natural orders occur.

Parts used.	Preparations into which it enters.
Balsam from the trunk purified	Tinctura Benzoini Composita.
Fossil resinous exudation.	
{ Dried transverse slices of the root . . . . . }	Tinctura Sumbul.
Dried leaves . . . . .	
{ Fruits freed from the brittle outer part of the pericarp . . . }	Confectio Sennæ.
Fresh and dried roots . . . . .	Extractum, Extractum Liquidum, Succus Taraxaci.
The oleo-resin . . . . .	Collodium Flexile.
Oil distilled from turpentine . . . . .	Linimentum, Linimentum Aceticum, and Terebenum.
{ Concrete oil from warm crushed seeds . . . . . }	All Suppositories except Glycerin.
Concrete oleo-resin . . . . .	Emplastrum Picis.
{ A crystalline substance from the oils . . . . . }	
{ Fresh and healthy thyroid gland . . . . . }	Liquor Thyroidei.
Dried stem . . . . .	{ Infusum Tinosporæ, Liquor Tinosporæ Concentratus, Tinctura Tinosporæ.
Dried root-bark . . . . .	Infusum Toddaliæ, Liquor Toddaliæ Concentratus.
Gummy exudation . . . . .	Mucilago, Glycerinum, Pulv. Tragac. Co.; Conf. Sulphuris; Pulv. Opii Co.; Mist. Cretæ and Guaiaci; Pil. Quinin. Sulph.; of the Mucilage, Lotio Hydrarg. Nigra.
Dried root and stem . . . . .	
Dried leaves . . . . .	
Dried inner bark . . . . .	
Younger bulbs . . . . .	{ Pilulæ Ipecacuanhæ c. Urginea, Acetum Urginæ, Oxymel Urginæ, Pilula Urginæ Composita, Syrupus Urginæ, and Tinctura Urginæ.
Dried leaves . . . . .	Infusum Uvæ Ursi.
Dried rhizome and rootlets . . . . .	Tinctura Valerianæ Indicæ Ammoniata.
Dried erect rhizome and roots . . . . .	Tinctura Valerianæ Ammoniata.
Dried fruit . . . . .	
Dried rhizome and rootlets . . . . .	
The alkaloid from cevadilla . . . . .	Unguentum Veratrinæ.
Dried bark . . . . .	Extractum Viburni Prunifolii Liquidum.
Scraped and dried rhizome . . . . .	Syrupus and Tinctura, Zingiberis. It is also used in some powders and other preparations.

with respect to botanical classification, and some alterations in the method of print-making the Table more useful to students, thus: quotation marks imply that the name has been used as a generic name; and an initial capital letter that the name has been retained, the name recommended by Lindley, for the sake of uniformity, has